

## Rabbit dietary supplementation with pale purple coneflower. 2. Effects on the performances, bacterial community, blood parameters and immunity of growing rabbits

A. Kovitvadhi<sup>1†</sup>, F. Gai<sup>2</sup>, S. Dabbou<sup>1</sup>, I. Ferrocino<sup>1</sup>, L. Rotolo<sup>1</sup>, M. Falzone<sup>3</sup>, C. Vignolini<sup>3</sup>, M. S. Gennero<sup>4</sup>, S. Bergagna<sup>4</sup>, Daniela Dezzutto<sup>4</sup>, R. Barbero<sup>4</sup>, P. Nebbia<sup>3</sup>, S. Rosati<sup>3</sup>, L. Cocolin<sup>1</sup>, I. Zoccarato<sup>1</sup> and L. Gasco<sup>1,2</sup>

<sup>1</sup>Department of Agricultural, Forest and Food Sciences, University of Turin, Largo P. Braccini 2, 10095 Grugliasco, Turin, Italy; <sup>2</sup>Institute of Science of Food Production, National Research Council, Largo P. Braccini 2, 10095 Grugliasco, Turin, Italy; <sup>3</sup>Department of Veterinary Sciences, University of Turin, Largo P. Braccini 2, 10095 Grugliasco, Turin, Italy; <sup>4</sup>Veterinary Medical Research Institute for Piemonte, Liguria and the Valle D'Aosta, via Bologna 148, 10154 Turin, Italy

(Received 11 June 2015; Accepted 3 December 2015; First published online 14 January 2016)

*Echinacea pallida* (EPAL), a herbaceous flowering plant with immunomodulatory properties, has been chosen to determine the pre- and post-supplementary effects on the growth performances, bacterial community, blood parameters and immunity of growing rabbits. The same Grimaud does (14-week-old) from the studied in the first part of this study were randomly divided into two groups (n = 50/group). The first group was fed a basal diet without supplementation (Control group, C) while the another group was fed a basal diet supplemented with 3 g EPAL/kg diet (Echinacea group, E). From the second parturition, 80 weaned kits (40 from the C does and 40 from the E does) were randomly assigned to four groups of 20 animals each and were fed a growing commercial diet supplemented with or without a 3 g EPAL/kg diet: the CC group (rabbits from the C does fed the control diet), CE group (rabbits from the C does fed the supplemented diet), EC (rabbits from the E does fed the control diet) and EE group (rabbits from the E does fed the supplemented diet). The dietary EPAL treatment did not affect the growth performance. Ten fattening rabbits from each group were selected to evaluate the bacterial community and blood parameters, while the remaining rabbits (n = 10/group) were used to study phagocytosis and the humoral immune response. The variability was evaluated from hard faeces at 35, 49 and 89 days, and the caecal content at 89 days. The variability of the bacterial community of the EE group was higher than that of the other groups. The phagocytic activity was higher in the CE and EE groups than in the CC and EC ones (30.9 and 29.7 v. 21.2 and 21.8%;  $P < 0.05$ ), whereas no statistically significant difference was observed for the blood parameters or humoral immune response against vaccination (rabbit haemorrhagic disease virus) at 95 days old which the serum was collected at 88, 102, 109, 116 and 123 days old. In conclusion, no impact of EPAL dietary supplementation has been observed on the growth performances, bacterial community, blood parameters or humoral immune responses in growing rabbits, except for an increase in phagocytic activities.

**Keywords:** bacterial community, blood parameter, *Echinacea pallida*, immunity, rabbit

### Implications

Dietary supplementation with an immunomodulatory herb, such as pale purple coneflower (*Echinacea pallida*: EPAL), could provide benefits to growing rabbits. Phagocytic activity, which is a defense mechanism against pathogens, was increased in the fattening rabbits fed the supplemented diets. However, dietary supplementation with EPAL had no impact on the growth performances, bacterial community, blood parameters or humoral immune responses in the growing rabbits.

### Introduction

The addition of medicinal plants with immune-stimulating properties, such as *Echinacea* species, is a valuable strategy as pointed out in the previous part of this study (Dabbou *et al.*, 2016). An improvement in growth performances and a reduction in the mortality rate were observed in growing rabbits fed diets with *Echinacea purpurea* (EPUR) supplementation (Ahmed *et al.*, 2008; Arafa *et al.*, 2010). The higher number of white blood cells has been reported in groups of rabbits, laying hens and fattening pigs treated with EPUR (Böhmer *et al.*, 2009; Arafa *et al.*, 2010). EPUR dietary

<sup>†</sup> E-mail: attawit.kovitvadhi@hotmail.com

addition has been shown to enhance phagocytic activity in fattening pigs (Böhmer *et al.*, 2009) and the level of specific antibody response against vaccination in poultry, quail, swine and rabbit (Maass *et al.*, 2005; Böhmer *et al.*, 2009; Ma *et al.*, 2009). To date, the research on animal studies has tended to focus on EPUR rather than *Echinacea angustifolia* (EANG) or *Echinacea pallida* (EPAL; Hermann *et al.*, 2003; Maass *et al.*, 2005; Böhmer *et al.*, 2009; Ma *et al.*, 2009; Arafa *et al.*, 2010; Sahin *et al.*, 2012) and it has been shown that there is a broad variation of the active compounds in the *Echinacea* species (Dabbou *et al.*, 2016). Therefore, the objective of this present study has been to evaluate pre- and postnatal effects on the performances, bacterial community, blood parameters and immunity of rabbits after the introduction of EPAL (pale purple coneflower) as a dietary addition. The consequences of EPAL dietary supplementation on rabbit does were presented in the previous part of this study (Dabbou *et al.*, 2016).

## Material and methods

### *Animals, housing, diets, experimental design and growth performances*

The trial was conducted at the experimental rabbit facility of the Department of Agricultural, Forest and Food Sciences in Carmagnola (Turin, Italy). Eighty kittens were taken from the second parturition of control group (C) and *Echinacea* group (E) of the previous study (Dabbou *et al.*, 2016). The kittens were weaned at 35-day old, were randomly separated into four groups of 20 and were housed in individual wire cages (0.41 m long × 0.30 m wide × 0.28 m high) when their mothers were 194-day old. The fattening rabbits were fed a growing commercial basal diet with or without the supplementation (3 g/kg of EPAL): the CC group (rabbits from the C does fed the control diet), CE group (rabbits from the C does fed the supplemented diet), EC (rabbits from the E does fed the control diet) and EE group (rabbits from the E does fed the supplemented diet). Feeds and clean water were provided *ad libitum*. The preparation and chemical analysis of the feeds and supplement (EPAL powder), which were obtained from the same provider, were performed as outlined in the previous part of this study (Dabbou *et al.*, 2016). The diet composition for the growing rabbits is illustrated in Table 1. Throughout the whole trial, the facility was climate and light controlled in order to maintain a temperature of  $22 \pm 2^\circ\text{C}$  and a 16L:8D photoperiod. Live weight and feed intake were measured at 35, 49 and 77 days old to calculate the average daily weight gain, average daily feed intake and feed conversion ratio, whereas morbidity and mortality rate were recorded according to Gidenne *et al.* (2009). Ten fattening rabbits were selected from each group to evaluate the bacterial community and blood parameters, while the remaining ten rabbits from each group were kept until 123 days old to study the phagocytic activity and specific antibody response. The timeline of the experimentation is illustrated in Figure 1.

**Table 1** Composition of the growing rabbit diets

	Diets	
	Control <sup>1</sup>	Treatment <sup>1</sup>
Chemical composition <sup>2</sup>		
Dry matter (DM)	89.8	89.8
CP (% DM)	17.1	17.3
Ether extract (% DM)	3	3
NDF (% DM)	39.4	39.6
ADF (% DM)	23.7	24
ADL (% DM)	6.6	6.6
Ash (% DM)	9.7	10.4
Starch (% DM)	12	12.3
Dry ground <i>Echinacea pallida</i> (g/kg)	0	3

<sup>1</sup>Minerals and vitamins in the growing diets: calcium 1% of DM, phosphorus 0.4% of DM, sodium 0.3% of DM, lysine 0.7% of DM, methionine 0.4% of DM, vitamin A 12 500 IU/kg, vitamin D3 1250 IU/kg, vitamin E 100 mg/kg, ferrous carbonate 704 mg/kg, copper sulphate 98 mg/kg, manganese oxide 209 mg/kg, zinc oxide 186 mg/kg, sodium selenite 0.57 mg/kg and potassium iodide 2.5 mg/kg. These data were provided by Ferrero Mangimi S.p.A. (Farigliano CN, Italy), who formulated and prepared the experimental diets.


<sup>2</sup>The experimental diets were analysed in the laboratory of the Department of Agricultural, Forest and Food Sciences, Turin, Italy.

### *Bacterial community*

The hard faeces of the growing rabbits were collected at 35, 49 and 89 days from five animals per group, whereas the caecal content samples ( $n = 5$  per group) were taken immediately after slaughtering at 89 days. The samples were kept in sterile polyethylene bags and immediately stored at  $-20^\circ\text{C}$  until analysis. The samples (10 g) were homogenized in 90 ml of Ringer's solution (Oxoid, Milan, Italy) for 2 min in a stomacher (LAB Blender 400 and Sto-circul-bag stomacher bags, PBI, Milan, Italy) at room temperature; a deposit was allowed to set for 1 min, and 1 ml of the supernatant was used for the DNA extraction. A Powersoil DNA kit (MO-BIO, Carlsbad, CA, USA) was used, according to the manufacturer's instructions. A total of 5  $\mu\text{l}$  of RNase (Promega, Milan, Italy) was added to the DNA and incubated at  $37^\circ\text{C}$  for 30 min before storage at  $-20^\circ\text{C}$ . DNA was quantified using a NanoDrop 1000 spectrophotometer (Thermo Scientific, MA, USA) and was standardized at 50 ng/ $\mu\text{l}$ . PCR was performed with the universal primers 338F-GC and 518R (Muyzer *et al.*, 1993; Invitrogen, Carlsbad, CA, USA) amplifying the variable V3 region of the 16S rRNA gene. The amplicons obtained (about 250 bp) were analysed by means of DGGE, with a Dcode universal mutation detection system (BioRad, Milan, Italy). The gels were run for 4 h at 200 V, stained with SYBR<sup>®</sup> Gold Nucleic Acid Gel Stain (Life technologies, Milan, Italy) for 30 min, and analysed under UV using UVipro Platinum 1.1 Gel Software (Eppendorf s.r.l., Milan, Italy).

### *Blood parameters*

Blood samples were taken from 10 rabbits per group at 89 days old during the slaughter process. Sterile tubes containing ethylenediaminetetraacetic acid were used to collect the samples for haematological analysis, which was conducted using an automated laser cell counter calibrated for rabbits



**Figure 1** The timeline of the experimental plan.

(MS4-S Hematology Analyzer, Melet Schloesing, Osny, France). A plain serum tube was instead used as a container to obtain serum for biochemistry analysis and protein electrophoresis, which was conducted using an automated photometer system (Screen Master Touch, Hospitex Diagnostics, Sesto Fiorentino, Italy) and a semi-automated agarose gel electrophoresis system (Sebia Hydrasys, Evry, France), respectively.

Changes in the number and species of white blood cells after infection and blood production disorders can be revealed by means of haematological parameters (Varga, 2014). The liver function (albumin), liver damage (aspartate aminotransferase and alanine aminotransferase), renal function (blood urea nitrogen and creatinine) and lipid metabolism (cholesterol and triglyceride) of the rabbits were established through an evaluation of the serum biochemistry (Varga, 2014). A serum protein fraction was used to introduce the an inflammatory response ( $\alpha$  1 globulin), metabolic enzymes ( $\alpha$  2 globulin), immunoglobulin G ( $\gamma$  globulin), iron-binding glycoprotein ( $\beta$  1 globulin) and immunoglobulin A and M ( $\beta$  2 globulin; Tizard, 2013). Therefore, these parameters were evaluated in this study to illustrate the effects of EPAL supplementation.

#### Phagocytic activity


At 88 days, blood samples were taken from the lateral saphenous vein of 10 rabbits per group. The samples were placed in sterile tubes containing lithium-heparin, kept at 4°C and immediately transported to the laboratory to perform a phagocytosis test, which involved a direct counting procedure using bacterial cells (Ragap *et al.*, 2012). The heparinized blood samples (100  $\mu$ l) were left until they reached room temperature (22°C), and then they were mixed with 10  $\mu$ l of diluted bacterial solution (100  $\mu$ l of broth with  $1 \times 10^{10}$  cells/ml of *Streptococcus canis* and 900  $\mu$ l of PBS, pH 7.2) in 14 ml sterile glass tubes (15  $\times$  125 mm). These mixtures were incubated at 37°C for 30 min. Blood samples, without the addition of bacterial cells, were used as a negative control. Immediately after incubation, all the samples were placed at 4°C to stop the reaction, and blood smearing was performed. After drying, the samples were stained with gram stain, according to the manufacturer's instructions (Sigma-aldrich®, Buchs, Switzerland). The results were evaluated under an oil immersion light microscope.

The neutrophils and monocytes that engulfed the gram positive cocci were counted as phagocytic cells. Phagocytic activity was calculated on the basis of the percentage of phagocytic cells observed (Ragap *et al.*, 2012). Triplicate counting was performed for each sample.

The phagocytosis test is able to confirm the response of sentinel cells (Macrophage and dendritic cells) and proinflammatory cytokine production against the pathogen invasion. After a bacterial pathogen passes the physical barrier and invades the host, sentinel cells can detect pathogen-associated molecular patterns on bacteria antigen through toll-like receptors. The sentinel cells then secrete vasoactive molecules and proinflammatory cytokines (interleukin 1, interleukin 6 and tumor necrotic factor  $\alpha$ ), which activate a systemic response and phagocytic activities (Tizard, 2013).

#### Specific humoral immune response

The rabbits were vaccinated against rabbit haemorrhagic disease virus (RHDV; IZOVAC MEVAX®, IZO S.p.A., Brescia, Italy) at 95 days old, and serum was collected at 88, 102, 109, 116 and 123 days old through centrifugation (1600  $\times$  g for 15 min) of the blood samples (collected from the lateral saphenous vein) after incubation at room temperature (22°C) for 4 h. The serum was stored at -20°C to evaluate the percentage of inhibition (PI) against the RHDV antibody, which was conducted by competitive ELISA (Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia, Brescia, Italy). The wells were coated with the antibody against RHDV. The serum samples (5  $\mu$ l) and RHDV antigen (25  $\mu$ l) were added to each well (1/10 serum dilution). Duplicates of the positive and negative controls were tested for each plate. The protocol recommendation by the Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia (Brescia, Italy) was followed. The colour was developed by adding 50  $\mu$ l of 3,3',5,5'-tetramethylbenzidine substrate, and the reaction was then stopped by blocking solution. An automated ELISA microplate reader was used to determine an optical density at 492 nm (Titertek Multiskan MCC/340, Labsystems, Helsinki, Finland). The results were expressed as PI on the basis of the optical density values according to the following formula:  $PI = [1 - (\text{optical density of the sample}) / (\text{average negative control optical density})] \times 100$ .



**Figure 2** Dendrogram of the PCR-DGGE profiles of the rabbit bacterial community from hard faeces samples (HF) at 35, 49 and 89 days, and caecal content samples (C) at 89 days between CC (rabbits fed basal diets from does fed the control diet), CE (rabbits fed treated diets from does fed the control diets), EC (rabbits fed basal diets from does fed the supplemented diets) and EE group (rabbits fed treated diets from does fed the supplemented diets).

In this study, the RHDV vaccine contained dead virus. Therefore, this vaccination only stimulated the humoral immune response (Tizard, 2013). Dead virus is engulfed, then is represented by major histocompatibility complex (MHC) class II and is presented to helper T cell by antigen presenting

cells which activates B cell differentiation to plasma cells for specific antibody production (Tizard, 2013). Therefore, the stimulation of the humoral immune response through vaccination can be used to confirm phagocytosis process, antigen presenting activity and antibody network.

#### Statistical analysis

All the statistical analyses were performed using the SPSS software package (IBM SPSS, 2012). The growth performances, blood parameters, phagocytic activity and PI were processed with one-way ANOVA (groups as the fixed factors) whose normal distribution was observed in the parametric data by Kolmogorov–Smirnov test. Significant differences in treatment means were established by Duncan's multiple range tests at  $P < 0.05$ . The entire fingerprint database of the bacterial community was created using Bionumerics version 5.1 software (Applied Maths, Sint Martens Latem, Belgium). A dendrogram of similarity was retrieved using the dice coefficient and unweighted pair group method for the arithmetic average clustering algorithm (Figure 2). The symmetrical matrix of the all pairwise distances among the proximities of the DGGE profiles of the rabbit's bacterial community was calculated using the Euclidean distance to establish the distance matrix. Two dimensions were chosen for the ordination. The distance matrix was used for nonmetric multidimensional scaling (nMDS) plots, which were performed in IBM SPSS (2012) using the PROXSCAL scaling algorithm. The nMDS plots of the bacterial communities from hard faeces at 35, 49 and 89 days and the caecal contents at 89 days are presented in Figure 3a–d, respectively. Kruskal's S-Stress was calculated. On nMDS plots, the near point shows higher similarity than the far point (Combes *et al.*, 2011).

## Results


#### Growth performances and bacterial community

No significant differences were found in growth performance (Table 2). There was no morbidity or mortality during the experimental period. The dendrogram and two-dimensional nMDS of the bacterial community of the rabbits were generated using the digitalized PCR-DGGE fingerprints obtaining from the DNA extracted directly from the hard faeces and caecal content samples (Figures 2 and 3, respectively). The cluster analysis (Figure 2) showed a shift in the microbial community, due to the an increase in age. Moreover, it was possible to observe a difference between the hard faeces at 89 day old and the caecal content fingerprints (Figure 2). Higher within-group variability was observed in the kittens from the treated groups, compared to the control (Figure 3a). No clear difference was observed between the CC, CE and EC samples, whereas the variability of the bacterial community of the EE group was higher than the others throughout the experimental period (Figure 3b–d).

#### Blood parameters and immunological test

The blood parameters, phagocytic activities and inhibition percentage of the competitive antibody are illustrated in Table 2.





**Figure 3** nMDS plot of the PCR-DGGE profiles of the rabbit bacterial community between CC (rabbits fed basal diets from does fed the control diets), CE (rabbits fed treated diets from does fed the control diets), EC (rabbits fed basal diets from does fed the supplemented diets) and EE group (rabbits fed treated diets from does fed the supplemented diets) from hard faeces samples (HF) at days 35 (a), 49 (b) and 89 (c), and samples of the caecal content at 89 days old (d).

All the evaluated parameters fell into the normality range (Varga, 2014). No differences were observed for the haematology, serum biochemistry or serum protein electrophoresis between the groups ( $P > 0.05$ ). An increase in phagocytic activity was observed in the rabbits fed the treated diets, compared to the rabbits fed the control diets (CE and EE v. CC and EC;  $P < 0.001$ ). Because of a specific antibody response against the vaccination, the inhibition percentage was lower than 25% 1 week before the vaccination in all the groups which was considered as negative (Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia, Brescia, Italy). A higher level of antibody response was observed 1 week after the vaccination. The inhibition percentage surged to above 80% in the 2nd week after the vaccination, and then became constant after this period. No significant difference was observed for the antibody production against RHDV between the treated groups, which were compared each week.

## Discussion

### Growth performances

The growth performance of the *Echinacea* supplemented groups did not show any significant differences. The present

results differ from those of Arafa *et al.* (2010) who found, in a similar study using EPUR at 130 mg/kg BW, a significant decrease in the mortality rate and an increase in the live weight in 6-week-old growing rabbits fed E diets ( $P < 0.05$ ). Dietary herb supplementation usually leads to an improvement in the flavour, which in turn accounts leads to an increase in feed ingestion and better performance (Franz *et al.*, 2010; Christaki *et al.*, 2012). Ahmed *et al.* (2008) highlighted a significant improvement in the final BW, daily weight gain and feed conversion ratio in growing rabbits, which had orally been given 7.5 mg of EPUR extracts/kg BW/day in liquid form. However, the outcomes of the above reported references are not exactly comparable with the present trial due to some dissimilarities in the experimental designs concerning: the tested *Echinacea* species, concentration of the supplement, administration route (oral by means of a liquid mixture), supplement preparation (extraction) and the supplemented periods in the doe diets. Positive outcomes on productive performance have been reported in rabbits after EPUR addition (Ahmed *et al.*, 2008; Arafa *et al.*, 2010), whereas studies conducted on other livestock species have not found any improvement (Hermann *et al.*, 2003; Maass *et al.*, 2005; Böhmer *et al.*, 2009; Sahin *et al.*, 2012).

**Table 2** Effects of pre- and postnatal dietary phytoadditives (*Echinacea pallida*; EPAL) on growth performances, blood parameters, phagocytic activity and specific antibody response in growing rabbits

Parameters	Groups				SEM	P-value
	CC	CE	EC	EE		
Live weight (g)						
At 35 days	884	889	888	882	5.90	0.97
At 49 days	1713	1711	1744	1716	8.87	0.51
At 77 days	3031	2997	3106	3041	18.1	0.19
Growth performance in 35–49 days						
ADFI (g/day)	134	138	140	139	1.19	0.32
ADG (g/day)	59.2	58.7	61.2	59.6	0.40	0.16
Feed conversion ratio	2.28	2.36	2.29	2.35	0.01	0.20
Growth performance in 49–77 days						
ADFI (g/day)	176	178	181	181	1.48	0.47
ADG (g/day)	45.4	44.4	46.9	45.7	0.45	0.25
Feed conversion ratio	3.87	4.03	3.88	3.98	0.03	0.28
Growth performance in 35–77 days						
ADFI (g/day)	162	164	167	167	1.28	0.36
ADG (g/day)	49.9	49.0	51.5	50.2	0.35	0.08
Feed conversion ratio	3.25	3.37	3.26	3.34	0.02	0.12
Haematology						
Erythrocytes ( $\times 10^{12}/l$ )	6.46	6.39	6.20	6.22	0.08	0.64
Haematocrit (%)	43.7	42.9	42.0	41.8	0.53	0.56
Haemoglobin (g/dl)	13.2	13.1	12.8	13.2	0.20	0.89
MCV (fl)	67.9	67.2	67.8	67.3	0.37	0.89
MCH (pg)	20.4	20.4	20.6	21.2	0.22	0.60
MCHC (g/dl)	30.1	30.4	30.5	31.6	0.30	0.37
Leukocyte ( $\times 10^9/l$ )	7.73	7.54	8.71	7.59	0.58	0.89
Neutrophils (%)	76.5	74.2	78.2	73.6	1.35	0.61
Lymphocytes (%)	15.4	16.7	14.1	15.2	0.93	0.82
Monocytes (%)	6.92	7.83	6.48	7.19	0.46	0.78
Eosinophils (%)	0.48	0.61	0.68	0.51	0.05	0.53
Basophils (%)	0.52	0.60	0.45	0.58	0.05	0.73
Serum biochemistry						
AST (U/l)	16.0	14.4	17.1	19.8	0.95	0.22
ALT (U/l)	42.2	43.6	40.3	43.2	2.39	0.96
BUN (mg/dl)	11.5	9.04	10.6	12.5	0.58	0.17
Creatinine (mg/dl)	1.11	1.12	1.18	1.16	0.02	0.69
Cholesterol (mg/dl)	33.5	29.6	36.5	36.0	1.92	0.62
Triglyceride (mg/dl)	56.8	54.9	61.8	51.0	3.08	0.68
Serum protein electrophoresis <sup>1</sup>						
Total protein (g/dl)	7.89	6.96	6.60	7.05	0.19	0.10
Albumin (g/dl)	5.81	5.25	4.87	5.36	0.14	0.15
Globulin (g/dl) <sup>2</sup>	1.89	1.71	1.54	1.70	0.04	0.06
Albumin/globulin	3.19	3.09	3.26	3.17	0.05	0.79
$\alpha$ 1 globulin (%)	1.40	1.14	0.97	1.13	0.08	0.31
$\alpha$ 2 globulin (%)	1.42	1.49	1.21	1.32	0.09	0.74
$\beta$ 1 globulin (%)	3.00	3.21	3.39	3.15	0.10	0.60
$\beta$ 2 globulin (%)	4.67	5.00	5.47	4.92	0.17	0.46
$\gamma$ globulin (%)	13.7	13.8	12.6	13.5	0.29	0.49
Phagocytic activity (%)	21.2 <sup>A</sup>	30.9 <sup>B</sup>	21.8 <sup>A</sup>	29.7 <sup>B</sup>	0.88	0.01
Percentage of inhibition against competitive antibody <sup>3</sup>						
88 days	10.0	10.9	9.39	8.63	0.86	0.80
102 days	31.0	37.9	29.1	33.1	3.35	0.81
109 days	91.1	90.1	89.4	89.9	0.85	0.92
116 days	93.8	93.9	93.9	93.4	0.27	0.91
123 days	94.1	93.7	94.7	94.3	0.18	0.31

CC = rabbits fed the basal diets from does fed the control diet; CE = rabbits fed diets with EPAL from does fed the control diet; EC = rabbits fed the basal diets from does fed diets with EPAL; EE = rabbits fed diets with EPAL from does fed diets with EPAL; ADFI = average daily feed intake; ADG = average daily weight gain; MCV = mean corpuscular volume; MCH = mean corpuscular haemoglobin; MCHC = mean corpuscular haemoglobin concentration; AST = aspartate aminotransferase; ALT = alanine aminotransferase; BUN = blood urea nitrogen.

<sup>1</sup>Each globulin fraction was expressed as a percentage of the total protein.

<sup>2</sup>The globulin was calculated as the difference between the total protein and albumin.

<sup>3</sup>The specific antibody production was performed by the vaccination against rabbit hemorrhagic disease virus at 95-day-old and the data is presented for each collection day.

### Bacterial community

The bacterial community in rabbits changes continuously from birth to adulthood. This ecosystem involves nutrient utilization, the immune system, development of the digestive tract and pathogen prevention (Combes *et al.*, 2013). The period from birth to 49 days (time window of permissiveness) is a critical period for rabbits, because several microorganisms present in an environment can randomly access the rabbit's digestive tract, depending on the diet formulation, management programme and/or environment conditions (Combes *et al.*, 2013). In the aforementioned study, the variability of the bacterial community in caecum at the neonatal stage was remarkable between each rabbit. Thereafter, this variability decreased continuously with age. Finally, the bacterial diversity was nearly similar between each rabbit (Combes *et al.*, 2011). A similar pattern has been observed in the present study, although hard faeces were used as the samples. The different characteristics between the caecal contents and the hard faeces provided in a dissimilarity of the bacterial community which was also reported by Michelland *et al.* (2010), as shown in Figure 2.

Nutrition is one of the most important factors that can influence the microbial community (Michelland *et al.*, 2011). In the present study, the variability between the kittens from the E does was higher at weaning than the kittens from the C does (Figure 3a). This result may be due to mother and/or diet effects, because the kittens lived in the same cage and were fed the same diet as their mother until weaning. If EPAL contained properties that could modify the bacterial community of does, the initial gut microbial composition of their kittens at weaning would also be changed. The addition of EPAL in the fattening period (CE group) or discontinuous supplementation (EC group) may not be sufficient to provoke effects on the bacterial community. Therefore, in order to modify the bacterial community in fattening rabbits through EPAL supplementation, it should be performed on their mother and the rabbits should be fed the supplemented diet continuously over the fattening period. The variability in the bacterial community between each rabbit of EE is higher than that of the other groups throughout the experimental period (Figure 3). At the moment, it is impossible to conclude whether the modification in EE is a positive or negative outcome. Further analysis on bacterial species should be performed to clarify this aspect.

### Blood parameters and immunological test

The physiological and biochemical status of animals can be expressed through haematology and blood chemistry. The collection should be performed more than 6 weeks after the supplementation, because no statistical difference between groups has been observed in samples that were collected at 3 weeks (Arafa *et al.*, 2010). In the present study, blood was taken from the rabbits after 54 days of supplementation. Therefore, a difference would have been observed, if the supplement had affected these parameters. A higher proliferation rate of spleen lymphocytes in EPAL supplemented mice has been reported in an *in vitro* study, but the

haematology indices were not influenced (Zhai *et al.*, 2007). EPUR supplementation in rabbits, laying hens and fattening pigs has been seen to increase the total amount of white blood cells, while in this study, EPAL in this study did not (Böhmer *et al.*, 2009; Arafa *et al.*, 2010). Therefore, the variation in the chemical compounds between EPUR and EPAL could be a possible explanation.

*Echinacea* genus herbs are particularly well known as immunostimulant agents, and they mainly affect innate immunity (Barnes *et al.*, 2005; Zhai *et al.*, 2007). The purified glycoproteins (arabinogalactans, fructofuranosides and heteroxylans), as well as alkylamides, especially isobutylamides and ketones (only in EPAL), in *Echinacea* plants appear to play important roles on phagocytic stimulation (Classen *et al.*, 2006; Kraus *et al.*, 2007). The supplementation of arabinogalactan-protein from an EPAL extract has been seen to increase the production of IL-6 and nitrite, which are by-products of activated macrophages (Classen *et al.*, 2006). An increase in phagocytic activity has been reported in fattening pigs after EPUR juice addition (Böhmer *et al.*, 2009), and in the present study after EPAL dietary integration. Therefore, the active chemical compounds in EPAL could be the cause of the improvement in phagocytosis.

As far as adaptive immunity is concerned, the EPAL extract significantly increased the production of IL-4 and IL-10 from the type 2 T helper cells and immunoglobulin M involved in humoral immune responses. A greater production of IL-2 and IFN- $\gamma$  from the type 1 T helper cells, which is associated with cell-mediated immunity, was also observed in the *Echinacea* treated group (Classen *et al.*, 2006; Zhai *et al.*, 2007).  $\beta$  2 globulin fractions should increase when immunoglobulin M production is higher. However, differences in the plant preparation before usage (extraction *v.* raw materials) and in the study design (*in vitro v. in vivo*) may alter the outcomes, as observed in the present study. A higher level of total globulin has been reported in rabbits after EPUR supplementation (Arafa *et al.*, 2010). The specific antibody response against inactivated vaccine as a cost effective way of assessing the network functionality of antigen presenting cells (via MHC class II), T helper cell activation and antibody production has been evaluated in this study and has here been shown not to be affected by the chosen treatments. Therefore, the selection of the *Echinacea* species (EPUR *v.* EPAL) and of the experimental design (with or without stress environment) can lead to diverse outcomes.

The alkamide, polysaccharide and proteoglycan in *Echinacea* plants have been suggested as active compounds that can stimulate antibody production (Egert and Beuscher, 1992; Hudson *et al.*, 2005). In an *in vitro* study, it has been shown that EANG extracts contain antiviral properties against the herpes simplex virus (HSV), the influenza virus (FV) and rhinovirus, whereas aqueous extracts of EPUR only show antiviral activities on HSV and FV, but no significant change has been found for EPAL extracts (Hudson *et al.*, 2005). Instead, the antibody response against the red blood cells of sheep by plaque-forming cell assays has not been affected by the supplementation of *Echinacea* extracts

(Zhai *et al.*, 2007). The level of specific antibody response against vaccination in EPUR treated animals has been shown to be greater than in animals fed control diets (Maass *et al.*, 2005; Böhmer *et al.*, 2009; Ma *et al.*, 2009), but some studies have not reported positive outcomes (Hermann *et al.*, 2003). The diversity in the active chemical compounds between *Echinacea* species may be the cause of the contrasting results. Further studies on the pharmacological activities of the different active compounds and on the synergistic effects are necessary to clarify these differences.

## Conclusion

An increase in phagocytosis has been observed in rabbits fed diets with EPAL supplementation, without any impact on growth performances, blood parameters, bacterial community or humoral immune response.

## Acknowledgements

This research was supported by the 'University of Torino (ex 60%)' grant (Es. fin. 2013–2015). The authors would like to thank Dr Claudio Malavasi (Biotrade snc®, Modena, Italy) for the dry ground EPAL samples, Dr Paolo Cerrina (Ferrero S.p.a Cuneo, Italy) for his support with the feed formulation, Mrs Vanda Malfatto for her technical support, and Mr Dario Sola, Mr Paolo Montersino and Mr Mario Colombano for the rabbit care.

## References

Ahmed HS, Kamel KI, El-Sabeiy ME and Zeitouny MH 2008. Effect of *Echinacea* extract supplementation on growth performance and hemo-biochemical traits of growing rabbits. *Egypt Poultry Science* 28, 1165–1180.

Arafa NMS, Salem SM and Farid OAHA 2010. Influence of *Echinacea* extract pre- or postnatal supplementation on immune and oxidative status of growing rabbits. *Italian Journal of Animal Science* 9, 338–343.

Barnes J, Anderson L, Gibbons S and Phillipson J 2005. *Echinacea* species (*Echinacea angustifolia* (DC.) Hell., *Echinacea pallida* (Nutt.) Nutt., *Echinacea purpurea* (L.) Moench): a review of their chemistry, pharmacology and clinical properties. *Journal of Pharmacy and Pharmacology* 57, 929–954.

Böhmer BM, Salisch H, Paulicks BR and Roth FX 2009. *Echinacea purpurea* as a potential immunostimulatory feed additive in laying hens and fattening pigs by intermittent application. *Livestock Science* 122, 81–85.

Christaki E, Bonos E, Giannenas I and Florou-Paneri P 2012. Aromatic plants as a source of bioactive compounds. *Agriculture* 2, 228–243.

Classen B, Thude S, Blaschek W, Wack M and Bodinet C 2006. Immunomodulatory effects of arabinogalactan-proteins from *Baptisia* and *Echinacea*. *Phytomedicine* 13, 688–694.

Combes S, Fortun-Lamothe L, Cauquil L and Gidenne T 2013. Engineering the rabbit digestive ecosystem to improve digestive health and efficacy. *Animal* 7, 1429–1439.

Combes S, Michelland R, Monteils V, Cauquil L, Soulié V, Tran N, Gidenne T and Fortun-Lamothe L 2011. Postnatal development of the rabbit caecal microbiota composition and activity. *FEMS Microbiology Ecology* 77, 680–689.

Dabbou S, Rotolo L, Kovitvadhi A, Bergagna S, Dezzutto D, Barbero R, Rubiolo P, Schiavone A, De Marco M, Helal A N, Zoccarato I and Gasco L 2016. Rabbit dietary supplementation with pale purple coneflower. 1. Effects on the reproductive performance and immune parameters of does. *Animal*, doi:10.1017/S1751731115002979.

Egert D and Beuscher N 1992. Studies on antigen specificity of immunoreactive arabinogalactan proteins extracted from *Baptisia tinctoria* and *Echinacea purpurea*. *Planta Medica* 58, 163–165.

Franz C, Baser KHC and Windisch W 2010. Essential oils and aromatic plants in animal feeding—a European perspective. a review. *Flavour and Fragrance Journal* 25, 327–340.

Gidenne T, Combes S, Feugier A, Jehl N, Arveux P, Boisot P, Briens C, Corrent E, Fortune H, Montessuy S and Verdelhan S 2009. Feed restriction strategy in the growing rabbit. 2. Impact on digestive health, growth and carcass characteristics. *Animal* 3, 509–515.

Hermann JR, Honeyman MS, Zimmerman JJ, Thacker BJ, Holden PJ and Chang CC 2003. Effect of dietary *Echinacea purpurea* on viremia and performance in porcine reproductive and respiratory syndrome virus-infected nursery pigs. *Journal of Animal Science* 81, 2139–2144.

Hudson J, Vimalanathan S, Kang L, Amigue VT, Livesey J and Arnason JT 2005. Characterization of antiviral activities in *Echinacea* root preparations. *Pharmaceutical Biology* 43, 790–796.

IBM SPSS 2012. IBM SPSS Statistics 20.0. SPSS Inc, Chicago, IL, USA.

Kraus GA, Bae J, Wu L and Wurtele E 2007. The synthesis and natural distribution of the major ketone constituents in *Echinacea pallida*. *Molecules* 12, 406–414.

Ma A, Shi W, Niu X, Wang M and Zhong X 2009. Effects of *Echinacea purpurea* extract on the immunological response to infectious bursal disease vaccine in broilers. *Frontiers of Agriculture in China* 3, 452–456.

Maass N, Bauer J, Paulicks BR, Böhmer BM and Roth-Maier DA 2005. Efficiency of *Echinacea purpurea* on performance and immune status in pigs. *Journal of Animal Physiology and Animal Nutrition* 89, 244–252.

Michelland RJ, Combes S, Monteils V, Cauquil L, Gidenne T and Fortun-Lamothe L 2010. Molecular analysis of the bacterial community in digestive tract of rabbit. *Anaerobe* 16, 61–65.

Michelland RJ, Combes S, Monteils V, Cauquil L, Gidenne T and Fortun-Lamothe L 2011. Rapid adaptation of the bacterial community in the growing rabbit caecum after a change in dietary fibre supply. *Animal* 5, 1761–1768.

Muyzer G, Waal E and Uitterlinden A 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology* 59, 695–700.

Ragap HM, Khalil RH and Mutawie HH 2012. Immunostimulant effects of dietary *Spirulina platensis* on tilapia *Oreochromis niloticus*. *Journal of Applied Pharmaceutical Science* 2, 26–31.

Sahin T, Ozlem K and Mehmet S 2012. Effects of ground *Echinacea* (*Echinacea purpurea*) supplementation quail diets on growth performance and carcass traits. *Kafkas Universitesi Veteriner Fakültesi Dergisi* 18, 15–19.

Tizard I 2013. *Veterinary immunology*, 9th edition. Elsevier Saunders, St. Louis, Missouri, USA.

Varga M 2014. *Text book of rabbit medicine*, 2nd edition. Butterworth–Heinemann, Oxford, UK.

Zhai Z, Liu Y, Wu L, Senchina DS, Wurtele ES, Murphy PA, Kohut ML and Cunnick JE 2007. Enhancement of innate and adaptive immune functions by multiple *Echinacea* species. *Journal of Medicinal Food* 10, 423–434.