



UNIVERSITY
OF TRENTO

DEPARTMENT OF INFORMATION AND COMMUNICATION TECHNOLOGY

38050 Povo – Trento (Italy), Via Sommarive 14
<http://www.dit.unitn.it>

KÖNIG'S EDGE COLORING THEOREM WITHOUT AUGMENTING
PATHS

Romeo Rizzi

October 1999

Technical Report # DIT-02-0057

Also : in *Journal of Graph Theory* 29 (1998) 87.

König's Edge Coloring Theorem without augmenting paths

Romeo Rizzi

October 15, 1999

Dipartimento di Matematica Pura ed Applicata, Università di Padova

Via Belzoni 7, 35131 Padova, Italy

`romeo@euler.math.unipd.it`

We give a simple, self-contained proof of the following basic fact [1, 2] in matching theory:

Theorem *Every bipartite regular multigraph is factorizable.*

Proof: Assume G to be a counterexample with the smallest number of edges. Then G is r -regular for some integer $r \geq 1$. Let $e = uv$ be any edge of G . In G , remove nodes u and v , then add a set of edges F as to obtain a bipartite multigraph G' which is r -regular. Note that $|F| < r$ and G' has less edges than G . By assumption G' contains r disjoint 1-factors. Since $|F| < r$, at least one of these 1-factors, say M' , is disjoint from F . Therefore $M = M' \cup \{e\}$ is a 1-factor of G . Moreover $G \setminus M$ has less edges than G and is factorizable. But then G is factorizable. \square

Note that the above proof does not use any alternating path argument.

References

- [1] D. König, Über Graphen und ihre Anwendung auf Determinantentheorie und Mengenlehre, Math. Ann. 77 (1916) 453–465.
- [2] D. König, Graphok és alkalmazásuk a determinánsok és a halmazok elméletére, Math. Termész. Ért. 34 (1916) 104–119.