


RESEARCH

Open Access



Oxygenation improvement and duration of prone positioning are associated with ICU mortality in mechanically ventilated COVID-19 patients

Silvia De Rosa^{1,2†}, Nicolò Sella^{3†}, Giacomo Bellani¹, Giuseppe Foti⁴, Andrea Cortegiani^{5,6}, Giulia Lorenzoni⁷, Dario Gregori⁷, Annalisa Boscolo^{3,8}, Lucia Cattin², Muhammed Elhadi⁹, Giorgio Fullin¹⁰, Eugenio Garofalo¹¹, Leonardo Gottin¹², Alberto Grassetto¹³, Salvatore Maurizio Maggiore^{14,15}, Elena Momesso¹⁶, Mario Peta¹⁷, Daniele Poole¹⁸, Roberto Rona⁴, Ivo Tiberio¹⁹, Andrea Zanoletti²⁰, Emanuele Rezoagli^{4†}, Paolo Navalesi^{3,8*†}  and for the SIAARTI Study Group

Abstract

Background Prone position has been diffusely applied in mechanically ventilated COVID-19 patients. Our aim is ascertaining the association between the physiologic response and the length of the first cycle of prone position and intensive care unit (ICU) mortality.

Methods International registry including COVID-19 adult patients who underwent prone positioning. We measured the difference for arterial partial pressure of oxygen to inspired fraction of oxygen ratio (PaO₂/FiO₂), ventilatory ratio, and respiratory system compliance (Cr_s) between baseline supine position and at either the end of the first cycle of prone position (Delta-PP) or re-supination (Delta-PostPP).

Results We enrolled 1816 patients from 53 centers. Delta-PP and Delta-PostPP for PaO₂/FiO₂ were both associated with ICU mortality [OR (95% CI) 0.48 (0.38, 0.59), and OR (95% CI) 0.60 (0.52, 0.68), respectively]. Ventilatory ratio had a non-linear relationship with ICU mortality for Delta-PP (p=0.022) and Delta-PostPP (p=0.004). Delta-PP, while not Delta-PostPP, for Cr_s was associated with ICU mortality [OR (95% CI) 0.80 (0.65, 0.98)]. The length of the first cycle of prone position showed an inverse relationship with ICU mortality [OR (95% CI) 0.82 (0.73, 0.91)]. At the multi-variable analysis, the duration of the first cycle of prone position, Delta-PP and Delta-PostPP for PaO₂/FiO₂, and Delta-PostPP for ventilatory ratio were independently associated with ICU mortality.

Conclusion In COVID-19 patients with acute respiratory failure receiving invasive mechanical ventilation and prone positioning, the physiological response to prone position is associated with ICU mortality. Prolonging the duration of the first cycle of prone position is associated with improved survival.

Keywords Prone position, Mechanical ventilation, Acute respiratory failure, Arterial partial pressure of oxygen to inspired fraction of oxygen ratio (PaO₂/FiO₂), Ventilatory ratio, Respiratory system compliance

[†]Silvia De Rosa and Nicolò Sella equally contributed, and also Emanuele Rezoagli and Paolo Navalesi equally contributed to this work.

*Correspondence:

Paolo Navalesi

paolo.navalesi@unipd.it

Full list of author information is available at the end of the article

Background

In patients with moderate-to-severe acute respiratory distress syndrome (ARDS) receiving invasive mechanical ventilation, prone position may improve arterial blood gases through different mechanisms such as rearrangement of the gas–tissue ratios along the dependent–nondependent regions of the lungs, enhanced ventilation/perfusion matching, and a more homogeneous distribution of lung stress and strain [1]. A large multicenter randomized trial found that prone positioning within 24 h after the diagnosis of ARDS and for at least 16 h for the first maneuver and 12 h for the following ones reduced both 28-day and 90-day mortality in moderate-to-severe ARDS patients [2]. Afterwards, other studies and meta-analyses confirmed a survival benefit of prone positioning in moderate-to-severe ARDS [3–5], so that since 2017 prone position has been strongly recommended with a high level of evidence for patients with severe or moderate-to-severe ARDS to reduce mortality [6–8].

Nonetheless, before the coronavirus disease 2019 (COVID-19) pandemic, prone position struggled to become part of the routine clinical practice for the management of patients with moderate-severe ARDS, where prone positioning was implemented in less than 1 out of 10 cases within 48 h [9] and not even performed in almost 70% of patients with severe ARDS prior to veno-venous extracorporeal membrane oxygenation (V-V ECMO) cannulation in experienced centers [10].

Quite the opposite, prone positioning has been diffusely adopted during the COVID-19 pandemic. In fact, prone position was used in 76% of 735 COVID-19-related ARDS patients included in a multicenter cohort in Spain [11], and in 61% of 1057 patients on invasive mechanical ventilation for COVID-19 acute respiratory failure in an Italian multicenter cohort study [12].

During the COVID-19 pandemic we set up an international multicenter registry, the Prone Positioning for Invasively Ventilated Patients with COVID-19 (PROVENT-C19) Registry [13], which aimed at ascertaining the association between Intensive Care Unit (ICU) mortality and the response to the first cycle of prone position, as assessed by physiological parameters [1] such as arterial oxygenation, dead space fraction estimates, and respiratory system compliance. Secondly, we aimed at profiling patients based on the physiological response to prone position in a large cohort of patients experiencing COVID-19 acute respiratory failure. Moreover, we investigated the associations between ICU mortality and both the length of the first cycle of prone position and the total number of prone position cycles.

Methods

The PROVENT-C19 is a multicenter, observational registry developed by the Veneto ICU Network [14], and endorsed by the Italian Society of Anesthesia, Analgesia, Resuscitation and Intensive Care, the European Society of Intensive Care Medicine, and the European Society of Anaesthesiology and Intensive Care [13]. There is no funding source.

The registry was designed following the Declaration of Helsinki and the study protocol was firstly approved by the Vicenza Ethics Committee (ID:22/21). Informed consent was obtained according to the European Union General Data Protection Regulation, the national regulations or local directives of each participating Institution. In cases the patient was incompetent because of critical illness or the use of sedative or anesthetic drugs, consent could be delayed, and a provision for delayed consent was applied: as soon as competent, each patient was fully informed on what had been done, and a written permission of using data collected was obtained. The patients or their legal surrogates were informed of their right to request that the study procedures be discontinued and their right to refuse the study-related use of their medical records.

The Strengthening the Reporting of Observational Studies in Epidemiology reporting guideline checklist for observational studies was used for reporting this study (Supplementary Material 1).

The registry included, either prospectively or retrospectively, consecutive adult patients who underwent invasive mechanical ventilation and prone positioning due to COVID-19 related acute respiratory failure, from December 31st 2019 to January 1st 2023. Patients were excluded if they refused the consent to participate or if they presented contraindications to prone position [1, 15].

Physiological response to prone position was evaluated at different time points (Fig. 1), calculating for arterial partial pressure of oxygen to fraction of inspired oxygen ratio ($\text{PaO}_2/\text{FiO}_2$), ventilatory ratio (as proxy for dead space) [16, 17], and static compliance of the respiratory system [Crs] [18–20]:

- Difference in prone position (Delta-PP): the difference between the last available values within the last 30 min in prone and supine position (prior to being turned prone), respectively;
- Difference after prone positioning (Delta-PostPP): the difference between the first available value within the first 30 min in supine position after the end of the prone position cycle and the last available value in supine position before being turned prone (in the last 30 min prior to being turned prone).




Fig. 1 Study design and timeline. These time windows were desirable but not obligatory. The real timing of variable collection was recorded and variables registered more than 30 min later than required by the study design were excluded. $\text{PaO}_2/\text{FiO}_2$ arterial partial pressure of oxygen to inspire fraction of oxygen ratio. Vent Ratio, ventilatory ratio. Crs, static compliance of the respiratory system

Patients were categorized according to:

- ICU survival;
- length of the first cycle of prone position in three groups: (1) short pronation (i.e., < 16 h); (2) long pronation (between 16 and 24 h); and (3) extended pronation (i.e., > 24 h).

Statistical analysis

Complete case analysis was performed.

Categorical data are presented as absolute numbers (n) and percentages (%). For continuous data medians and I and III quartiles are used. The distribution of categorical variables was compared with the Pearson's Chi-squared test or the Fisher's exact test, whichever appropriate, while the distribution of continuous variables was compared using Wilcoxon rank sum test. To account for multiplicity of testing, Benjamini–Hochberg correction was employed.

Univariable logistic regression models were employed to assess the association between patients' characteristics or physiological response to prone position and ICU mortality. For nonlinear association, restricted cubic splines were used. To account for clustering effect within the same center, the Huber-White method was employed.

A multivariable logistic regression model was used to evaluate the association between physiological response to prone position and ICU mortality after adjusting for relevant confounders, including the ICU admission period.

A subgroup analysis was conducted to assess the consistency of findings in retrospective and prospective data: logistic regression models for ICU mortality were fitted, including interaction terms between Delta-PP and recruitment type (prospective vs. retrospective) and between Delta-PostPP and recruitment type (prospective vs. retrospective).

All statistical tests were 2-tailed, and statistical significance was defined as $p < 0.05$. All analyses have been conducted using R 4.0.3 (Vienna, Austria).

Additional details on methods and statistics are provided in the Supplementary Material 2.

Results

A total of 1816 patients were enrolled from 53 international centers (Supplementary Material 3 and Supplementary Material 4). The baseline characteristics of the study population are shown in Table 1 and Supplementary Material 5, while the main clinical outcomes are listed in Table 2. Overall, 837 patients (47.7%) were discharged alive from ICU, while 916 patients (52.3%) died

Table 1 Demographic and baseline characteristics of the study population

Characteristics	Overall N	ICU survivors n=837 (47.7%)	ICU non-survivors n=916 (52.3%)	OR (95% CI)	P-value
<i>Demographics</i>					
Age, years	1752	61 (53, 69)	69 (62, 74)	2.47 (1.92, 3.18)	< 0.001
Male, n (%)	1750	576 (69.0)	659 (72.0)	1.16 (0.98, 1.36)	0.077
BMI, kg/m ²	1667	29 (26, 33)	28 (25, 31)	0.94 (0.84, 1.05)	0.269
Pandemic wave, n (%):	1753				
- 1st wave (January 2020–August 2020)		94 (11.0)	70 (7.6)	–	–
- 2nd wave (September 2020–March 2021)		362 (43.0)	462 (50.0)	1.71 (1.22, 2.41)	0.002
- 3rd wave (April 2021–December 2022)		381 (46.0)	384 (42.0)	1.35 (0.96, 1.91)	0.081
<i>Comorbidities</i>					
COPD, n(%)	1751	41 (4.9)	127 (14.0)	3.12 (2.01, 4.84)	< 0.001
Arterial hypertension, n (%)	1752	393 (47.0)	580 (63.0)	1.94 (1.54, 2.44)	< 0.001
Chronic heart failure, n (%)	1752	69 (8.3)	164 (18.0)	2.42 (1.69, 3.46)	< 0.001
Cerebral vasculopathy, n (%)	1752	23 (2.8)	48 (5.2)	1.95 (1.12, 3.42)	0.019
Diabetes mellitus, n (%)	1752	172 (21.0)	270 (29.0)	1.61 (1.26, 2.06)	< 0.001
Chronic kidney disease, n (%)	1751	11 (1.3)	67 (7.3)	5.93 (2.78, 12.62)	< 0.001
Home renal replacement therapy, n (%)	1751	5 (0.6)	11 (1.2)	0.31 (0.08, 1.24)	0.097
Chronic liver failure, n (%)	1752	9 (1.1)	24 (2.6)	2.47 (1.18, 5.17)	0.016
Cancer, n (%)	1752	26 (3.1)	53 (5.8)	1.91 (1.11, 3.30)	0.020
Immunological deficiency, n (%)	1751	41 (4.9)	91 (9.9)	2.14 (1.43, 3.20)	< 0.001
<i>Before ICU admission</i>					
COVID-19 vaccination, n (%)	1232	83 (14.0)	99 (15.0)	1.07 (0.71, 1.60)	0.750
Hospitalization before ICU admission, days	1752	2 (1, 5)	3 (1, 7)	1.00 (0.99, 1.00)	0.372
Corticosteroids before ICU admission, n(%)	1745	566 (68.0)	655 (72.0)	1.22 (0.80, 1.86)	0.354
Anticoagulant therapy before ICU admission, n (%)	1744	542 (65.0)	610 (67.0)	1.10 (0.76, 1.61)	0.611
Non-invasive respiratory support before IMV, n (%)	1749	728 (87.0)	790 (87.0)	0.97 (0.65, 1.46)	0.881
Non-invasive respiratory support before IMV, days	1326	2 (1, 4)	4 (2, 7)	1.57 (1.28, 1.93)	< 0.001
<i>ICU admission</i>					
IMV at ICU admission, n (%)	1751	213 (25.0)	306 (33.0)	1.41 (0.85, 2.35)	0.184
PaO ₂ /FiO ₂ at ICU admission, mmHg	1742	90 (69, 118)	84 (65, 114)	0.92 (0.82, 1.02)	0.108
Glasgow Coma Scale at ICU admission	1711	15 (15, 15)	15 (14, 15)	0.45 (0.13, 1.57)	0.212
SOFA at ICU admission	1720	4 (3, 5)	4 (4, 7)	1.54 (1.25, 1.90)	< 0.001
White Blood Cells at ICU admission, × 10 ⁹ /L	1735	10 (7, 13)	11 (8, 15)	1.00 (0.99, 1.01)	0.315
CRP at ICU admission, mg/L	1413	48 (13, 130)	38 (11, 125)	0.97 (0.76, 1.25)	0.839
Procalcitonin at ICU admission, mcg/L	1244	0.20 (0.10, 0.50)	0.26 (0.12, 0.78)	1.01 (1.00, 1.02)	0.189
D-Dimer at ICU admission, mcg/L	1178	912 (354, 2020)	1099 (35, 3898)	1.03 (0.99, 1.07)	0.124

Results of the univariable analysis

Data are median (I quartile–III quartile) for continuous variables and absolute numbers (percentages) for categorical variables. Results of the univariable analyses are reported as Odds Ratio (OR), 95% Confidence Interval (CI), *p*-value

ICU intensive care unit, OR Odds Ratio, 95%CI 95% Confidence Interval, BMI body mass index, COPD chronic obstructive pulmonary disease, IMV invasive mechanical ventilation, PaO₂/FiO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio, SOFA sequential organ failure assessment, CRP C-reactive protein

during the ICU stay. The risk factors associated with ICU mortality are reported in Tables 1 and 2.

Physiological response to prone position

The response to the first cycle of prone position on the basis of PaO₂/FiO₂ was available for 1655 patients. As depicted in Fig. 2A, B, both Delta-PP and Delta-PostPP

for PaO₂/FiO₂ showed an inverse linear relationship with ICU mortality [OR (95% CI) 0.48 (0.38, 0.59), *p* < 0.001, and OR (95% CI) 0.60 (0.52, 0.68), *p* < 0.001, respectively]. By plotting Delta-PostPP for PaO₂/FiO₂ versus Delta-PP for PaO₂/FiO₂, patients were divided in four quadrants according to their oxygenation responses (improvement or worsening) to the first cycle of prone position

Table 2 Main clinical outcomes of the study population

Characteristics	Overall N	ICU survivors n=837 (47.7%)	ICU non-survivors n=916 (52.3%)	OR (95% CI)	P-value
<i>Outcomes</i>					
ICU LOS, days	1748	16 (10, 28)	13 (8, 22)	0.90 (0.79, 1.02)	0.091
Hospital LOS, days	1713	35 (22, 52)	19 (12, 28)	0.45 (0.30, 0.67)	<0.001
28-day ventilator free days	992	15 (3, 20)	0 (0, 0)	0.58 (0.50, 0.64)	<0.001
Tracheostomy, n (%)	1751	265 (32.0)	197 (22.0)	0.59 (0.40, 0.87)	0.007
Renal replacement therapy, n (%)	1741	33 (4.0)	151 (17.0)	4.81 (2.89, 8.01)	<0.001
ECMO or ECCO ₂ R, n (%)	1727	19 (2.3)	27 (3.0)	1.30 (0.54, 3.14)	0.565
iNO, n (%)	1631	49 (6.2)	52 (6.2)	1.00 (0.56, 1.79)	0.998
Cycles of prone position, n (%)	1753	2 (2, 4)	3 (2, 5)	1.06 (0.92, 1.11)	0.102
Overall time in prone position, hours	1651	64 (40, 98)	63 (38, 102)	1.02 (0.85, 1.23)	0.835
Hospital mortality, n (%)	1720	39 (4.8)	\	\	\

Results of the univariable analysis

Data are median (I quartile-III quartile) for continuous variables and absolute numbers (percentages) for categorical variables. Results of the univariable analyses are reported as Odds Ratio (OR), 95% Confidence Interval (CI), *p*-value

ICU intensive care unit, OR Odds Ratio, 95%CI 95% Confidence Interval, LOS length of stay, ECMO extracorporeal membrane oxygenation, ECCO₂R extracorporeal carbon dioxide removal, iNO inhaled nitric oxide

(Supplementary Material 6). The group who experienced an increase in PaO₂/FiO₂ both at the end of prone position (Delta-PP > 0 mmHg) and at re-supination (Delta-PostPP > 0 mmHg) presented the lower ICU mortality [47.0 (44.0–50.0) %]. Worth remarking, patients underwent further cycles of prone positioning in the vast majority of cases, either in case Delta-PP for PaO₂/FiO₂ improved (92%) or not (87%) within the first cycle of prone position.

Secondly, the response to the first cycle of prone position with respect to the ventilatory ratio was assessed in 1509 patients. As depicted in Fig. 2C, D, both Delta-PP and Delta-PostPP for the ventilatory ratio showed a non-linear relationship with ICU mortality (*p*-value

for non-linearity = 0.022 and 0.004, respectively), with a linear increase of the risk of death in ICU at increasing values of Delta-PP and Delta-PostPP for ventilatory ratio only for values greater than 0.04 and 0.09, respectively.

Finally, the response to the first cycle of prone position on the basis of Crs was available in 469 patients. As depicted in Fig. 2E, Delta-PP for Crs showed an inverse linear relationship with ICU mortality [OR (95% CI) 0.80 (0.65, 0.98), *p* = 0.034]. Delta-PostPP for Crs exhibited a trend towards an inverse relationship with ICU mortality, but the association was not significant [OR (95% CI) 0.84 (0.66, 1.06), *p* = 0.067] (Fig. 2F).

(See figure on next page.)

Fig. 2 Relationship between ICU mortality and the response to the first cycle of prone position on the basis of arterial oxygenation at the end of the prone position cycle (Delta-PP) (A) and after re-supination (Delta-PostPP) (B), on the basis of ventilatory ratio at the end of the prone position cycle (Delta-PP) (C) and after re-supination (Delta-PostPP) (D), and on the basis of respiratory system static compliance at the end of the prone position cycle (Delta-PP) (E) and after re-supination (Delta-PostPP) (F). The x-axis shows the predictor and the y-axis shows the effect of the predictor on the outcome in log-odds. The solid line represents the regression line estimated by the logistic regression model, and the gray bands show the confidence intervals. When the relationship was found to be non-linear, the change-point was identified (red dotted line). Log-odds quantify the influence of factors on the probability of an outcome: a positive value indicates that the probability of the event increases with the factor, while a negative value suggests a decrease. If we suppose the logistic regression model gives a log-odds for Delta-PP for PaO₂/FiO₂ of -0.05, this log-odds value indicates that for each additional point of Delta-PP for PaO₂/FiO₂, the log-odds of death in the ICU decreases by 0.05. In panel A, the odds ratio of 0.48 for ICU mortality associated with an increase of Delta-PP for PaO₂/FiO₂ indicates that an improvement in the PaO₂/FiO₂ ratio within the interquartile range (i.e., from 26 to 124 mmHg) significantly reduces the odds of ICU mortality. Specifically, this increase from the 25th to the 75th percentile is associated with a 52% reduction in the odds of ICU mortality. The same rule could be applied for the interpretation of the linear relationships depicted in panels B, E, and F. When the relationship is non-linear, as in panels C and D, the odds ratio for ICU mortality refers to an increase of the variable within the range indicated in the table below the chart. For example, in panel C, the odds ratio of 1.27 for ICU mortality associated with the increase of Delta-PP for ventilatory ratio from 0.0 to 0.5 indicates that the odds of ICU mortality are significantly increased if Delta-PP for ventilatory ratio passes from 0.0 to 0.5. PaO₂/FiO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio. Crs, static compliance of the respiratory system. OR, odds ratio. 95%CI, 95% confidence interval




Fig. 2 (See legend on previous page.)

Duration and frequency of prone position cycles

The length of the first cycle of prone position showed a linear inverse relationship with ICU mortality [OR (95% CI) 0.82 (0.73, 0.91), $p=0.001$], while the frequency of the prone position cycles was not associated with ICU mortality [OR (95% CI) 1.13 (0.98, 1.31), $p=0.087$] (Fig. 3). ICU mortality did not differ significantly between those who were not treated with further cycles of prone positioning after the first one and those who underwent multiple cycles (46% vs 53%, respectively, $p=0.186$).

The baseline and clinical characteristics and the main outcomes of patients according to the short, long or extended pronation group are shown in Supplementary Material 7. Worth remarking, short pronation was more frequently adopted during the first wave of COVID-19 pandemic (January 2020–August 2020), accounting for 44.1% of the overall patients registered in this period ($p<0.001$), while extended pronation was more extensively applied during the third wave (April 2021–December 2023), accounting for 17.4% of the overall patients recorded in this period ($p<0.001$). The short pronation group experienced significantly lower Delta-PP for PaO₂/FiO₂ [45 (6–95) mmHg] compared to the long [73 (33–125) mmHg] and extended pronation group [72 (30, 125) mmHg] ($p<0.001$). Likewise, the extended pronation group showed the greatest Delta-PostPP for PaO₂/FiO₂ [39 (11–85) mmHg] compared to the other two groups [17 (– 1–58) mmHg,

and 28 (– 1–66) mmHg, respectively] ($p<0.001$). Conversely, no differences between the three groups were identified in Delta-PP and Delta-PostPP for the ventilatory ratio and Crs.

As depicted in Supplementary Material 8, a non-linear relationship was found between the length of the first cycle in prone position and the oxygenation response as assessed by Delta-PP for PaO₂/FiO₂ (p -value for non-linearity <0.001), with a linear increase of Delta-PP at increasing duration up to 19 h. The length of the first cycle of prone position and Delta-PostPP for PaO₂/FiO₂ showed a direct linear relationship [OR (95% CI) 7.21 (3.20, 11.23), $p<0.001$].

Multivariable analysis for ICU mortality

At the multivariable analysis, in the model 1 both the duration of the first cycle of prone position [OR (95% CI) 0.84 (0.72, 0.96)] and Delta-PP for PaO₂/FiO₂ [OR (95% CI) 0.51 (0.41, 0.63)] were confirmed to be independently associated with ICU mortality, while not Delta-PP for ventilatory ratio [OR (95% CI) 1.07 (0.95, 1.20)]. In the model 2 the duration of the first cycle of prone position [OR (95% CI) 0.82 (0.71, 0.95)], Delta-PostPP for PaO₂/FiO₂ [OR (95% CI) 0.58 (0.47, 0.71)], and Delta-PostPP for ventilatory ratio [OR (95% CI) 1.20 (1.05, 1.37)] were all independently associated with ICU mortality (Table 3).




Fig. 3 Relationship between ICU mortality and the length of the first cycle of prone position (**A**) and the total numbers of prone position cycles (**B**). The x-axis shows the predictor and the y-axis shows the effect of the predictor on the outcome in log-odds. The solid line represents the regression line estimated by the logistic regression model, and the gray bands show the confidence intervals. Log-odds quantify the influence of factors on the probability of an outcome: a positive value indicates that the probability of the event increases with the factor, while a negative value suggests a decrease. If we suppose the logistic regression model gives a log-odds for the length of prone position of -0.05 , this log-odds value indicates that for each additional point of prone position duration, the log-odds of death in the ICU decreases by 0.05. OR, odds ratio. 95%CI, 95% confidence interval

Table 3 Results of the multivariable analysis for ICU mortality

n = 945	Variable interquartile range	ODDS ratio	95% CI
<i>MODEL 1</i>			
Age, years	57, 73		
Arterial hypertension (Ref. No)	\		
Chronic heart failure (Ref. No)	\		
Pandemic wave	\		
SOFA at ICU admission (unit increase)	3, 6		
D-Dimer at ICU admission, mcg/L	285, 2737		
Length of the 1st cycle in prone position, hours	16, 25	0.84	0.72, 0.96
Delta-PP for PaO ₂ /FiO ₂ , mmHg	26, 124	0.51	0.41, 0.63
Delta-PP for ventilatory ratio	− 0.15, 0.28	1.07	0.95, 1.20
<i>MODEL 2</i>			
Age, years	57, 73		
Arterial hypertension (Ref. No)	\		
Chronic heart failure (Ref. No)	\		
Pandemic wave	\		
SOFA at ICU admission (unit increase)	3, 6		
D-Dimer at ICU admission, mcg/L	285, 2737		
Length of the 1st cycle in prone position, hours	16, 25	0.82	0.71, 0.95
Delta-PostPP for PaO ₂ /FiO ₂ , mmHg	1, 70	0.58	0.47, 0.71
Delta-PostPP for ventilatory ratio	− 0.15, 0.38	1.20	1.05, 1.37

Data are Odds Ratio, 95% Confidence Interval (CI)

95%CI 95% Confidence Interval, SOFA sequential organ failure assessment, ICU intensive care unit, PaO₂/FiO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio

Subgroup analysis according to recruitment type

As reported in Supplementary Material 9, the effects of Delta-PP and Delta-PostPP on ICU mortality did not differ between the retrospective and prospective subgroups.

Discussion

In this multicenter international registry, exploring a large cohort of mechanically ventilated COVID-19 patients with acute respiratory failure, we found the physiological response to prone position to be associated with ICU mortality. The oxygenation improvement, observed either at the end of the first cycle of prone position or after re-supination, was inversely associated with ICU mortality, with the lowest death rate in patients experiencing oxygenation improvement both during prone position and after re-supination. Moreover, we showed that the worsening of either dead space or respiratory system compliance during the first cycle of prone position was associated with a higher risk of ICU mortality. Finally, in our study population the longer the duration of the first prone position cycle, the smaller the ICU mortality. After adjustment for clinically meaningful variables, both an improvement in oxygenation and the duration of the first prone positioning confirmed to be independently associated with ICU mortality.

In COVID-19 patients, Camporota and colleagues reported that the increase in oxygenation upon prone position was an independent predictor of ICU survival [21]. Imputed mechanisms of improved oxygenation during prone position include a higher ratio between the resolution of dorsal atelectasis and the onset of ventral atelectasis—as described by computed tomography data [22]—and improved ventilation perfusion matching [15, 23]. In our population we confirmed the positive effects of prone position on oxygenation and its independent association with ICU mortality. These findings differ from the secondary analysis of the Proseva trial by Albert et al. [24], who did not observe any association between the improvement in arterial oxygenation during prone positioning and survival in 232 patients randomized to receive prone position for severe ARDS. Differences in sample size, etiology of respiratory failure, and timing of oxygenation assessment might be factors explaining this discrepancy. Although our results show an association between improved oxygenation at End-PP and mortality, we cannot infer causality, and we do not suggest that clinical decision making be based on oxygenation response.

Furthermore, we observed a significant effect of prone position in improving oxygenation—as compared to baseline supine position—after re-supination. Of note,

the improved oxygenation upon prone position showed the lowest mortality when associated with a persistent improvement in oxygenation after re-supination.

This effect was probably obtained through an improvement in respiratory mechanics during prone position, such as an increased end expiratory lung volume and a decreased strain, as reported by Dilken et al. [25]. In keeping with these findings, we observed that an improved Crs linearly predicted a decrease in ICU mortality. This is in line with what has already been observed in severe ARDS patients, who showed improved survival when prone position improved Crs [26]. The main mechanism advocated to explain the improvement of Crs in prone position is the recruitment of the dorsal region of the lung during prone position [1].

In keeping with Gattinoni et al. who showed the response on carbon dioxide clearance after prone positioning to be directly correlated with survival in non-COVID-19 ARDS patients [27], we found ICU mortality to be linearly associated with a decreasing efficiency of alveolar ventilation and we observed a strong signal towards an increased ICU mortality in patients with greater ventilatory ratio after the first cycle of prone position, for both Delta-PP and Delta-PostPP. For instance, ventilatory ratio is a readily accessible bedside respiratory variable that functions as a proxy for physiologic dead space. Like dead space, ventilatory ratio also is correlated with mortality in ARDS, being an elevated ventilatory ratio associated with poorer outcomes in patients with ARDS [16, 28]. Recently, da Cruz et al. described the presence of 2 different COVID-19 phenotypes based on longitudinal respiratory variables and the one associated with a higher mortality rate was characterized by a higher ventilatory ratio during the first prone session [29]. Our findings, although associative, corroborate the potential beneficial implications of assessing the response in ventilatory ratio during and after prone positioning on outcomes. The increase in physiologic dead space as a response to prone positioning—as indicated by a higher ventilatory ratio—may introduce the concept of negative responders to prone positioning [27], which remains to be confirmed by future dedicated trials with causative design. If so, this new insight in the physiological response to prone position may have two clinical implications: (1) the increase in ventilatory ratio consequent to prone positioning may be utilized as a prognostic variable to infer an increase in mortality; (2) continuation of prone positioning should be carefully evaluated after a response in increased ventilatory ratio. This may serve as a future area of investigation to explore whether the increase of dead space secondary to lung recruitment maneuvers (e.g., prone positioning) may guide lung

recruitment or explain an increased mortality in respiratory failure.

Since the cornerstone study by Guerin et al., which led to the guidelines recommendation on applying one 16-h session followed by 12-h sessions of prone position in ARDS patients [2], the COVID-19 pandemic introduced the implementation of prolonged/extended duration of prone position, which resulted to be relatively safe and potentially more effective [1, 30–32].

Despite a trend toward a higher rate of pressure ulcers in the extended prone position subgroup (≥ 24 h) as compared with subgroups receiving shorter durations, not achieving, nonetheless, statistical significance, our data overall confirm that prolonged/extended prone position does not add harm. We found that prolonged prone position significantly improved oxygenation both for Delta-PP and Delta-PostPP, indicating an exposure-dependent response to prone position in this cohort of patients [33]. Noteworthy, while the duration of the first cycle of prone positioning was linearly associated to a higher response in oxygenation during prone position up to a plateau of 19 h, it was linearly associated with a constantly increasing oxygenation after re-supination.

More importantly, our data demonstrated for the first time in a large population an inverse linear relationship between length of application of the first prone position and ICU mortality, in keeping with the findings by Okin et al. who showed in mechanically ventilated COVID-19 hypoxemic patients that cycles ≥ 24 h reduced short- and long-term mortality [30].

This international multicenter clinical registry has strengths, including (1) the large sample size; (2) the high numbers of centers involved in the study which helps a high generalizability of the study findings; and (3) the noticeable data granularity.

We would be remiss not to mention some of the limitations of our study. First, it is a real-life registry that bears all the limitations of this study design. In particular, the study did not comprehensively explore potential confounding factors and sources of bias, such as variations in patient management protocols (e.g., implementation of awake prone positioning) and inconsistencies in data recording practices across different healthcare facilities. However, to limit the cohort effect due to the dynamics of disease characteristics and clinical management of critically ill COVID-19 patients over the time course of the pandemic [34], the multivariable analysis was corrected to account for the potential confounding effect of the ICU admission period. Second, since the registry included either prospective or retrospective data, information bias related to the retrospective nature of some of the collected data cannot be ruled out. Third, since our study focused on COVID-19 patients only, whether

or not these results may be extended to other patient populations remains to be ascertained. However, allowing for the first time the collection of large amounts of data in an homogenous population of patients with acute hypoxemic respiratory failure caused by the same disease, COVID-19 represented a good model for studying acute respiratory failure secondary to viral pneumonia, despite some specific characteristics of the COVID-19. Furthermore, differences in hospital resources and surge during the COVID-19 pandemic among different hospitals cannot be excluded and this may have been linked to patients' outcomes. Indeed, variations in the duration of the first cycle may be driven by variations in "quality" and capacity of care in each ICU: if an ICU is short-staffed, overworked, or facing a surge of patients, personnel may be not be able to prone for long periods, thus patients could be more likely to die just because staff is overwhelmed and less attentive to react to complications. Whether or not this may have played a role in the number and duration of prone position cycles during the daily care and their potential impact on ICU mortality cannot be ascertained. Last, considering the design of the study, mechanisms underlying the physiological and clinical response to prone positioning are not investigated. These limitations underscore the need for cautious interpretation of the findings and highlight areas for further research to enhance our understanding of the topic.

Implication for practice

The findings of this large multicenter study may have implications for clinical practice. Indeed, our results seem to suggest that either an improvement of arterial oxygenation at the end of the first cycle of prone position, especially when maintained after re-supination, or an increase of Crs in prone position seem to be favorable prognostic indexes for hypoxemic patients with acute respiratory failure receiving invasive mechanical ventilation and prone positioning.

Furthermore, because a longer duration of the first prone position cycle is associated with a greater improvement in arterial oxygenation after re-supination and a lower ICU mortality without major adverse effects, it may be reasonable to maintain a patient as long as possible in prone position in order to maximize its beneficial effects and reduce ICU mortality, and it is justified to address the lengthening of PP in large scale randomized clinical trials.

Conclusions

In this large international multicenter clinical registry including COVID-19 patients with acute respiratory failure receiving invasive mechanical ventilation and undergoing prone positioning, we found the physiological

response to prone position to be associated with ICU mortality. Furthermore, prolonging the duration of the first cycle of prone position seemed to improve oxygenation and survival. Whether or not these results may be extended to other patient populations remains to be ascertained.

Abbreviations

ARDS	Acute respiratory distress syndrome (ARDS)
COVID-19	Coronavirus disease 2019
V-V ECMO	Veno-venous extracorporeal membrane oxygenation
ICU	Intensive care unit
PaO ₂ /FIO ₂	Arterial partial pressure of oxygen to fraction of inspired oxygen ratio
Crs	Static compliance of the respiratory system
Delta-PP	Difference in prone position
Delta-PostPP	Difference after prone positioning

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13613-025-01438-y>.

Supplementary Material 1. The Strengthening the Reporting of Observational studies in Epidemiology reporting guideline checklist

Supplementary Material 2. Extended Methods

Supplementary Material 3. Patients enrollment among the participating centers

Supplementary Material 4. Study patients flowchart. Abbreviations. ICU, intensive care unit. PaO₂/FIO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio. *Combined enrollment indicates patients who were included in the registry after the end of the first cycle of prone position, but before hospital discharge, thus part of the data has been collected retrospectively, while other information has been registered prospectively.

Supplementary Material 5. Main clinical characteristics of the study population. Data are median for continuous variables and absolute numbers for categorical variables

Supplementary Material 6. Relationship between the arterial oxygenation response to the first cycle of prone positioning evaluated at the end of prone position before and after being turned supine. Abbreviations. PaO₂/FIO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio. 95%CI, 95% confidence interval. ICU, intensive care unit.

Supplementary Material 7. Baseline characteristics and main clinical outcomes of the study population stratified according to the length of the first cycle in prone position. Data are median for continuous variables and absolute numbers for categorical variables

Supplementary Material 8. Relationship between the length of the first cycle in prone position and the arterial oxygenation response to the first cycle of prone positioning evaluated at the end of prone position before and after re-supination. The x-axis shows the predictor and the y-axis shows the effect of the predictor on the outcome. The solid line represents the regression line estimated by the logistic regression model, and the gray bands show the confidence intervals. When the relationship was found to be non-linear, the change-point was identified. Abbreviations. PaO₂/FIO₂ arterial partial pressure of oxygen to inspire fraction of oxygen ratio. OR, odds ratio. 95%CI, 95% confidence interval.

Supplementary Material 9. Subgroup analysis on ICU mortality according to recruitment type

Acknowledgements

We acknowledge the role of the Scientific Secretariat of the Italian Society of Anesthesia Analgesia Resuscitation and Intensive Care (SIAARTI) in inviting centers and collecting adhesions, supporting the investigators with local

ethical approval processes, and for logistic support throughout the study period. We also acknowledge the role of the European Society of Intensive Care Medicine (ESICM), and of the European Society of Anaesthesiology and Intensive Care (ESAIC) for the endorsement in spreading the Registry internationally. We are indebted with all ICU personnel who made this work possible. Members of the SIAARTI Study Group, to be searchable through their individual PubMed and Scopus records, are listed below.

Marco Abastanotti: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Mohamed Abdel-Maboud Abdel-Maboud: Internal Medicine, Al-Hussein University Hospital, Cairo, Egypt; Abdelfatah Abdellateef Abdelmageed: Department of Emergency Medicine, Hajj Al-Mardi Mohiuddin Teaching Hospital, Omdurman, Sudan; Eissa Abdullah: Anesthesia, Surgical Intensive Care Unit and Pain Management, Ain Shams University hospitals (El-Demerdash Hospital), Cairo, Egypt; Ahmed Mohammed Abodina: Isolation Center, Gheryan Central Hospital, Gheryan, Libya; Aymen Abuelyamen: General Medicine, Ibra Hospital, Ibra, Sultanate of Oman; Abdurraouf Abusalama: ICU Department, Swany Health Isolation Center, Tripoli, Libya; Tareq Abdalla Abuzaid: Infectious Disease, Misurata Medical Center, Misurata, Libya; Romina Aceto: Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Stefano Addesa: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Daniela Alampi: Department of Surgical and Medical Science and Translational Medicine, Sapienza University of Rome, Rome, Italy; Amer Aldhalia: Internal Medicine, Nasr City Hospital For Health Insurance, Cairo, Egypt; Risoli Alessio: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Maytham Al-Juaifari: Emergency Department, Al-Amal Hospital, Najaf, Iraq; Raja Ahmed Alqandouz: Infectious Disease, Misurata Medical Center, Misurata, Libya; Mohammed Al-Sadawi: Department of Medicine, Stony Brook Medicine University, Stony Brook, USA; Marco Anderloni: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Cannone Andrea: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Enrico Andriolo: UOC Anestesia e Rianimazione, San Valentino Hospital, Montebelluna, Italy; Botto Anna: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Benvenuto Antonini: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Gian Marco Anzellotti: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Matteo Aritzu: Anesthesia and Critical Care Unit, Ospedale di Belluno, Belluno, Italy; Ahmed K. Awad: Anesthesia, Surgical Intensive Care Unit and Pain Management, Ain Shams University hospitals (El-Demerdash Hospital), Cairo, Egypt; Flavio Badii: UOC Anestesia e Rianimazione, Ospedale di Vittorio Veneto, Vittorio Veneto (TV), Italia; Hibah Bileid Bakeer: Isolation Center, Gheryan Central Hospital, Gheryan, Libya; Aziza Bakri: Internal Medicine, Tishreen University Hospital, Lattakia, Syria; Andrea Ballin: Anesthesia and Intensive Care Unit, University Hospital of Padua, Padua, Italy; Massimo Barattini: Anesthesia and Intensive Care Unit, Department of Emergency and Critical Care, Santa Maria Nuova Hospital, Florence, Italy; Mattia Barotti: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Mara Bassi: Department of Medicine, University of Padua, Padua, Italy; Mattia Bellandi: Department of Anesthesia and Intensive Care, ASST Nord-Milano - Bassini Hospital, Cinisello Balsamo, Italy; Marzia Bellin: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Agrippino Bellissima: UOC ANESTESIA E RIANIMAZIONE—ARNAS GARIBALDI CATANIA- CATANIA- ITALY; Annalisa Benini: Department of Emergency and Intensive Care, Area Critica Generale, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Francesco Berruto: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, University of Turin.; Giacomo Berta: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, University of Turin.; Marco Berti: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Emanuela Biagoni: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Eugenio Biamonte: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Giacomo Bianchetti: Anesthesia and Intensive Care, Manerbio

Hospital, Manerbio, Italy; Andrea Bianchin: UOC Anestesia e Rianimazione, San Valentino Hospital, Montebelluna, Italy; Manuela Biassetto: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Muhannud Binnawara: Isolation Department, Tripoli Central Hospital, Tripoli, Libya; Maria Bisi: Institute of Anesthesia and Intensive Care, University of Padova, Padova, Italy; Maria Maddalena Bitondo: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Nicoletta Boffa: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Michela Bombino: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Benedetta Bonazzi: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Elisa Bonetta: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Elisa Boni: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Sara Borga: Anestesia e rianimazione, Ospedale di Ciriè, via Battitore, 7/9, 10073, Ciriè, Turin.; Vincenzo Bosco: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Gloria Boscolo: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Veronica Bozzon: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Luca Brazzi: Department of Surgical Sciences, University of Turin, Turin, Italy; Department of Anaesthesia, Critical Care and Emergency, Città Della Salute e Della Scienza Hospital, Turin, Italy; Alessandro Bristot: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Niccolò Brumana: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Andrea Bruni: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Chiara Brusca: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Stefano Busani: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Guido Bussone: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, Turin.; Pietro Caironi: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, University of Turin.; Tiffany Calocero: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Matteo Campagnolo: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Claudine Canepa: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, University of Turin.; Riccardo Carlon: UOC Anestesia e Rianimazione, Ospedale di Vittorio Veneto, Vittorio Veneto (TV), Italia; Alfonso Carrara: Department of Biomedical Science, Humanitas University, Milan, Italy; Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Antonio Castelli: Department of Anesthesiology and Intensive Care Unit, ASST Fatebenefratelli Sacco, "Luigi Sacco" Hospital—Polo Universitario, Università degli Studi di Milano, Milan, Italy; Giulia Catalisano: Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Italy; Martina Cavinato: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Francesca Ceccaroni: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Maurizio Cecconi: Department of Biomedical Science, Humanitas University, Milan, Italy; Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Martina Cedrone: Department of Surgical Sciences, University of Turin, Turin, Italy; Marcello Ceola Graziadei: UOC di Anestesia e Terapia Intensiva Cardio-Toraco-Vascolare, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Matteo Cesana: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Nicola Cilloni: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Enrico Colombo: Neurointensive Care Unit, Department of Emergency and Intensive Care, San Gerardo Hospital, ASST-Monza, Italy; Riccardo Colombo: Department of Anesthesiology and Intensive Care Unit, ASST Fatebenefratelli Sacco, "Luigi Sacco" Hospital—Polo Universitario, Università degli Studi di Milano, Milan, Italy; Irene Coloretti: Anesthesia and Intensive Care Department, University

Hospital of Modena, Modena; Sabrina Congedi: Department of Medicine, University of Padua, Padua, Italy; Angela Corea: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Nicola Cornacchia: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Silvia Corrado: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Valentina Cricca: Department of translational medicine, Anesthesia and Intensive Care, Sant' Anna hospital, University of Ferrara, Italy; Chiara Crivellari: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Wanda Cursio: Department of Anesthesia and Intensive Care, University of Padua, Italy; Ernesto Dalla Mora: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Lorenzo Dall'Ara: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Vinicio Danzi: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Luca Davicco: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, Turin.; Alessandro De Cassai: Institute of Anesthesia and Intensive Care, University Hospital of Padua, Padua, Italy; Alessandro Devigili: UOC di Anestesia e Terapia Intensiva Cardio-Toraco-Vascolare, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Salvatore Di Blasi: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, Turin.; Antonino Di Fede: Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Italy; Antonia Di Giannantonio: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Pierluigi Di Giannatale: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Luna Di Matteo: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Giovanni Di Noto: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Luca Di Tizio: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Katia Donadello: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia ; Chiara Donatelli: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Deyaa Elden Elbakheet: General Medicine, Ibra Hospital, Ibra, Sultanate of Oman; Gallo Elisa: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Moataz Maher Emara: Faculty of Medicine, Mansoura University COVID isolation hospital, Mansoura, Egypt; Haneen Esaadi: Isolation Department, Tripoli Central Hospital, Tripoli, Libya; Mirca Fabbris: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Massimo Ferluga: Department of Anesthesia and Intensive Care, Cattinara University Hospital, Trieste, Italy ; Luigi Ferrante: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Francesco Filippone: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Francesco Filippone: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Elena Finotto: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Ilaria Fior: Department of Anesthesia and Intensive Care, University of Padua, Italy; Edoardo Forin: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Tommaso Fossali: Department of Anesthesiology and Intensive Care Unit, ASST Fatebenefratelli Sacco, "Luigi Sacco" Hospital—Polo Universitario, Università degli Studi di Milano, Milan, Italy; Rosa Fracchia: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Salvo Francesco: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Alessandra Gioia Francesconi: Operative Unit of Anesthesia and Intensive Therapy in Cardiosurgery, Department of Emergency and Admission, Anesthesia and Critical Areas, Umberto I Policlinic, Sapienza University, Rome, Italy; Nicola Franchetti: Department of Medicine, University of Padua, Padua, Italy; Sara Frisella: Department of Surgical, Oncological and Oral Science (Di.Chir.On.S.), University of Palermo, Palermo, Italy; Regina Frontera: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Giorgio Fullin: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Roberto Fumagalli: School of Medicine and Surgery, University of Milano-Bicocca, Milano, Italy, Department of Anesthesia and Critical Care, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy ; Elisa Furlani: Department of Medicine, University of Padua, Padua, Italy; Lorenzo Gamberini: Anesthesia, Intensive

Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Elisa Gamberini: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Emiliano Gamberini: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Leonardo Gandolfi: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Bernardetta Ganzerla: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Silvia Gasperi: Department of Anesthesia and Intensive Care, University of Padua, Italy; Ilenia Gatto: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Federico Geraldini: Institute of Anesthesia and Intensive Care, University Hospital of Padua, Padua, Italy; Monica Geremia: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Marco Giani: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Massimo Girardoni: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Lorenzo Giuntoli: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Ilaria Godi: Anesthesia and Intensive Care Unit, University Hospital of Padua, Padua, Italy; Gianlorenzo Golino: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Beatrice Gottardi: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Antonio Grande: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Massimiliano Greco: Department of Biomedical Science, Humanitas University, Milan, Italy; Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Daniela Guerra: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Amedeo Guzzardella: IRCCS Ca'Granda Ospedale Maggiore Policlinico di Milano; Abdurrahman Haddud: ICU Department, Swany Health Isolation Center, Tripoli, Libya; Hashim Talib Hashim: Al-Nassiryah teaching hospital, Nasiriyah, Iraq; Aliae AR Mohamed Hussein: Chest Department, Assiut University Hospitals, Assiut, Egypt; Blangetti Ilaria: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Teresa Iob: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Giovanni Carmine Iovino: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Mariachiara Ippolito: Department of Precision Medicine in Medical, Surgical and Critical Care (Me. Pre.C.C.), University of Palermo, Italy; Stefano Isgrò: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Cristina Jovinelli: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Abdulmuhamyin Khalleefah: Isolation Center, Gheryan Central Hospital, Gheryan, Libya; Ali Abdulsasir Kredan: ICU Department, Swany Health Isolation Center, Tripoli, Libya; Riccardo La Rosa: Department of translational medicine, Anesthesia and Intensive Care, Sant' Anna hospital, University of Ferrara, Italy; Claudio La Spisa: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Luca Landolfi: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Thomas Langer: School of Medicine and Surgery, University of Milano-Bicocca, Milano, Italy, Department of Anesthesia and Critical Care, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy; Annalisa Leroze: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; UOC Anestesia e Rianimazione, Ospedale Marcello Magalisi, ULSS 9 Scaligera, Verona, Italia; Federico Linassi: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Jacopo Lion: Department of Anesthesia and Intensive Care, University of Padua, Italy; Giulia Lo Scudato: Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Italy; Antonella Lombardo: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Federico Longhini: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Græcia University, Catanzaro, Italy; Marta Lubian: Department of Biomedical Science, Humanitas University, Milan, Italy; Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Giulia Luccarelli: Anesthesia e Rianimazione in urgenza—Azienda ospedali riuniti marche nord -San Salvatore Centrale—Pesaro; Alberto Lucchini: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Aurora Magliocca: Department of Medical Physiopathology and Transplants, University of Milan, Milano,

Italy; Giorgio Maiorelli: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Ilaria Mariani: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Anna Marinello: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Francesco Marrazzo: Department of Anesthesia and Critical Care, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy; Marina Alessandra Martin: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Nicolo Martinetti: Department of Biomedical Science, Humanitas University, Milan, Italy; Department of Anesthesiology and Intensive Care, IRCCS Humanitas Research Hospital, Milan, Italy; Ettore Martinez: Department of Emergency and Intensive Care, Area Critica Generale, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Alvise Martini: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Marilena Matteo: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; James Mattson: Department of Medicine, Stony Brook Medicine University, Stony Brook, USA; Jessica Giuseppina Maugeri: UOC ANESTESIA E RIANIMAZIONE—ARNAS GARIBALDI CATANIA- CATANIA- ITALY; Federica Mazzanti: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Francesca Medici: Department of Medicine, University of Padua, Padua, Italy; Luca Melchior: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Lorenza Menato: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Beatrice Milan: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Maiseloon Mogahed: Faculty of Medicine, Mansoura University COVID isolation hospital, Mansoura, Egypt; Francesco Mojoli: Anesthesia and Intensive Care Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo Pavia; Silvia Mongodi: Anesthesia and Intensive Care Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo Pavia; Jonathan Montomoli: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Giorgia Montrucchio: Department of Surgical Sciences, University of Turin, Turin, Italy; Department of Anaesthesia, Critical Care and Emergency, Città Della Salute e Della Scienza Hospital, Turin, Italy; Lorenza Moretto: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Giuseppe Moschini: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Elena Munari: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Giuseppe Neri: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro, Italy; Rosella Nicoletti: UOC Anesthesia and Intensive Care, "Madonna delle Grazie" Hospital, Matera, Italy; Alice Nova: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Nicoletta Nuzzo: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Andrea Maria Olivieri: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Sara Olivieri: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Sebastiano Ongaro: Department of Medicine, University of Padua, Padua, Italy; Anita Orlando: Anesthesia and Intensive Care Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo Pavia; Eman Othman: Isolation Department, Tripoli Central Hospital, Tripoli, Libya; Davide Ottolina: Department of Anesthesiology and Intensive Care Unit, ASST Fatebenefratelli Sacco, "Luigi Sacco" Hospital—Polo Universitario, Università degli Studi di Milano, Milan, Italy; Michele Pagani: Anesthesia and Intensive Care Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo Pavia; Giacomo Paluzzano: Department of Anesthesia and Intensive Care, Cattinara University Hospital, Trieste, Italy; Giulio Panciera: Anesthesia and Intensive Care UOC, 'Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy; Valesano Paolo: Anesthesiology and Intensive Care Unit, Regina Montis Regalis Hospital, Mondovì, ITALY; Francesco Papaccio: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Marcella Parente: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Zoe Parimisi: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Laura Pasin: Institute of Anesthesia and Intensive Care, Azienda Ospedaliera Universitaria di Padova, Padova, Italy;

Lucia Alessandra Pasqua: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro, Italy; Federica Pavan: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Matteo Perona: Department of Medicine, University of Padua, Padua, Italy; Paolo Persona: Institute of Anesthesia and Intensive Care, University Hospital of Padua, Padua, Italy; Tommaso Petteuzzo: Institute of Anesthesia and Intensive Care, University Hospital of Padua, Padua, Italy; Angelo Pezzi: Department of Anesthesia and Intensive Care, ASST Nord-Milano -Bassini Hospital, Cinisello Balsamo, Italy; Elisa Pistollato: Department of Medicine, University of Padua, Padua, Italy; Enrico Polati: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Melissa Polo Fritz: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Gilda Ponzone: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Matteo Pozzi: Department of Emergency and Intensive Care, Terapia intensiva cardio-toracovascolare, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Antonella Prandini: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Nicoletta Predonzani: Department of Anesthesia and Intensive Care, Cattinara University Hospital, Trieste, Italy; Chiara Premoli: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Davide Raimondi Cominesi: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Jacopo Rama: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia; Linda Ramahi: Department of Anesthesia and Intensive Care, Infermi Hospital, AUSL ROMAGNA, Rimini, Italy; Ginevra Randon: Department of Medicine, University of Padua, Padua, Italy; Marta Repishti: School of Medicine and Surgery, University of Milano-Bicocca, Milano, Italy; Andrea Restivo: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Mara Ricci: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Veronica Rizzello: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Erika Roat: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena; Monica Rocco: Department of Surgical and Medical Science and Translational Medicine, Sapienza University of Rome, Rome, Italy; Andrea Rodi: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Egle Rondelli: Department of Emergency and Intensive Care, Terapia intensiva e Semintensiva adulti e pediatrica, Fondazione IRCCS San Gerardo dei Tintori, Monza, Italy; Vincenzo Russotto: Department of anesthesia and intensive care, University hospital San Luigi Gonzaga, via Regione Gonzole, 10, 10043, Orbassano, University of Turin.; Debora Saggioro: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Francesco Saggiotti: Department of Anesthesia and Intensive Care, ASST Nord-Milano -Bassini Hospital, Cinisello Balsamo, Italy; Abdurraouf Said: Department of Anaesthesia and Intensive Care, Tripoli University Hospital (Tripoli Medical Center), Tripoli, Libya; Gabriele Sales: Department of Surgical Sciences, University of Turin, Turin, Italy; Department of Anaesthesia, Critical Care and Emergency, Città Della Salute e Della Scienza Hospital, Turin, Italy; Abdulhamid Ahmed Saliga: Infectious Disease, Misurata Medical Center, Misurata, Libya; Reem Salih: General Medicine, Ibra Hospital, Ibra, Sultanate of Oman; Michele Salvagno: Department of Medicine, University of Padua, Padua, Italy; Antonio Massimo Sammarco: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Jacopo Santi: UOC Anestesia e Rianimazione, AULSS8 Berica , Ospedale San Bortolo, Vicenza, Italia; Carola Santi: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Elisabetta Saraceni: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Gaetano Scaramuzzo: Department of translational medicine, Anesthesia and Intensive Care, Sant' Anna hospital, University of Ferrara, Italy; Chiara Schiavolin: Department of Medicine, University of Padua, Padua, Italy; Valerio Schinetti: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Vittorio Schweiger: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia, Dip. di Scienze Chirurgiche, Odontostomatologiche e Materno Infantili, Università degli Studi di Verona, Verona, Italia ; Elena Serafini: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Teodora Serana: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Luca Serano: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Laila Esnoussi Shalabi:

Isolation Center, Gheryan Central Hospital, Gheryan, Libya; Haitam Shames: Isolation Center, Gheryan Central Hospital, Gheryan, Libya; Salvatore Simari: UOC di Anestesia e Rianimazione B, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Caterina Simoni: Department of Medicine, University of Padua, Padua, Italy; Simone Smiraglia: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Fabio Soccorsi: UOC di Anestesia e Terapia Intensiva Cardio-Toraco-Vascolare, Dip di Emergenza e Terapie Intensive, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italia; Alessandra Soragni: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Silvana Sorrentino: Department of Anesthesia and Intensive Care, Ospedale Dell'Angelo, Mestre-Venice, Italy; Savino Spadaro: Department of translational medicine, Anesthesia and Intensive Care, Sant' Anna hospital, University of Ferrara, Italy; Stefano Spina: Department of Anesthesia and Critical Care, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy; Marta Talamonti: Anesthesia and Intensive Care Department, University Hospital of Modena, Modena, Italy; Francesco Talarico: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Chiara Natalia Tartivita: Anesthesia, Intensive Care and Prehospital Emergency, Maggiore Hospital, Bologna, IT; Tommaso Tenaglia: S.C. Anestesia e Rianimazione ASLTO4, Ivrea, Italy; Francesco Terranova: Anesthesia and Intensive Care Unit, University Hospital of Padua, Padua, Italy; Denise Testini: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Fabio Toffoletto: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Anna Toniolo: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Morena Tonzar: Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave; Giulia Torsello: Department of Surgical Sciences, University of Turin, Turin, Italy; Tommaso Travaglini: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Fabrizio Tritapepe: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Luigi Tritapepe: Operative Unit of Anesthesia and Intensive Therapy in Cardiosurgery, Department of Emergency and Admission, Anesthesia and Critical Areas, Umberto I Policlinic, Sapienza University, Rome, Italy; Letizia Troisi: Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro, Italy; Fabrizio Turvani: Anestesia e rianimazione, Ospedale di Rivoli, via Rivalta 29, 10098 Rivoli, Turin; Lucrezia Urso: Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Italy; Rita Vaia Liouras: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Paolo Valente: UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italia; Maria Sole Vallecoccia: Anesthesia and Intensive Care Unit, Department of Emergency and Critical Care, Santa Maria Nuova Hospital, Florence, Italy; Paola Vergano: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Sara Vergine: AOUI Anestesia e Rianimazione 1, S. Chiara, APSS Trento, Trento, Italia; Luigi Vetrugno: Department of Anesthesiology and Intensive Care, S.S. Annunziata Hospital, Chieti, Italy; Vanessa Zambelli: School of Medicine and Surgery, University of Milano-Bicocca, Monza, Italy; Mostafa Zanaty: Faculty of Nursing, Mansoura University COVID isolation hospital, Mansoura, Egypt; Andrea Zanoletti: Anesthesia and Intensive Care, Manerbio Hospital, Manerbio, Italy; Francesco Zarantonello: Institute of Anesthesia and Intensive Care, University Hospital of Padua, Padua, Italy; Michela Zardin: AOUI Anestesia e Rianimazione 1, S. Chiara, APSS Trento, Trento, Italia. UOC Anestesia e Rianimazione Ospedale valli del Noce Cles, APSS Trento, Trento, Italia; Francesca Zini: Anesthesia and Critical Care Unit, Ospedale di Belluno, Belluno, Italy; Eugenia Zoppellaro: Anesthesia and Intensive Care UOC, Pronto Soccorso, Anestesia e Terapie Intensive' Department, Aulss2 Treviso Hospital, Treviso, Italy.

Author contributions

All authors directly accessed and verified the underlying data reported in the manuscript, and accepted responsibility to submit for publication. All authors approved the final version of the manuscript to be published. All authors agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Substantial contributions to the conception or design of the work: SDR, NS, GL, LC, GF, EM, MP, DP, ER, PN. The acquisition, analysis, or interpretation of data for the work: GB, GF, AC, DG, AB, ME, EG, LG, AG, SMM, RR, IT, AZ. Drafting the work: SDR, NS, GL, AB, ME, LC, GF, EG, LG, AG, EM, MP, DP, RR, IT, AZ, ER. Reviewing it critically for important intellectual content: GB, GF, AC, DG, SMM, PN.

Funding

None to declare.

Availability of data and materials

Individual participant data that underlie the results reported in this article, after de-identification, data dictionary, study protocol, statistical analysis plan, informed consent form, and analytic code will be available to any researchers who provide a methodologically sound proposal, immediately following publication and without end date. Proposals should be directed to the corresponding author. To gain access, data requestors will need to sign a data access agreement.

Declarations

Ethics approval and consent to participate

The registry was designed following the Declaration of Helsinki and the study protocol was firstly approved by the Ethics Committee of the Saint Bortolo Hospital, Vicenza, Italy (Study ID Numbers: 22/21). Patient consent was obtained according to the national regulations of each participating Institution. In cases the patient was incompetent because of critical illness or the use of sedative or anesthetic drugs, consent could be delayed, and a provision for delayed consent was applied: as soon as competent, each patient was fully informed on what had been done, and a written permission of using data collected was obtained. The patients or their legal surrogates were informed of their right to request that the study procedures be discontinued and their right to refuse the study-related use of their medical records.

Consent for publication

Not applicable.

Competing interests

We declare no competing interests.

Author details

¹Centre for Medical Sciences-CISMed, University of Trento, Trento, Italy. ²UOC Anestesia e Rianimazione, AULSS8 Berica, Ospedale San Bortolo, Vicenza, Italy. ³Institute of Anesthesia and Intensive Care, Padova University Hospital, Padua, Italy. ⁴Department of Medicine and Surgery, University of Milano-Bicocca, San Gerardo Hospital, Monza, Italy. ⁵Department of Precision Medicine in Medical, Surgical and Critical Care (Me.Pre.C.C.), University of Palermo, Palermo, Italy. ⁶Department of Anesthesia, Intensive Care and Emergency, Policlinico Paolo Giaccone, Palermo, Italy. ⁷Unit of Biostatistics, Epidemiology and Public Health, Department of Cardiac, Thoracic and Vascular Sciences, University of Padova, Padua, Italy. ⁸Department of Medicine (DIMED), University of Padua, Via Vincenzo Gallucci 13, 35125 Padua, PD, Italy. ⁹Faculty of Medicine, University of Tripoli, Tripoli, Libya. ¹⁰Anesthesia and Intensive Care, Ospedale All'Angelo, Mestre, Italy. ¹¹Anaesthesia and Intensive Care, Department of Medical and Surgical Sciences, Magna Graecia University, Catanzaro, Italy. ¹²Dipartimento di Emergenza e Terapie Intensive, UOC di Anestesia e Terapia Intensiva Cardio-Toraco-Vascolare, Azienda Ospedaliera Universitaria Integrata di Verona, Verona, Italy. ¹³UOC Anestesia e Rianimazione, Ospedale di Vittorio Veneto, Vittorio Veneto, TV, Italy. ¹⁴University Department of Innovative Technologies in Medicine and Dentistry, Gabriele d'Annunzio University of Chieti-Pescara, Chieti, Italy. ¹⁵Department of Anesthesiology, Critical Care Medicine and Emergency, S.S. Annunziata Hospital, Chieti, Italy. ¹⁶Anaesthesia and Intensive Care Unit, Ospedali di San Donà di Piave e Jesolo, San Donà di Piave, Italy. ¹⁷Department of Anesthesia and Intensive Care, Santa Maria dei Battuti-Ca' Foncello Hospital, Treviso, Italy. ¹⁸Anesthesia and Critical Care Unit, Ospedale di Belluno, Belluno, Italy. ¹⁹UOC Anestesia e Rianimazione, Padova University Hospital, Padua, Italy. ²⁰Anesthesia and Intensive Care Unit, Manerbio Hospital, Manerbio, Italy.

Received: 8 November 2024 Accepted: 13 January 2025

Published online: 28 January 2025

References

- Guérin C, Albert RK, Beitler J, et al. Prone position in ARDS patients: why, when, how and for whom. *Intensive Care Med.* 2020;46(12):2385–96.

2. Guérin C, Reignier J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med*. 2013;368(23):2159–68.
3. Sud S, Friedrich JO, Adhikari NK, et al. Effect of prone positioning during mechanical ventilation on mortality among patients with acute respiratory distress syndrome: a systematic review and meta-analysis. *CMAJ*. 2014;186(10):E381–90.
4. Munshi L, DeSorbo L, Adhikari NKJ, et al. Prone position for acute respiratory distress syndrome. A systematic review and meta-analysis. *Ann Am Thorac Soc*. 2017;14(Suppl4):S280–8.
5. Bloomfield R, Noble DW, Sudlow A. Prone position for acute respiratory failure in adults. *Cochrane Database Syst Rev*. 2015;2015(11): CD008095.
6. Fan E, Del Sorbo L, Goligher EC, et al. An Official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine Clinical Practice Guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2017;195(9):1253–63.
7. Grasselli G, Calfee CS, Camporota L, et al. ESICM guidelines on acute respiratory distress syndrome: definition, phenotyping and respiratory support strategies. *Intensive Care Med*. 2023;49(7):727–59.
8. Qadir N, Sahetya S, Munshi L, et al. An update on management of adult patients with acute respiratory distress syndrome: an official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2024;209(1):24–36.
9. Duggal A, Rezoagli E, Pham T, et al. Patterns of use of adjunctive therapies in patients with early moderate to severe ARDS: insights from the LUNG SAFE Study. *Chest*. 2020;157(6):1497–505.
10. Li X, Scales DC, Kavanagh BP. Unproven and expensive before proven and cheap: extracorporeal membrane oxygenation versus prone position in acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2018;197(8):991–3.
11. Ferrando C, Suarez-Sipmann F, Mellado-Artigas R, et al. Clinical features, ventilatory management, and outcome of ARDS caused by COVID-19 are similar to other causes of ARDS. *Intensive Care Med*. 2020;46(12):2200–11.
12. Langer T, Brioni M, Guzzardella A, et al. Prone position in intubated, mechanically ventilated patients with COVID-19: a multi-centric study of more than 1000 patients. *Crit Care*. 2021;25(1):128.
13. De Rosa S, Sella N, Rezoagli E, et al. The PROVENT-C19 registry: a study protocol for international multicenter SIAARTI registry on the use of prone positioning in mechanically ventilated patients with COVID-19 ARDS. *PLoS ONE*. 2022;17(12): e0276261.
14. Pasin L, Sella N, Correale C, et al. Regional COVID-19 Network for Coordination of SARS-CoV-2 outbreak in Veneto. *Italy J Cardiothorac Vasc Anesth*. 2020;34(9):2341–5.
15. Zarantonello F, Sella N, Pettenuzzo T, et al. Early physiologic effects of prone positioning in COVID-19 acute respiratory distress syndrome. *Anesthesiology*. 2022;137(3):327–39.
16. Sinha P, Calfee CS, Beitler JR, et al. Physiologic analysis and clinical performance of the ventilatory ratio in acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2019;199(3):333–41.
17. Zheng M. Dead space ventilation-related indices: bedside tools to evaluate the ventilation and perfusion relationship in patients with acute respiratory distress syndrome. *Crit Care*. 2023;27(1):46.
18. Chen L, Chen GQ, Shore K, et al. Implementing a bedside assessment of respiratory mechanics in patients with acute respiratory distress syndrome. *Crit Care*. 2017;21(1):84.
19. Boscolo A, Sella N, Lorenzoni G, et al. Static compliance and driving pressure are associated with ICU mortality in intubated COVID-19 ARDS. *Crit Care*. 2021;25(1):263.
20. Tonetti T, Grasselli G, Rucci P, et al. Synergistic effect of static compliance and d-dimers to predict outcome of patients with COVID-19-ARDS: a prospective multicenter study. *Biomedicines*. 2021;9(9):1228.
21. Camporota L, Sanderson B, Chiumello D, et al. Prone position in COVID-19 and -COVID-19 acute respiratory distress syndrome: an international multicenter observational comparative study. *Crit Care Med*. 2022;50(4):633–43.
22. Rossi S, Palumbo MM, Sverzellati N, et al. Mechanisms of oxygenation responses to proning and recruitment in COVID-19 pneumonia. *Intensive Care Med*. 2022;48(1):56–66.
23. Zarantonello F, Andreatta G, Sella N, Navalesi P. Prone position and lung ventilation and perfusion matching in acute respiratory failure due to COVID-19. *Am J Respir Crit Care Med*. 2020;202(2):278–9.
24. Albert RK, Keniston A, Baboi L, Ayzac L, Guérin C, Proseva Investigators. Prone position-induced improvement in gas exchange does not predict improved survival in the acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2014;189(4):494–6.
25. Dilken O, Rezoagli E, Yartaş Dumanlı G, Ürkmez S, Demirkıran O, Dikmen Y. Effect of prone positioning on end-expiratory lung volume, strain and oxygenation change over time in COVID-19 acute respiratory distress syndrome: a prospective physiological study. *Front Med (Lausanne)*. 2022;2(9):1056766.
26. Giani M, Rezoagli E, Guervilly C, et al. Timing of prone positioning during venovenous extracorporeal membrane oxygenation for acute respiratory distress syndrome. *Crit Care Med*. 2023;51(1):25–35.
27. Gattinoni L, Vagginelli F, Carlesso E, et al. Decrease in PaCO₂ with prone position is predictive of improved outcome in acute respiratory distress syndrome. *Crit Care Med*. 2003;31(12):2727–33.
28. Morales-Quinteros L, Schultz MJ, Bringué J, et al. Estimated dead space fraction and the ventilatory ratio are associated with mortality in early ARDS. *Ann Intensive Care*. 2019;9(1):128.
29. da Cruz MR, Azambuja P, Torres KSC, Lima-Setta F, Japiassú AM, Medeiros DM. Identification and validation of respiratory subphenotypes in patients with COVID-19 acute respiratory distress syndrome undergoing prone position. *Ann Intensive Care*. 2024;14(1):178.
30. Okin D, Huang CY, Alba GA, et al. prolonged prone position ventilation is associated with reduced mortality in intubated COVID-19 patients. *Chest*. 2023;163(3):533–42.
31. Cornejo RA, Montoya J, Gajardo AJ, et al. Continuous prolonged prone positioning in COVID-19-related ARDS: a multicenter cohort study from Chile. *Ann Intensive Care*. 2022;12(1):109.
32. Carsetti A, Damia Paciarini A, Marini B, Pantanetti S, Adrario E, Donati A. Prolonged prone position ventilation for SARS-CoV-2 patients is feasible and effective. *Crit Care*. 2020;24(1):225.
33. Jochmans S, Mazerand S, Chelly J, et al. Duration of prone position sessions: a prospective cohort study. *Ann Intensive Care*. 2020;10(1):66.
34. Wendel-Garcia PD, Moser A, Jeitziner MM, et al. Dynamics of disease characteristics and clinical management of critically ill COVID-19 patients over the time course of the pandemic: an analysis of the prospective, international, multicentre RISC-19-ICU registry. *Crit Care*. 2022;26(1):199.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.