

Functional connectivity fingerprints of the frontal eye field and inferior frontal junction suggest spatial versus nonspatial processing in the prefrontal cortex

Orhan Soyuhos^{1,2}  | Daniel Baldauf¹ 

¹Centre for Mind/Brain Sciences (CIMEC), University of Trento, Trento, Italy

²Center for Neuroscience, University of California, Davis, California, USA

Correspondence

Daniel Baldauf, Center for Neuroscience, University of California, Davis, California, USA.

Email: daniel.baldauf@unitn.it

Funding information

Ministero degli Affari Esteri e della Cooperazione Internazionale; Fondazione Cassa Di Risparmio Di Trento E Rovereto

Edited by: Min Xu

Abstract

Neuroimaging evidence suggests that the frontal eye field (FEF) and inferior frontal junction (IFJ) govern the encoding of spatial and nonspatial (such as feature- or object-based) representations, respectively, both during visual attention and working memory tasks. However, it is still unclear whether such contrasting functional segregation is also reflected in their underlying functional connectivity patterns. Here, we hypothesized that FEF has predominant functional coupling with spatiotopically organized regions in the dorsal ('where') visual stream whereas IFJ has predominant functional connectivity with the ventral ('what') visual stream. We applied seed-based functional connectivity analyses to temporally high-resolving resting-state magnetoencephalography (MEG) recordings. We parcellated the brain according to the multimodal Glasser atlas and tested, for various frequency bands, whether the spontaneous activity of each parcel in the ventral and dorsal visual pathway has predominant functional connectivity with FEF or IFJ. The results show that FEF has a robust power correlation with the dorsal visual pathway in beta and gamma bands. In contrast, anterior IFJ (IFJa) has a strong power coupling with the ventral visual stream in delta, beta and gamma oscillations. Moreover, while FEF is phase-coupled with the superior parietal lobe in the beta band, IFJa is phase-coupled with the middle and inferior temporal cortex in delta and gamma oscillations. We argue that these intrinsic connectivity fingerprints are congruent with each brain region's function. Therefore, we conclude that FEF and IFJ have dissociable connectivity patterns that fit their

Abbreviations: dwPLI, debiased weighted phase lag index; FDR, false discovery rate; FEF, frontal eye field; FEF-IFJa, the contrasting functional connectivity patterns of FEF and IFJa; FEF-IFJp, the contrasting functional connectivity patterns of FEF and IFJp; HCP, human connectome project; ICA, independent component analysis; iCOH, imaginary part of the coherency; IFJa, anterior inferior frontal junction; IFJp, posterior inferior frontal junction; IPS, intraparietal sulcus; IT, inferior temporal; MEG, magnetoencephalography; MMP1, multimodal parcellation atlas; MRI, magnetic resonance imaging; oPEC, orthogonalized power envelope correlation; PDC, partial directed coherence; PFC, prefrontal cortex; plPFC, posterior lateral prefrontal cortex; rMEG, resting-state magnetoencephalography; ROI, region of interest; SCEF, supplementary and cingulate eye fields; SPC, superior parietal cortex; SPL, superior parietal lobule; VPA, ventral pre-arcuate. The abbreviations for the regions of interest in dorsal and ventral visual streams are presented in Table 1.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *European Journal of Neuroscience* published by Federation of European Neuroscience Societies and John Wiley & Sons Ltd.

respective functional roles in spatial versus nonspatial top-down attention and working memory control.

KEYWORDS

brain connectivity, magnetoencephalography, visual attention, visual pathways, working memory

1 | INTRODUCTION

At any moment, we face an overwhelming stream of rich information from the surrounding environment. Our brain must continuously re-evaluate all the incoming sensory stimulation to prioritize what is relevant to our behavioural goals (Baldauf & Deubel, 2010; Desimone & Duncan, 1995; Fecteau & Munoz, 2006). The prefrontal cortex (PFC), the maestro in orchestrating a range of cognitive functions in the human brain, plays an essential role in this respect (Duncan, 2001; Fuster, 2001; Miller & Cohen, 2001; Rossi et al., 2008). It crucially contributes to the central functions of selective attention and working memory by top-down modulating priority for incoming stimuli according to the context of the ongoing task demands (Jerde et al., 2012; Ptak, 2012). Its distinct control circuits' interareal communications with the rest of the cortex enable the brain to focus its limited perceptual resources on specific locations in space or on certain features or objects (Baldauf & Desimone, 2014; Bichot et al., 2019; Corbetta & Shulman, 2002; Gregoriou et al., 2009; Gross et al., 2004; Mangun, 1995; Siegel et al., 2008). In this context, human neuroimaging techniques and invasive recordings in primates suggest that the representational content encoded during visual selective attention and working memory can further segregate the functional specialization within PFC (Carrasco, 2011; Cavada & Goldman-Rakic, 1989; Giesbrecht et al., 2003; Goldman-Rakic, 1988; McCarthy et al., 1996; Paneri & Gregoriou, 2017; Romanski, 2004; Scialidhe et al., 1999; Wilson et al., 1993). In particular, two specific control sites located in the posterior lateral PFC (plPFC), namely, the frontal eye field (FEF) (Kelley et al., 2008; Petit & Pouget, 2019; Vernet et al., 2014) and the inferior frontal junction (IFJ) (Baldauf & Desimone, 2014; Brass et al., 2005; Derrfuss et al., 2005; Muhle-Karbe et al., 2014), show differential neural activity depending on the stimulus and task domain.

Experiments in which attention is covertly directed to a visuospatial target have led to increased neural activity in FEF, apart from its oculomotor function (Armstrong et al., 2009; Thompson et al., 2005; Thompson & Bichot, 2005). Additionally, muscimol injection and suprathreshold microstimulation techniques revealed FEF's crucial role in spatial attention (Moore &

Armstrong, 2003; Moore & Fallah, 2004; Wardak et al., 2006). In this respect, Thompson and Bichot et al. (2005) reported the existence of a visual salience map in primate FEF and noted that its initial neural activities are not selective for nonspatial features such as colour or shape, but that their activity evolves over time to select the target location. Along these lines, human neuroimaging and monkey studies indicated that FEF exerts top-down regulatory influences on the extrastriate visual cortex and increases the response gain to retinotopically corresponding targets directly or indirectly through its connections to the intraparietal sulcus (IPS) (Bressler et al., 2008; Gregoriou et al., 2009; Gregoriou, Gotts, & Desimone, 2012; Kelley et al., 2008; Merrikhi et al., 2017). Moreover, transcranial magnetic stimulation studies advanced these correlational findings with causal evidence and pointed out FEF's key function in the top-down control of visuospatial attention (Heinen et al., 2017; Marshall et al., 2015; Veniero et al., 2021).

In contrast, experimental designs that require attention to nonspatial features or object representations have consistently invoked neural responses in the inferior frontal cortex around IFJ. This evidence mainly accumulated from human neuroimaging studies due to the lack of corresponding sulcal landmarks in the nonhuman primate brain (Bedini & Baldauf, 2021; Donahue et al., 2018). Nevertheless, the studies on the ventral pre-arcuate of monkey PFC (VPA; regarded as the homolog of IFJ) have provided important insights into the region's functioning. For instance, Bichot et al. (2015) reported VPA as the source of feature selection. They showed that the inactivation of VPA reduces the feature-based modulation in ipsilateral V4 and impairs monkeys' performance in detecting target objects in the contralateral visual field (Bichot et al., 2019). Likewise, human neuroimaging and transcranial magnetic stimulation studies reported a top-down modulatory role of IFJ in object- and feature-based attention and working memory (Baldauf & Desimone, 2014; Meyyappan et al., 2021; Zanto et al., 2010, 2011; Zhang et al., 2018). For instance, Zanto et al. (2010) found that the functional coupling between the right IFJ and V4 was predictive of the extent of attentional modulation during attention to colour. Additionally, they demonstrated that the perturbation of IFJ weakens working memory performance for colour features (Zanto et al., 2011). Following

this, through directional phase coupling, Baldauf and Desimone (2014) additionally showed that IFJ is the source of attentional modulation of object representations in the inferior temporal (IT) cortex and the driver of non-spatial, object-based attention.

Overall, the evidence from task-based studies suggests that the encoding of to-be-attended or to-be-remembered representational contents functionally segregates PFC and reveals two distinct control circuits in the dorsal and ventral PFC: FEF and IFJ. While FEF is involved in the top-down control of spatial attention and working memory, IFJ contributes to the top-down control of nonspatial (feature- and object-based) attention and working memory (for a review, see Bedini & Baldauf, 2021). Nonetheless, it is still unclear to what extent this functionally specific contrast is reflected in these regions' underlying functional connectivity profiles.

1.1 | Research question and hypothesis

The current study aimed to analyse the intrinsic functional connectivity fingerprints of FEF and IFJ to support their functional segregation within PFC. We argue that FEF and IFJ's interareal communication with regions in the dorsal ('where') and ventral ('what') visual pathways (Goodale & Milner, 1992; Mishkin et al., 1983) would be a prerequisite for their functional involvement in spatial and nonspatial attention and working memory control, respectively; similarly to the structural connectivity fingerprints of areas in IT cortex being predictive of their functional selectivity (Osher et al., 2016; Saygin et al., 2011). In this respect, we hypothesize that FEF has predominant functional connectivity with spatiotopically organized areas in the dorsal visual stream, whereas IFJ has predominant functional connectivity with feature- or object-encoding areas in the ventral visual pathway. To this end, we performed region of interest (ROI) analyses and explored FEF and IFJ's functional connectivity to the rest of the brain. Additionally, we inquired about the dominant directionality of these interactions between FEF/IFJ and the visual streams. To this aim, we used various functional connectivity metrics (both phase- and power-based) and epoch lengths to consider their effects on the consistency of results.

2 | MATERIALS AND METHODS

2.1 | Participants

We benefited from the 1200 Subjects Release of Human Connectome Project (HCP) dataset that includes resting-state magnetoencephalography (MEG) recordings and

structural magnetic resonance imaging (MRI) of 95 healthy participants aged between 22 and 35 (Larson-Prior et al., 2013; Van Essen et al., 2013). The Washington University institutional review board approved data acquisition protocols for HCP. Each subject was asked for verbal consent during the initial telephone screening and required a signed consent document right before data acquisition. Thirty-six participants are reported to be monozygotic and dizygotic twins. Out of 95 subjects, we selected 55 independent ones (26 female) after randomly excluding one of the pairs from each twin and checking the data quality. The anonymized dataset is provided in ConnectomeDB (db.humanconnectome.org; Hodge et al., 2016).

2.2 | Procedure

Subjects typically spent 2 or 3 days completing the HCP protocol that includes behavioural tests, resting-state functional MRI, task-evoked functional MRI, diffusion-weighted MRI, and task-evoked MEG in addition to structural MRI and resting-state MEG (rMEG) sessions. MEG recordings always took place before MRI sessions to prevent the potential effect of residual magnetization caused by the MRI scanner on the superconducting MEG sensors (Van Essen et al., 2013).

2.3 | MRI data acquisition and preprocessing

Washington University—University of Minnesota Consortium of the Human Connectome Project performed structural MRI on a customized Siemens 3T 'Connectome Skyra' at Washington University (Uğurbil et al., 2013). They acquired two high-resolution T1-weighted (3D MPRAGE; .7 mm isotropic; FOV, 224 × 224; 256 sagittal slices; TR = 2400 ms; TE = 2.14 ms; inversion time [TI], 1000 ms; 8° flip angle) and T2-weighted (T2w; 3D T2-SPACE; .7 mm isotropic; FOV, 224 × 224; 256 sagittal slices; TR = 3200 ms; TE = 565 ms; variable flip angle) anatomical images of participants using a standard 32-channel Siemens receive head coil. Experts reviewed the quality of scans, considering the tissue contrast, blurriness and artefacts. If necessary additional scans were arranged for a second time. Also, a radiologist examined the existence of any neuro-anatomical anomalies. While the normal variations were noted, subjects with abnormal brain anatomy were excluded from the dataset.

ConnectomeDB provides preprocessed structural data resulting from the HCP structural pipeline (Glasser et al., 2013). The second part of this pipeline is mainly

based on a modified version of FreeSurfer's recon-all script to reconstruct a two-dimensional cortical surface from a three-dimensional volume (Dale et al., 1999; Fischl, Sereno, & Dale, 1999). We used the resulting files provided inside the Structural Extended Preprocessed package in the ConnectomeDB for the source reconstruction of MEG data.

2.4 | MEG data acquisition

The rMEG recordings were acquired on a whole-head Magnes 3600 scanner (4D Neuroimaging, San Diego, CA, USA) in a magnetically shielded room at Saint Louis University (Larson-Prior et al., 2013). The MEG scanner had 248 magnetometers and 23 reference channels (18 magnetometers and five first-order gradiometers). The data were recorded with a sampling rate of 2034.5101 Hz, delta encoded, and saved in a 4D file format. Electrooculography, electrocardiography and electromyography signals were also measured simultaneously to enable offline artefact rejection. Moreover, the experimenters used five head position indicators to track the head position in reference to the MEG sensors and marked the anatomical landmarks of nasion and the preauricular points to coregister the SQUID array to each individual's structural MRI. They used a Polhemus Fastrak-III system for the spatial digitization of localizer coils and fiducial points. Additional points defining the subject's head shape (about 2400 points) were also digitized to fine-tune the co-registration.

During the rMEG scanning, subjects lay in a supine position. They were instructed to keep their heads still while fixating a red crosshair at the center of the projection. Their head positions were recorded before and after every three consecutive sessions. Each run took approximately 6 min. Before rMEG, the empty room and participant noise recordings were also acquired to monitor the sensor noise and the noise stemming from the participant due to residual magnetization effects.

2.5 | MEG data preprocessing

The ASCII text files that annotate the indexes of bad channels, bad segments and non-brain components are provided inside the Preprocessed Resting-State package in the ConnectomeDB. To ensure the reproducibility of our results, we decided to utilize them for preprocessing rMEG data. First, the noisy channels and the bad data segments were marked following a visual inspection. Later, the correlation between signals from adjacent sensors and the ratio of their variance to their neighbours were calculated to mark the outliers as bad

channels. Additionally, z-scores for all time points in each channel were computed to identify the segments whose score was higher than a given threshold. Next, the clipping and muscle-related artefacts were detected using `ft_artifact_clip.m` and `ft_artifact_muscle.m` functions from the FieldTrip toolbox (Oostenveld et al., 2010). Furthermore, an iterative independent component analysis (ICA) was performed to detect faulty channels and bad segments in the raw data by using the list of bad channels and segments identified beforehand as initial input. In subsequent iterations of this ICA, the input was updated by the bad sensors or segments found in the previous iteration until the ICA algorithm could no longer identify any further bad channels/segments.

In the next step of the MEG preprocessing pipeline, ICA was used this time to remove the artefacts related to eye movements, blinks, and heartbeats from the signal. To this end, first, a bandpass (1.3–150 Hz) and a notch Butterworth (59–61 and 119–121 Hz) filter were applied to the MEG, electrocardiography and electrooculography data. Next, the previously identified bad channels, and segments were cut out from these three recordings. Afterward, ICA was implemented on the preprocessed MEG data 20 times based on different seed values. The power spectral densities and power time courses were calculated for the independent components of each repetition and corresponding time series of electrocardiography and electrooculography channels. An automatic classification algorithm detected artefacts based on the correlation between independent components and electrocardiography/electrooculography recordings. It classified independent components into the brain and non-brain components while also considering pink noise and the noise stemming from the power supply (60 and 120 Hz). The classifier's performance was later confirmed by visual inspection. Ultimately, the best iteration with the highest ratio of brain to artefact components from 20 repetitions was selected. The indices of its brain and non-brain components were reported in ASCII text files. After the removal of the bad channels, bad segments and artefacts from the rMEG data, we parsed the continuous times series into non-overlapping epochs of either 2, 5 or 10 s to consider the epoch length's effect on the connectivity measures (Brookes et al., 2011; Fraschini et al., 2016).

2.6 | Source reconstruction of MEG signals

For each epoch length, we utilized the Brainstorm toolbox (Tadel et al., 2011) to reconstruct source-level activation on the native cortex of each subject. We first

corrected the DC offset of preprocessed rMEG data and downsampled it to 300 Hz. To estimate sensor noise, we computed the noise covariance matrix from raw empty room recording after removing its DC bias. Additionally, we calculated the lead field matrix for an evenly distributed grid of 15,002 cortical sources by fitting a single sphere under each sensor (Huang et al., 1999). Based on these gain and noise covariance matrices, we applied the minimum-norm estimation method to inverse-model activations in source space (Baillet et al., 2001; Dale et al., 2000). Here, we chose minimum-norm estimation over beamformers because it was shown to perform better for spatially distant sources (Hincapié et al., 2017). Next, the source level activity for each subject was projected to an MNI average brain template (FreeSurfer fsaverage; Fischl, Sereno, Tootell, & Dale, 1999) and parcellated by the multimodal parcellation atlas (MMP1; Glasser et al., 2016). Finally, we calculated the scout activations for 360 regions by taking the maximum absolute value across all vertices per scout and multiplying with its sign. We saved them as Fieldtrip structures to perform functional connectivity analyses in the next step.

The MEG Anatomy package provided in the ConnectomeDB enables the automatic co-registration of MEG sensors with the structural MRI. However, it does not allow projecting sources to a standard brain (such as fsaverage). Because we aimed to interpolate source-level activation of each subject onto the fsaverage template for the group-level analysis, we used the pial surface provided in the Structural Extended Preprocessed package to reconstruct sources. In this way, we could also increase the resolution of the cortex from 8004 to 15,002 vertices for a better representation of cortical folds. However, we still had to use the MEG Anatomy package to retrieve the coordinates of fiducial points for each subject because the

individual MRIs were defaced, and the fiducial points were not stated explicitly.

2.7 | Functional connectivity and directionality analyses

In the second step, we conducted seed-based functional connectivity analyses with seed regions of FEF, anterior IFJ (IFJa), and posterior IFJ (IFJp) in both hemispheres by employing the FieldTrip toolbox (Oostenveld et al., 2010) (Figure 1). We first carried out frequency analyses on time series data over all epochs while using the multiple tapers based on the Slepian sequence. The frequencies of interest were the delta (1–4 Hz), theta (4–8 Hz), alpha (8–13 Hz), beta (13–30 Hz) and gamma (30–100 Hz) bands with the number of spectral smoothing boxes of 1, 1, 1, 2 and 10, respectively. We computed complex Fourier spectra between all brain parcels for each frequency band. Based on this Fourier representation of data, we calculated the functional connectivity between each of the six seed regions and the rest of the 359 brain parcels. We decided to use both phase- and amplitude-based functional connectivity measures to take into account the possible distinct roles of phase- and amplitude-coupling (Daffertshofer et al., 2018; Siems & Siegel, 2020). In particular, we chose the imaginary part of the Coherency (iCOH) (Nolte et al., 2004), debiased weighted phase lag index (dwPLI) (Vinck et al., 2011), and orthogonalized power envelope correlation (oPEC) (Hipp et al., 2012) because these metrics are corrected for spatial leakage and have higher group-level repeatability compared with other methods (Bastos & Schoffelen, 2016; Brookes et al., 2011; Colclough et al., 2016; Duan et al., 2021).




FIGURE 1 The locations of seed regions from the MMP1 atlas for the functional connectivity analyses.

We performed specific ROI and whole-brain exploratory analyses based on the connectivity metrics mentioned above. For ROI analyses, the target MMP1 parcels in Table 1 were chosen based on a systematic bibliographical search of the relevant literature (Felleman & Van Essen, 1991; Glasser et al., 2016; Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Mishkin et al., 1983) and the list reported in Zimmermann et al. (2018). We first selected the mid-higher visual areas that belong to the dorsal or ventral visual pathway. Additionally, we used the retinotopy atlas by Wang et al. (2015) to identify putative retinotopic parcels of the MMP1 atlas and include them in our definition of the dorsal visual stream, as the topographic organization is generally considered a defining feature of this stream. We excluded parcels near the medial wall, parcels from the superior temporal lobe, and the temporoparietal junction because they seem to be characterized by distinctive connective profiles (Kravitz et al., 2011, 2013; Mars et al., 2011). This led to a total of 33 regions with specific visual properties and selectivity to visual features (Table 1). Next, we tested whether the spontaneous activity of each parcel in the ventral and dorsal visual pathway has higher functional connectivity with FEF or IFJ. Following this, we additionally questioned whether the regions that show preferential connectivity, neither with FEF nor with IFJ, have a statistically significant intrinsic functional connectivity with any of these seed regions when tested against the mean of whole-brain connectivity values per frequency band.

Finally, we analysed the dominant directional interactions between the seed regions in PFC and the target areas in the visual cortex during the resting state. To this end, we conducted directionality analyses between FEF, IFJa, IFJp, and all the ROIs in Table 1. First, we computed partial directed coherence (PDC; Baccalá & Sameshima, 2001) of each unidirectional interaction between seed regions and visual streams. Then, for each pair, we subtracted the PDC values of one unidirectional connection from the other and tested its significance against zero. We chose PDC from a range of directionality measures because it mainly considers direct interactions between regions, is generally regarded as insensitive to source leakage and shows high group-level repeatability, for instance, in comparison with the phase slope index (Colclough et al., 2016).

2.8 | Statistical analyses

We implemented paired Wilcoxon signed-rank tests for each type of analysis because the obtained connectivity values had a non-Gaussian distribution. For the same

reason, we did not apply z -transformation to the correlation matrix resulting from the oPEC metric. Additionally, we used the false discovery rate (FDR) method to correct for multiple comparisons (Benjamini & Hochberg, 1995; Groppe, 2021). In the hypothesis-driven ROI analyses, we corrected multiple comparisons for 16 ventral and 17 dorsal regions in Table 1 (+3 seed regions for PDC measure). In contrast, we constrained the significance test for the whole-brain exploratory analyses to one hemisphere and accordingly corrected multiple comparisons for 180 regions. To visualize FEF and IFJ's predominant functional connectivities across frequency bands or hemispheres, we converted the adjusted p values to z -scores. We summed the z -score of each parcel and divided them into the number of times they were significant. Importantly, we applied paired significance tests separately for FEF, IFJa and IFJp per region. We did not average the connectivity values of IFJa and IFJp before statistical analyses because we suspected them of having different functional connectivity fingerprints based on previous evidence (Baker et al., 2018; Bedini & Baldauf, 2021).

2.9 | Ground truth analyses

We performed ground truth analyses to ensure that our analysis pipeline was implemented correctly and provided valid results. First, we replicated the results from Hipp et al. (2012) paper for the oPEC metric. In the beta band (13–30 Hz), we calculated the power correlation of the left and right MT with the rest of the brain. We could successfully replicate area MT's seed-based connectivity and reproduce their paper's third figure by using a different dataset, source-reconstruction method (minimum norm estimation rather than beamforming), and our own processing pipeline. We did not apply any statistical mask to the figure shared in the Supporting information (Figure S1). Additionally, as it has been done in the resting MRI data of non-human primates, we tested the existence of a functional coupling between the supplementary eye field and FEF. We computed the seed-based connectivity of area supplementary and cingulate eye fields (SCEF) and observed its ipsilateral power and phase coupling with the right FEF, specifically in the beta band (one-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected for 180 ROIs) (Figure S23). Notably, the best match between fMRI and MEG functional connectivity findings has also previously been reported in the beta band (Hall et al., 2014). This precise functional connectivity between the right FEF and SCEF in the beta band further confirms the reliability of our findings.

TABLE 1 The list of parcels from the MMP1 atlas for ROI analyses.

Parcel	Description	Visual stream	Topographical organization
Dorsal visual pathway			
V6	Sixth visual area	Dorsal (Glasser et al., 2016)	✓ (Kravitz et al., 2011)
V6A	Area V6A	Dorsal (Glasser et al., 2016; Zimmermann et al., 2018)	✓ (Kravitz et al., 2011; Wang et al., 2015)
V7	Seventh visual area	Dorsal (Glasser et al., 2016)	✓ (Glasser et al., 2016; Wang et al., 2015)
IPS1	Intraparietal sulcus area 1	Dorsal (Glasser et al., 2016; Zimmermann et al., 2018)	✓ (Wang et al., 2015)
IP1	Area intraparietal 1	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
MIP	Medial intraparietal area	Dorsal (Zimmermann et al., 2018)	✓ (Kravitz et al., 2011; Wang et al., 2015)
VIP	Ventral intraparietal complex	Dorsal (Zimmermann et al., 2018)	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
LIPd	Area lateral intraparietal dorsal	—	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
LIPv	Area lateral intraparietal ventral	Dorsal (Zimmermann et al., 2018)	✓ (Felleman & Van Essen, 1991; Kravitz et al., 2011; Wang et al., 2015)
7AL	Lateral area 7A	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
7PC	Area 7PC	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
7PL	Lateral area 7P	Dorsal (Zimmermann et al., 2018)	✓ (Wang et al., 2015)
V3A	Area V3A	Dorsal (Glasser et al., 2016; Goodale & Milner, 1992)	✓ (Felleman & Van Essen, 1991; Glasser et al., 2016; Kravitz et al., 2013; Wang et al., 2015)
V3B	Area V3B	Dorsal (Glasser et al., 2016)	✓ (Wang et al., 2015)
V3CD	Area V3CD	—	✓ (Wang et al., 2015)
LO3	Area lateral occipital 3	—	✓ (Wang et al., 2015)
MT	Middle temporal area	Dorsal (Goodale & Milner, 1992)	✓ (Felleman & Van Essen, 1991; Glasser et al., 2016; Kravitz et al., 2013)
Ventral visual pathway			
V8	Eighth visual area	Ventral (Glasser et al., 2016)	—
VMV1	Ventromedial visual area 1	Ventral (Glasser et al., 2016)	—
VMV2	Ventromedial visual area 2	Ventral (Glasser et al., 2016)	—
VMV3	Ventromedial visual area 3	Ventral (Glasser et al., 2016)	—
PHT	Area PHT	Ventral (Zimmermann et al., 2018)	—
PH	Area PH	Ventral (Zimmermann et al., 2018)	—
TE1a	Area TE1 anterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	—
TE1m	Area TE1 middle	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	—
TE1p	Area TE1 posterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	—
TE2a	Area TE2 anterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	—

(Continues)

TABLE 1 (Continued)

Parcel	Description	Visual stream	Topographical organization
TE2p	Area TE2 posterior	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011, 2013; Zimmermann et al., 2018)	—
TF	Area TF	Ventral (Goodale & Milner, 1992; Kravitz et al., 2011; Zimmermann et al., 2018)	—
TGv	Area TG ventral	Ventral (Kravitz et al., 2013; Zimmermann et al., 2018)	—
FFC	Fusiform face complex	Ventral (Glasser et al., 2016)	—
PIT	Posterior Infero temporal	Ventral (Glasser et al., 2016)	—
VVC	Ventral visual complex	Ventral (Glasser et al., 2016)	—

Note: We selected region of interests in the visual cortex as belonging to the ventral or dorsal visual stream by considering previous arguments for their inclusion in the dorsal/ventral pathway, receptive field sizes, and the existence of a topographical organization.

3 | RESULTS

We first report the outcomes for 2-s epoch segmentation and ipsilateral functional connectivity. We detail the results for ROI analyses with respect to phase- and power-based functional connectivity metrics. Additionally, we present PDC in the resting state to expand on the direction of the interaction. While the contrasting functional connectivity patterns of FEF versus IFJa ('FEF-IFJa') are presented in Figures 2 and 3, the illustrations for the contrasting functional connectivity patterns of FEF versus IFJp ('FEF-IFJp') are available in the Figures S2–S3. Moreover, the supplementary document provides all three seed regions' intrinsic functional connectivity profiles against the mean (Figure S4), the findings for 5- and 10-s epoch segmentations (Figures S5–S6), contralateral connectivity (Figure S7) and whole-brain connectivity analyses (Figures S18–22). The abbreviations for the cortical area names throughout the paper are mainly based on Glasser et al. (2016).

3.1 | Predominant functional connectivities

All three connectivity metrics demonstrated that FEF has robust functional connectivity with the regions in the dorsal visual stream, whereas IFJa has stronger functional connectivity with the areas in the ventral visual stream independently of lateralization and frequency band (Figure 2; two-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected). The direct contrast between the ipsilateral connectivity fingerprints of FEF and IFJa revealed that FEF has a robust power correlation with the dorsal visual stream in beta and gamma bands. In contrast, IFJa has a strong power coupling with the

ventral visual stream in delta, beta and gamma oscillations. Moreover, both iCOH and dwPLI metrics showed that FEF is phase coupled with the superior parietal lobe (SPL) and IPS in the beta band, while IFJa is phase coupled with the inferior and middle temporal cortex by delta and gamma oscillations (Figure 3). Importantly, the anterior and posterior parts of IFJ have different connectivity patterns. Although the direct contrast between the ipsilateral connectivity fingerprints of FEF and IFJp also indicates FEF's higher functional connectivity to SPL and IPS, we observed that IFJp has weaker functional connectivity to the middle and ventromedial temporal cortex, particularly in the alpha and beta frequencies. Nevertheless, it still achieves a higher power correlation with the ventral visual areas in the delta frequency compared with FEF (Figures S2 and S3).

The phase-based metrics showed that the significant parcels for FEF-IFJa are lateralized in the left hemisphere. The iCOH and dwPLI metrics did not reveal any predominant coupling for FEF or IFJa with the selected ROIs in the right hemisphere. On the contrary, while the strong power correlation for FEF-IFJa was evident in both hemispheres, the predominant power coupling between the left IFJa and ventral visual stream was weaker than in the right hemisphere (Figure S3). Additionally, we observed an apparent dissociation within the higher frequencies for the power coupling between visual streams and pPPFC in the right hemisphere. The right FEF and IFJa functionally segregate the right visual pathways into dorsal and ventral streams in beta and gamma oscillations. Overall, the results from both phase- and power-based metrics illustrate that FEF and the areas in the dorsal visual stream predominantly communicate in beta frequency. In contrast, IFJa and the regions in the ventral visual stream are functionally connected through delta and gamma oscillations.




FIGURE 2 The predominant functional connectivity maps of frontal eye field (FEF) versus anterior inferior frontal junction (IFJa) across frequency bands and hemispheres for (A) oPEC, (B) iCOH and (C) dwPLI metrics (two-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected for 33 ROIs). The results shown here are based on 2-s epoch segmentation. Figures S8–S15 demonstrate the distribution of FEF–IFJa’s values for each region of interest (ROI).




FIGURE 3 The frequency-specific predominant functional connectivities for frontal eye field–anterior inferior frontal junction (FEF–IFJa) and FEF–posterior IFJ (IFJp). While the contrasting functional connectivity fingerprints of FEF versus IFJa are shown with darker red and blue colours, the contrasting functional connectivity fingerprints of FEF versus IFJp are illustrated with brighter red and blue colours, respectively. The width of a line reflects its relative z-score of the connectivity to other nodes (two-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected for 33 regions of interest [ROIs]). The results shown here are based on 2-s epoch segmentation (Kassebaum, 2021).

For FEF-IFJp, we observed predominant phase coupling only for FEF in the alpha and beta bands, in the left and right hemispheres, respectively (Figure S3). Furthermore, the iCOH and dwPLI measures showed that areas PHT, TE1m and VMV1 in the ventral stream have a more robust phase relation with FEF than IFJp. Like the contrast between FEF and IFJa, a predominant power correlation exists in both hemispheres also for FEF-IFJp. However, we cannot see any predominant power coupling between IFJp and the ventral visual stream in higher frequencies. Also, the left VMV2 in the ventral visual pathway has a higher power correlation with the left FEF rather than IFJp. Finally, IFJa and IFJp's contrasting functional connectivity patterns in the ventral visual stream revealed that the ventral regions have robust phase and power coupling with IFJa than IFJp independently of lateralization and frequency band (Figure 4; two-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected). IFJa has a predominant phase relation with areas PH, TE1a, TE1m and TE2p. Besides, it has a stronger power correlation with areas V8, VMV2, VMV3, PH, TE1a, TE1m, TE1p, TE2a, TE2p, TF, TGv, FFC and VVC compared with IFJp.

Noticeably, areas V7, V3B, MT, V8, PH, FFC, PIT and VVC do not show predominant connectivity either with FEF, IFJa or IFJp (Figures 2 and S2). In this respect, our additional analyses for FEF, IFJa and IFJp's seed-based intrinsic functional connectivities (one-sided Wilcoxon signed-rank test against the average, $p < .05$, FDR-corrected) revealed that V7, V3B, MT, V8 and PIT do not have statistically significant power or phase coupling with any of these three seed regions (Figure S4). Meanwhile, PH has a significant power correlation both with FEF and IFJa. FFC is significantly power-coupled with IFJa and IFJp, but not FEF. VVC has significant phase coupling with IFJa. Overall, we observed that all three

seed regions have a robust power correlation and statistically significant spontaneous phase coupling with the parcels in SPL, IPS and the ventral visual stream. In particular, the functional coupling between area 7PC and the seed regions is highly significant for both phase- and power-based metrics.

3.2 | Influences of chosen epoch length

We observed that 2-, 5-, and 10-s epoch segmentation affects distinct frequency bands and connectivity metrics differently. In the delta band, IFJa has a consistent robust power correlation with the ventral visual stream. However, its dominance in the ventral pathway decreases in beta and gamma oscillations with increasing segmentation length. In contrast, FEF's power coupling with the dorsal visual stream is consistent in beta and gamma frequencies (Figure S5). Additionally, the results for iCOH are less consistent as the epoch length increases and the number of trials decreases per recording session (Figure S6). Nevertheless, specific predominant functional couplings stand out across different epoch segmentation and connectivity measures: FEF has robust functional connectivity with the areas VIP, LIPd, 7AL and 7PC in the dorsal visual stream in higher frequencies whereas IFJa is predominantly coupled with the areas TE1, TE2, TF and TGv in the ventral visual stream in lower frequencies.

3.3 | Directional interactions between visual streams and pIPFC

Directionality analyses in the frequency domain revealed intrinsic unidirectional information flows between seed




FIGURE 4 The predominant functional connectivity maps of IFJa versus IFJp in the ventral visual stream, across frequency bands and hemispheres for iCOH and oPEC metrics (two-sided Wilcoxon signed-rank test, $p < .05$, FDR-corrected for 16 ROIs). The results shown here are based on 2-s epoch segmentation.




FIGURE 5 The partial directed coherence between frontal eye field (FEF), inferior frontal junction (IFJ) and visual streams. the figure shows the directional influences between seed regions (FEF, anterior IFJ [IFJa] and FEF) and visual streams. While the green regions receive inputs, the magenta ones send outputs to a seed region (two-sided Wilcoxon signed-rank test, $p < .001$, FDR-corrected for 33 + 3 regions of interest [ROIs]). The results shown here are based on 2-s epoch segmentation.

regions and the visual streams (Figure 5; two-sided Wilcoxon signed-rank test, $p < .001$, FDR-corrected). Most remarkably, PDC showed that directional interactions from SPL and IPS to seed regions are significantly more potent than in the other direction. In contrast, directional influences from seed regions to the middle and IT cortex are more robust than in the other direction, mainly in delta, theta and beta frequency bands. In other words, the rMEG time series of the regions in plPFC is predicted by the intrinsic neural activity of the regions in the superior parietal cortex (SPC; SPL and IPS) and themselves predictive of the spontaneous activity of the areas in the ventral visual stream. In the alpha band, we instead observed that the seed regions, particularly IFJa and IFJp, predict the intrinsic neural activity in both visual streams except areas 7AL, 7PL and VIP in the dorsal stream. The direction of information flow from seed regions to these areas located in the medial part of SPC is

not significantly stronger than the other direction for any oscillatory band. Likewise, no region in the ventral stream has a directional influence over any seed region, except left TE1m in the gamma band.

3.4 | Directional interactions among FEF, IFJa and IFJp

Lastly, the directionality analyses between FEF, IFJa, and IFJp indicated a direction of information flow from IFJp to the other two seed regions (Figure 6). The PDC values from IFJp to IFJa and IFJp to FEF are statistically stronger than in the other directions in both hemispheres. While left IFJp feeds information to left FEF and left IFJa mainly in the theta band, the same direction of interaction in the right hemisphere is driven through both alpha and beta oscillations. Notably, the directed interaction




FIGURE 6 The partial directed coherence between the seed regions within prefrontal cortex (PFC). The figure shows the directional influences among frontal eye field (FEF), anterior inferior frontal junction (IFJa) and posterior inferior frontal junction (IFJp) in both hemispheres. The width of the arrows reflects z-scores on top of them (two-sided Wilcoxon signed-rank test, FDR-corrected for 33 + 3 ROIs; see Figure 5). The distribution of subjects' PDC value for each significant directional influence is illustrated in Figures S16 and S17 (Bechtold, 2016).

from the right IFJp to the right FEF in the alpha band is highly significant ($p < .00001$; Figure S16). Besides, we observe that right FEF and IFJa have reciprocal connections in various frequency bands. The direction of spontaneous activity from FEF to IFJa and IFJa to FEF are subserved by delta and alpha oscillations, respectively. In the left hemisphere, we only see unidirectional interactions from FEF to IFJa in the delta and gamma bands but not vice versa. Noticeably, the delta frequency band underlies FEF's directional influence over IFJa in both hemispheres ($p < .005$).

4 | DISCUSSION

The current study investigated the intrinsic functional connectivity profiles of FEF and IFJ, two specific control circuits in the dorsal and ventral PFC, to provide additional evidence for their respective top-down functions in spatial versus nonspatial (feature and object-based) attention and working memory control. Our results demonstrated that FEF has a more robust power and phase coupling with the spatiotopically organized regions in the dorsal visual stream. In contrast, IFJa has more substantial power and phase relation with the feature or object encoding regions in the ventral visual pathway (Figure 2). These results suggest that in humans, the 'where' and 'what' visual streams (Goodale & Milner, 1992; Mishkin et al., 1983) might extend all the way into plPFC (Goldman-Rakic, 1996; Macko et al., 1982; Wilson et al., 1993). The spatial versus nonspatial to-be-attended or to-be-remembered stimulus domain might respectively segregate plPFC into functionally specific dorsal versus ventral visual pathways.

4.1 | FEF versus IFJa

FEF has robust functional connectivity with the areas VIP, LIPd, 7AL and 7PC across different epoch lengths and connectivity metrics in the dorsal stream. In this respect, studies on monkeys reported that area VIP is involved in the analysis of visual motion (Colby et al., 1993; Duhamel et al., 1998), multisensory space representation (Schlack et al., 2005), visuomotor control (Culham et al., 2006), self-motion (Bremmer, 2005; Chen et al., 2011; Schlack et al., 2002) and motion encoding in near-extraperosnal space (Schlack et al., 2003). While area LIP is also to some degree selective to shape (Serenio & Maunsell, 1998), researchers at large showed its role in spatial attention (Bisley & Goldberg, 2003; Thakral & Slotnick, 2009), the choice of saccade target (Andersen et al., 1990; Barash et al., 1991; Blatt et al., 1990; Kusunoki et al., 2000; Kusunoki & Goldberg, 2003) and the creation of priority maps (Gottlieb, 2007; Ipata et al., 2009; Sprague & Serences, 2013). Finally, it is evident that SPL (areas 7AL and 7PC) takes part in shifting spatial attention (Behrmann et al., 2004; Caspari et al., 2018; Corbetta et al., 1995; Kelley et al., 2008; Molenberghs et al., 2007; Spadone et al., 2021; Vandenberghe et al., 2001; Yantis et al., 2002) and encoding spatial coordinates for planned reach movements (Baldauf et al., 2008; Connolly et al., 2003). Besides, these results match FEF's deterministic tractography (Baker et al., 2018). FEF's white matter connections accordingly terminate at the intraparietal sulcus, nearby SPL.

In contrast to FEF, IFJa has a stronger functional coupling with the areas TE1a, TE2a, TF and TGv in the ventral visual stream. These areas roughly correspond to

the anterior middle temporal gyrus (area TE1a), infero-temporal cortex (areas TE2a and TF), and temporal pole (area TGv). Notably, studies associated the anterior middle temporal gyrus with social cognition (Xu et al., 2020), language and semantic processing (Ferreira et al., 2015; Gorno-Tempini & Price, 2001; Rossell et al., 2001; Siuda-Krzywicka et al., 2021; Walker et al., 2011). On this note, this particular functional coupling with the anterior temporal cortex might be due to mind-wandering during the resting-state scan (Chou et al., 2017). In comparison, lesions and neuroimaging studies suggest that the temporal pole takes part in face recognition by associating person-specific stored memories with the perceptual representation of people's faces (de Vries & Baldauf, 2019; Kamps et al., 2019; Landi et al., 2021; Olson et al., 2007). Additionally, a relational processing task adapted from Smith et al. (2007) revealed that, in contrast to TGd, TGv is activated vs deactivated in relational primary contrasts when subjects were asked to differentiate objects based on their shapes or textures (Glasser et al., 2016). Lastly, the IT cortex has a direct role in object recognition (Kreiman et al., 2006; Lehky & Tanaka, 2016; Tanaka, 1996; Yamane et al., 2008). Studies on monkeys and humans both show the IT cortex's involvement in object-based attention and working memory (Baldauf & Desimone, 2014; Kar et al., 2019; Miller et al., 1996). Furthermore, IFJa, but not FEF, has distinctive white matter connectivity with the temporal cortex, which is in accordance with its predominant functional coupling reported here. IFJa's deterministic tractography demonstrates its structural connectivity to the middle and temporal gyrus (Baker et al., 2018). Also, IFJ's probabilistic diffusion-weighted imaging tractography is especially well connected with several functional areas in the IT cortex, which was not the case for FEF (Baldauf & Desimone, 2014). Overall, FEF and IFJa's respective predominant functional coupling with SPC and the temporal cortex in the resting state is congruent with their underlying structural connectivity and respective functional specialization in task-based studies.

It is important to note that FEF has statistically significant functional coupling with the areas in the dorsal visual stream when its connectivity contrasted against the mean of the whole-brain connectivity matrix for each frequency band. Also, IFJa and IFJp have significant coupling with the dorsal areas when contrasted against the respective average connectivity values (Figure S4). That is to say, FEF and IFJ do not have exclusive functional connectivity with the areas in the dorsal or ventral visual pathways. Similarly, some studies indicated FEF and IFJ's role in spatial and nonspatial tasks, respectively (Asplund et al., 2010; Heinen et al., 2014; Liu, 2016; Liu et al., 2011; Tamber-Rosenau et al., 2018; Zhou & Desimone, 2011).

For instance, Liu (2016) could decode the attended object from the neural activity of FEF in a nonspatial attention task, whereas Tamber-Rosenau et al. (2018) found transient activity in IFJ during a spatial attention task. Moreover, Heinen et al. (2014) provided causal evidence for FEF's role in feature-based attention by applying offline transcranial magnetic stimulation. However, these studies did not directly compare the predominant involvement of IFJ versus FEF. These regions might have transitive relations with either visual stream through each other's functional coupling. In this context, our results rather show that FEF and IFJa have dominant intrinsic relations with the dorsal and ventral regions, respectively, when their functional connectivities are contrasted against each other. Together with the previous evidence discussed in the previous two paragraphs, our results suggest that FEF and IFJa are predominantly responsible for spatial versus nonspatial information processing in the pPPFC during visual attention and working memory tasks.

4.2 | Frequency-specific interactions

Different oscillations are associated with different cognitive functions (Başar et al., 2001; Güntekin & Başar, 2016; Harmony et al., 1996). Thanks to the high temporal and spatial resolution of MEG, we could reveal the frequency-specific signatures of the functional coupling between FEF, IFJ, and visual pathways (Figure 3). While these frequency-specific interactions in resting state data hint at general communication patterns between the involved areas, it is important to bear in mind that those interactions can rapidly change either in strength, dominant frequency band, and/or directionality as specific attentional or working memory tasks evolve.

In the resting state, we found substantial delta oscillatory coupling between IFJ and the areas in the ventral visual stream. In this respect, Harmony et al. (1996) observed that attention to internal representations leads to increased delta power during mental processing, such as the Sternberg paradigm that measures working memory performance. They suggested that delta oscillations carry attentional modulations from the frontal cortex to distant neuronal networks (Harmony, 2013) such that it might have a role in suppressing internal or external distractors that interfere with the ongoing mental state. Noticeably, these statements align with IFJ's role in cognitive control (Brass et al., 2005; Derrfuss et al., 2005) and selective attention (Baldauf & Desimone, 2014; Bedini & Baldauf, 2021). Also, for example, Zarjam et al. (2011) could successfully classify working memory load levels based on the delta activity in frontal electroencephalography channels. Moreover, they linked delta

synchronization to subjects' internal concentration. Because functional coupling between distant regions requires a slower 'clock' (Jensen & Colgin, 2007), delta oscillations might subserve the intrinsic functional connectivity between IFJ and the areas in the ventral visual stream in the absence of attention to features or objects. This baseline information flow can potentially pave the way for task-evoked activity (Cole et al., 2016) during selective attention and working memory.

In the theta band, we did not find any significant region that predominantly power correlates with FEF or IFJ. Additionally, phase-based measures did not reveal any consistent results. The lack of predominant coupling in the theta band might be because theta oscillations are instead linked to learning and memory through cortico-hippocampal interactions (Başar et al., 2001; Buzsáki, 2002; Goutagny et al., 2009; Herweg et al., 2020; Lega et al., 2012; Lubenov & Siapas, 2009; O'Neill et al., 2013; Siapas et al., 2005). However, it is important to note that several task-based studies linked theta oscillation to visual working memory (Liebe et al., 2012; Muthukrishnan et al., 2020; Sauseng et al., 2010) and willed attention (Rajan et al., 2019).

On the other side, we found predominant functional connectivity between FEF and SPC (areas VIP, LIPd, 7AL and 7PC) in the alpha band. Some of this alpha band activity in the dorsal connections might be attributed to general arousal (Barry et al., 2020; Brancaccio et al., 2020; Cantero et al., 1999; Pivik & Harman, 1995) or visual mental imagery (Xie et al., 2020) during resting state. However, alpha oscillations are also known as neural signatures in task-based attention studies (Babiloni et al., 2006; Bae & Luck, 2019; Bagherzadeh et al., 2020; Capotosto et al., 2009; de Vries et al., 2019; de Vries et al., 2021; Geng, 2014; Klimesch et al., 2007; Mazaheri et al., 2011; Noah et al., 2020; Rihs et al., 2007; Sauseng et al., 2005; Thut et al., 2006; Toscani et al., 2010; van Diepen et al., 2016).

Beta synchronization, which we found more strongly involved in the dorsal connections of FEF, is associated with motor inhibition (Gilbertson et al., 2005; Picazio et al., 2014) and termination (Heinrichs-Graham et al., 2017; Kilavik et al., 2013; Pfurtscheller et al., 1996). In contrast, its desynchronization is linked to movement preparation and execution (Kühn et al., 2004; Pfurtscheller et al., 1997; Pfurtscheller & Lopes da Silva, 1999; Zhang et al., 2008). Even so, the entrainment of beta-band activity through transcranial alternating-current stimulation could slow down voluntary movements in healthy subjects (Pogosyan et al., 2009). In this respect, Engel and Fries (2010) suggested that beta oscillations signal the status quo, that is, the maintenance of the current sensorimotor set. They claimed that the intention or prediction to hold on to the present state

leads to robust beta band coupling. Accordingly, we reason that the robust beta oscillatory coupling between FEF and SPC stems in part from the suppression of eye movements during fixation on the crosshair. It might reflect the inhibition of the oculomotion, sustained attention, and the status quo during the resting state. Thereby, movement or visuomovement cells in FEF (Gregoriou et al., 2012) likely drive FEF's intrinsic beta synchronization with SPC. In this respect, it is important to note that the deficits in motor response inhibition are linked to attention deficit hyperactivity disorder (Mostofsky et al., 2001). Compared with control subjects, people with attention deficit hyperactivity disorder perform particularly poorly in anti-saccade tasks that require suppressing reflexive eye movements. They also have difficulties maintaining steady fixations for a prolonged time (Munoz et al., 2003; Munoz & Everling, 2004). Consequently, it has been argued that the disruption in FEF and SPC's functional coupling in the beta band might indicate a specific deficit in endogenous visuospatial attention (Caldani et al., 2020).

Engel and Fries (2010) proposed that while beta-band activity signals the maintenance of the status quo, gamma-band oscillations indicate the readiness to change it. In line with this statement, several studies reported task-induced gamma synchrony in object perception, attention and working memory (Baldauf & Desimone, 2014; Fries, 2005, 2009, 2015; Gregoriou et al., 2009; Howard et al., 2003; Tallon-Baudry et al., 1998; Tallon-Baudry & Bertrand, 1999). Despite its specific role in the task-based activity, all three connectivity measures of our present analysis revealed spontaneous functional connectivity between IFJ and the areas in the ventral visual stream in the gamma band. We suggest that, in particular, IFJ and IT cortex's intrinsic gamma coupling hints at their task-evoked functional connectivity during object-based attention (Baldauf & Desimone, 2014). However, it is also likely that this spontaneous synchronization between IFJ and the ventral stream is driven by mind-wandering or imagery in the scanner (Albers et al., 2013; de Borst et al., 2012; Dentico et al., 2014; Dijkstra et al., 2017, 2018; Pearson, 2019). Finally, the predominant functional coupling between IFJ and the ventral visual stream in the rest might to some extent also be driven by the cross-frequency coupling of delta and gamma rhythms (Bruns & Eckhorn, 2004; Canolty & Knight, 2010).

4.3 | IFJa versus IFJp

The functional connectivity analyses (FEF-IFJa and FEF-IFJp) did not reveal identical predominant intrinsic connectivities for IFJa and IFJp (Figures 2, 3 and S2). While

IFJp had weaker functional coupling, IFJa showed stronger predominant connectivity with the ventral visual stream compared with FEF. Additionally, the follow-up analysis (IFJp–IFJa) indicated that IFJa has a more robust functional relation with the ventral visual areas than IFJp (Figure 4). Because IFJa and IFJp have different intrinsic connectivity fingerprints, we suggest that they are functionally distinct regions (Passingham et al., 2002), as already emphasized in MMP1 (Glasser et al., 2016). Several studies also have provided evidence in this direction. For instance, Zanto et al. (2011) pointed out that the right IFJ's dorsal and ventral parts are respectively involved in motion and colour processing (see also Schwedhelm et al., 2020). Moreover, Asplund et al. (2010) proposed that IFJ might have a role in both stimulus-driven and goal-directed attention through its participation in the ventral and dorsal attention networks. Likewise, the deterministic tractography revealed distinct white matter tracts for IFJa and IFJp. IFJa has stronger structural connectivity with the middle and IT gyrus (areas TE1a, TE1m and TE2a) than IFJp (Baker et al., 2018). Following the previous evidence, the distinct predominant connectivities of IFJa and IFJp in the current study attribute different functional roles to these two neighboring regions. At the same time, they also provide strong evidence against the notion that our connectivity results could potentially suffer from a proximity bias (because IFJa is more distant from the visual cortex than IFJp).

4.4 | Area MT

FEF, IFJa or IFJp did not have statistically significant coupling with several regions, including area MT when their connectivities contrasted against each other (Figures 2 and S2) or the average connectivity value per frequency band (Figure S4). Similarly, Hipp et al. (2012) did not find a power correlation between MT and the areas in the frontal cortex when they tested MT's seed-based connectivity in the resting state against the average correlation for a 16-Hz carrier frequency. In particular, the absence of functional coupling between FEF and MT is surprising given the fact that there are direct anatomical pathways (Stanton et al., 1995) and functional links between them (Heinen et al., 2014; Silvanto et al., 2006). Here, we attribute MT's statistically nonsignificant coupling with FEF to the following reasons. First, area MT is involved not only in processing spatial information but also involved in feature information such as the direction of motion (Treue & Trujillo, 1999). In this respect, it is reasonable to observe that MT has similar strength in its functional connectivity either with FEF, IFJa or IFJp. Additionally, area MT is highly connected with early

visual areas compared with the frontal cortex (Figure S1). For this reason, when its functional connectivity is tested against the whole-brain mean connectivity, the frontal areas might not have statistically higher connectivity to survive the statistical test for a respective frequency band. Therefore, we do not propose that there is no functional link between MT and FEF. Instead, our results suggest that their functional coupling is not statistically strong enough to be revealed in direct contrast to MT's strong local connections.

4.5 | Directionality of interactions

In addition to bidirectional functional coupling, we investigated unidirectional interactions between pIPFC and visual streams in the resting state. The majority of regions in Table 1 exhibited a clear directionality to the respectively connected seed region in the lower frequencies. Distinctively, the partial directed coherence revealed directed interactions from the medial part of SPC (areas 7AL, 7PL and VIP) to FEF and IFJ in delta and theta oscillations (Figure 5). Because in the delta and theta range the neural time series of medial SPC are statistically more successful in predicting the neural activity in FEF and IFJ than vice versa, we can suggest that SPC drive the spontaneous activity in pIPFC during the resting state. This unidirectional connectivity during rest might reflect similar task-related influences of SPC over pIPFC during attention and working memory (Chen et al., 2020; Spadone et al., 2021). Moreover, we observe a frequency-specific direction of interaction in the beta band from parietal areas VIP and 7PL to FEF. A recent study by Spadone et al. (2021) similarly indicated that contrasting stay with shift cues reveals a robust direction of information flow from the right ventral IPS to the right dorsal FEF, specifically in the beta band. Because subjects had to fixate during rMEG, the maintenance of attention and the status quo of beta oscillations could have led to the results we observed in the current study. The feedback influences of IPS onto FEF in the beta band might signal to sustain eye movements on the crosshair. Finding such directionality from the parietal cortex to FEF in the beta band is somewhat surprising and speaks against the general notion that the beta band solely provides top-down signals. For these reasons, it is important to note that the direction of an oscillatory coupling is often task dependent (Bastos et al., 2015). Therefore, the direction of interaction between SPC and FEF might change depending on the current task demands. While SPC's highly significant directional influence over pIPFC during the resting state suggests that it might provide some sort of spatial attention signals to FEF, Spadone

et al. (2021) also showed that the contrast between shift versus stay cues reveals feedback connections from the right FEF to the right SPL.

Additionally, we found a highly significant direction of interaction from IFJp to FEF and IFJa in both hemispheres (Figure 6). Within the right hemisphere, alpha and beta oscillations govern these intrinsic directional influences from IFJp to the other seed regions. In the left hemisphere, the same direction of information flow is also supported by theta, in addition to alpha and beta bands. Although the direction of interaction may change depending on task-related activity, it is clear that IFJp feeds information to FEF and IFJa in the resting state. Taken together with the previously mentioned result that IFJp has overall less functional connectivity with the ventral visual cortex (Figure 4), this may hint at its role in providing precursor activity to the output units in FEF and IFJa, which then channel that information to the visual cortex. Interestingly, also in non-human primate recordings, similar interactions have been observed between VPA (the non-human homolog of IFJ) and FEF in visual search tasks: Signals from VPA to FEF helped guide eye movements to likely targets, and the inactivation of VPA selectively eliminated feature attention signals in FEF (while leaving spatial attention and eye movement control unaffected), such that the animals could no longer find targets during visual search (Bichot et al., 2019). In addition to this, we observe a reverse interaction from FEF to IFJa in delta and gamma bands (Figure 6). Strikingly, the predominant coupling between IFJa and the ventral visual areas also occurs in the same frequency bands (Figure 3). In a visual search task, FEF might potentially exchange information about the behavioural relevance of a target's feature with IFJa and modulate IFJa's oscillatory coupling with the ventral visual areas. However, further research is needed to reveal whether FEF still influences IFJ in task-based settings because unidirectional interactions can also be observed because of the asymmetries in the noise levels (Nolte et al., 2004).

4.6 | Power- and phase-based connectivity

Power-based metrics compute functional connectivity between two neuronal oscillations based on their amplitudes' envelope correlation whereas phased-based measures instead consider the consistent relative phase difference between them (Siegel et al., 2012). In this sense, phase-based measures such as iCOH or dwPLI are more delicate and more conservative while attributing a genuine functional coupling in comparison to power-based metrics to avoid Type I errors (Nolte et al., 2004; Stam et al., 2007; Vinck et al., 2011). However, they both have

also been found to be less replicable on the group level and less consistent within and between subjects in resting state data (Colclough et al., 2016). On the other hand, the orthogonalized power envelope correlations (Brookes et al., 2011; Hipp et al., 2012) are shown to be the most replicable methods among several alternative metrics, and, importantly, they are insensitive to spurious coupling stemming from field-spread (Colclough et al., 2016; Duan et al., 2021). In light of these findings, we state that the robustness of power- and phase-based measures in the current study relates to the nature of these metrics' computation. Compared with power correlation, the lower number of significant phase couplings for iCOH and dwPLI might be due to their detailed measurement. Additionally, power- and phase-coupling might potentially reflect partly distinct neuronal mechanisms (Daffertshofer et al., 2018; Siems & Siegel, 2020). For instance, a recent study by Cattai et al. (2021) suggests that power correlation between cortical regions codes for a basic substrate of neural communication. In contrast, their phase synchronization favours information binding through coherence (Fries, 2015). In this context, the sparse number of significant regions for iCOH and dwPLI methods compared with the oPEC metric might also be due to the task-free design of these rMEG recordings. Because there is no explicit competition between areas to communicate one piece of information over the other, the spontaneous neural activity in the resting state might rather reveal the baseline power synchronization to be paved for the precise phase coupling during task settings. For this reason, we might observe that the intrinsic functional connectivity for power-based metrics has higher group-level repeatability and greater within- and between-subject consistency compared with phase-based measures in the resting state (Colclough et al., 2016).

4.7 | Functional lateralization

Both task-based and task-free neuroimaging studies have reported distinct functional lateralization for the dorsal and ventral attentional networks. While the dorsal attentional network including FEF has a bilateral organization, the ventral attentional network including the ventral frontal cortex is lateralized in the right hemisphere (Baldauf & Desimone, 2014; Fox et al., 2006; Vossel et al., 2014). In our study, we observed that FEF had a robust bilateral predominant power correlation with the dorsal visual stream whereas IFJa's predominant power relation with the ventral visual stream was stronger in the right hemisphere. In this respect, these results are well in accordance with the previous findings in the literature. On the other hand, the predominant phase relation of FEF and IFJa with the visual streams was lateralized in the left

hemisphere (Figure S3). These results seem not to be along the same line as the previous findings that suggest a right-lateralized ventral attentional network. However, when we look into the main nodes of the ventral attentional networks, we see that our target areas in the ventral visual stream are not part of the ventral attentional network presented in previous findings (Fox et al., 2006; Vossel et al., 2014). The ventral attentional network rather consists of more dorsally located regions in the temporo-parietal junction and superior temporal sulcus in addition to the ventral frontal cortex. Therefore, the lateralized functional connectivity between IFJa and the ventral visual pathway, which was in either the right (power correlation) or left (phase coupling) hemisphere, does not strictly contradict previous findings but rather provides new additional evidence for their coupling-specific functional lateralizations in the resting state.

4.8 | Future research directions

The resting-state neuroimaging data are time and cost efficient because they ease the acquisition of large datasets and enable analyses of multiple hypotheses over task-free recordings. Nevertheless, it is vital to support the findings from resting-state recordings with task-based settings. To this end, the current study attempted to provide the first direct contrast between the functional connectivity profiles of FEF and IFJ to prompt further research on these regions' functional integration with the rest of the brain. Future studies should compare the functional connectivity fingerprints of these regions in separate experimental paradigms that can isolate the to-be-attend or to-be-remembered spatial versus nonspatial (feature or object) representational content. In many natural tasks, however, both structures work together and readily exchange information. In visual search, for example, nonspatial information about the search template is conveyed from homolog structures in ventrolateral PFC to FEF, which then organizes sequential eye movements to inspect potential targets (Bichot et al., 2015). Similarly, in tasks where feature or object information is bound to specific spatial locations, the communication between IFJ and FEF allows to bind nonspatial information about features or object type and pins it to a specific location. In consequence, such information about features and object type might then also be decoded from FEF (see, e.g., Meyyappan et al., 2021; Rajan et al., 2021). Thereby, it is critical to use special experimental designs in which feature or object information is not in conjunction with spatial representations but separated (e.g., Baldauf & Desimone, 2014), in order to isolate the specific contributions of FEF and IFJ to the encoding of spatial and nonspatial information, respectively.

At the same time, it is an important goal for future studies to relate their neural measurements with the behavioural scores to affirm the proposed functional role of FEF and IFJ in the current study. For example, a task-based neuroimaging study can investigate the correlation between specific time windows and behavioural performance. We would hypothesize that FEF's, rather than IFJ's, functional coupling with the areas in the dorsal stream correlates with the behavioural performance in spatial attention and working memory tasks whereas IFJ's functional connectivity with the regions in the ventral pathway would predict the behavioural scores in feature-based or object-based attention and working memory paradigms. Additionally, it is also crucial to contrast FEF and IFJ's underlying structural connectivity patterns, ideally by using probabilistic rather than deterministic tractography because the latter underestimates the existing long-range white matter tracts (Bedini et al., 2021). Overall, the direct contrast between the intrinsic, functional and anatomical 'connectional fingerprints' (Passingham et al., 2002) of FEF and IFJ would provide further evidence for their functional integration with other cortical regions and a dorsal versus ventral functional segregation within pIPFC as an extension of 'where' and 'what' visual pathways.

5 | CONCLUSIONS

The current study extensively analysed the frequency-specific functional connectivity profiles of FEF and IFJ in the resting state. We presented these regions' intrinsic reciprocal functional connectivities, their dominant directional interactions, and predominant couplings compared with each other for the whole brain and for specific ROIs in the visual streams. As we hypothesized, our ROI analyses revealed that the areas in the dorsal visual stream have a more robust functional coupling with FEF, particularly in beta oscillations. In contrast, the regions in the ventral visual stream have statistically greater functional connectivity with IFJa, particularly in delta and gamma bands. In light of these and previous findings (Cohen et al., 2008; Cole et al., 2014; Fox et al., 2006; Glasser et al., 2016; Mars et al., 2011; Power et al., 2014; Smith et al., 2009; Tavor et al., 2016; Yeo et al., 2011), we argue that intrinsic connectivity patterns are congruent with each brain region's function. Therefore, we conclude that FEF's functional connectivity with the spatiotopically organized regions in SPL and IPS suggests its role in spatial attention and working memory. In contrast, IFJa's functional coupling with feature- or object-encoding areas in the temporal cortex indicates its role in nonspatial attention and working memory. Importantly, these dissociative predominant connectivity profiles of FEF and IFJa

provide strong evidence that the PFC is also organized in a dorsal and ventral stream (Goldman-Rakic, 1996).

AUTHOR CONTRIBUTIONS

O. S. preprocessed and analysed MEG recordings, wrote the initial draft for the manuscript and prepared the table and figures. D. B. conceived the study; supervised the data analyses, writing process and preparation of the figures; and revised the final manuscript.

ACKNOWLEDGEMENTS

This research is supported by *Ministero degli Affari Esteri e della Cooperazione Internazionale* (to O. S.) and *Fondazione Cassa Di Risparmio Di Trento E Rovereto* (to D. B.). Open Access Funding provided by Università degli Studi di Trento within the CRUI-CARE Agreement.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data used in the current study are provided and publicly made available by the 1200 Subjects Release of Human Connectome Project (Larson-Prior et al., 2013; Van Essen et al., 2013) in ConnectomeDB (db.humanconnectome.org; Hodge et al., 2016).

ORCID

Orhan Soyuhos  <https://orcid.org/0000-0002-2858-3979>

Daniel Baldauf  <https://orcid.org/0000-0001-5764-506X>

PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/ejn.15936>.

REFERENCES

- Albers, A. M., Kok, P., Toni, I., Dijkerman, H. C., & de Lange, F. P. (2013). Shared representations for working memory and mental imagery in early visual cortex. *Current Biology*, 23(15), 1427–1431. <https://doi.org/10.1016/j.cub.2013.05.065>
- Andersen, R. A., Bracewell, R. M., Barash, S., Gnadt, J. W., & Fogassi, L. (1990). Eye position effects on visual, memory, and saccade-related activity in areas LIP and 7a of macaque. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 10(4), 1176–1196. <https://doi.org/10.1523/JNEUROSCI.10-04-01176.1990>
- Armstrong, K. M., Chang, M. H., & Moore, T. (2009). Selection and maintenance of spatial information by frontal eye field neurons. *Journal of Neuroscience*, 29(50), 15621–15629. <https://doi.org/10.1523/JNEUROSCI.4465-09.2009>
- Asplund, C. L., Todd, J. J., Snyder, A. P., & Marois, R. (2010). A central role for the lateral prefrontal cortex in goal-directed and stimulus-driven attention. *Nature Neuroscience*, 13(4), 507–512. <https://doi.org/10.1038/nn.2509>
- Babiloni, C., Vecchio, F., Bultrini, A., Luca Romani, G., & Rossini, P. M. (2006). Pre- and Poststimulus alpha rhythms are related to conscious visual perception: A high-resolution EEG study. *Cerebral Cortex*, 16(12), 1690–1700. <https://doi.org/10.1093/cercor/bhj104>
- Baccalá, L. A., & Sameshima, K. (2001). Partial directed coherence: A new concept in neural structure determination. *Biological Cybernetics*, 84(6), 463–474. <https://doi.org/10.1007/PL00007990>
- Bae, G.-Y., & Luck, S. J. (2019). Decoding motion direction using the topography of sustained ERPs and alpha oscillations. *NeuroImage*, 184, 242–255. <https://doi.org/10.1016/j.neuroimage.2018.09.029>
- Bagherzadeh, Y., Baldauf, D., Pantazis, D., & Desimone, R. (2020). Alpha synchrony and the neurofeedback control of spatial attention. *Neuron*, 105(3), 577–587.e5. <https://doi.org/10.1016/j.neuron.2019.11.001>
- Baillet, S., Mosher, J. C., & Leahy, R. M. (2001). Electromagnetic brain mapping. *IEEE Signal Processing Magazine*, 18(6), 14–30. <https://doi.org/10.1109/79.962275>
- Baker, C. M., Burks, J. D., Briggs, R. G., Conner, A. K., Glenn, C. A., Sali, G., McCoy, T. M., Battiste, J. D., O'Donoghue, D. L., & Sughrue, M. E. (2018). A connectomic atlas of the human cerebrum—Chapter 1: Introduction, methods, and significance. *Operative neurosurgery (Hagerstown, Md.)*, 15, S1–S9. <https://doi.org/10.1093/ons/opy253>
- Baldauf, D., Cui, H., & Andersen, R. A. (2008). The posterior parietal cortex encodes in parallel both goals for double-reach sequences. *Journal of Neuroscience*, 28(40), 10081–10089. <https://doi.org/10.1523/JNEUROSCI.3423-08.2008>
- Baldauf, D., & Desimone, R. (2014). Neural mechanisms of object-based attention. *Science (New York, N.Y.)*, 344(6182), 424–427. <https://doi.org/10.1126/science.1247003>
- Baldauf, D., & Deubel, H. (2010). Attentional landscapes in reaching and grasping. *Vision Research*, 50(11), 999–1013. <https://doi.org/10.1016/j.visres.2010.02.008>
- Barash, S., Bracewell, R. M., Fogassi, L., Gnadt, J. W., & Andersen, R. A. (1991). Saccade-related activity in the lateral intraparietal area. I. Temporal properties; comparison with area 7a. *Journal of Neurophysiology*, 66(3), 1095–1108. <https://doi.org/10.1152/jn.1991.66.3.1095>
- Barry, R. J., De Blasio, F. M., Fogarty, J. S., & Clarke, A. R. (2020). Natural alpha frequency components in resting EEG and their relation to arousal. *Clinical Neurophysiology*, 131(1), 205–212. <https://doi.org/10.1016/j.clinph.2019.10.018>
- Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and theta oscillations govern cognitive processes. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, 39(2–3), 241–248. [https://doi.org/10.1016/s0167-8760\(00\)00145-8](https://doi.org/10.1016/s0167-8760(00)00145-8)
- Bastos, A. M., & Schoffelen, J.-M. (2016). A tutorial review of functional connectivity analysis methods and their interpretational pitfalls. *Frontiers in Systems Neuroscience*, 9, 175. <https://doi.org/10.3389/fnsys.2015.00175>
- Bastos, A. M., Vezoli, J., Bosman, C. A., Schoffelen, J.-M., Oostenveld, R., Dowdall, J. R., De Weerd, P., Kennedy, H., & Fries, P. (2015). Visual areas exert feedforward and feedback influences through distinct frequency channels. *Neuron*, 85(2), 390–401. <https://doi.org/10.1016/j.neuron.2014.12.018>
- Bechtold, B. (2016). Violin plots for Matlab, Github Project. <https://doi.org/10.5281/zenodo.4559847>

- Bedini, M., & Baldauf, D. (2021). Structure, function and connectivity fingerprints of the frontal eye field versus the inferior frontal junction: A comprehensive comparison. *European Journal of Neuroscience*, *54*(4), 5462–5506. <https://doi.org/10.1111/ejn.15393>
- Bedini, M., Olivetti, E., Avesani, P., & Baldauf, D. (2021). The anatomical pathways underlying spatial versus non-spatial attention. *Cognitive Processing*, *22*(1), 35–36.
- Behrmann, M., Geng, J. J., & Shomstein, S. (2004). Parietal cortex and attention. *Current Opinion in Neurobiology*, *14*(2), 212–217. <https://doi.org/10.1016/j.conb.2004.03.012>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B: Methodological*, *57*(1), 289–300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Bichot, N. P., Heard, M. T., DeGennaro, E. M., & Desimone, R. (2015). A source for feature-based attention in the prefrontal cortex. *Neuron*, *88*(4), 832–844.
- Bichot, N. P., Rossi, A. F., & Desimone, R. (2005). Parallel and serial neural mechanisms for visual search in macaque area V4. *Science*, *308*, 529–534. <https://doi.org/10.1126/science.1109676>
- Bichot, N. P., Xu, R., Ghadooshahy, A., Williams, M. L., & Desimone, R. (2019). The role of prefrontal cortex in the control of feature attention in area V4. *Nature Communications*, *10*(1), 5727. <https://doi.org/10.1038/s41467-019-13761-7>
- Bisley, J. W., & Goldberg, M. E. (2003). Neuronal activity in the lateral intraparietal area and spatial attention. *Science*, *299*(5603), 81–86. <https://doi.org/10.1126/science.1077395>
- Blatt, G. J., Andersen, R. A., & Stoner, G. R. (1990). Visual receptive field organization and cortico-cortical connections of the lateral intraparietal area (area LIP) in the macaque. *The Journal of Comparative Neurology*, *299*(4), 421–445. <https://doi.org/10.1002/cne.902990404>
- Brancaccio, A., Tabarelli, D., Bigica, M., & Baldauf, D. (2020). Cortical source localization of sleep-stage specific oscillatory activity. *Scientific Reports*, *10*(1), 6976. <https://doi.org/10.1038/s41598-020-63933-5>
- Brass, M., Derrfuss, J., Forstmann, B., & von Cramon, D. Y. (2005). The role of the inferior frontal junction area in cognitive control. *Trends in Cognitive Sciences*, *9*(7), 314–316. <https://doi.org/10.1016/j.tics.2005.05.001>
- Bremmer, F. (2005). Navigation in space—The role of the macaque ventral intraparietal area. *The Journal of Physiology*, *566*(1), 29–35. <https://doi.org/10.1113/jphysiol.2005.082552>
- Bressler, S. L., Tang, W., Sylvester, C. M., Shulman, G. L., & Corbetta, M. (2008). Top-down control of human visual cortex by frontal and parietal cortex in anticipatory visual spatial attention. *Journal of Neuroscience*, *28*(40), 10056–10061. <https://doi.org/10.1523/JNEUROSCI.1776-08.2008>
- Brookes, M. J., Hale, J. R., Zumer, J. M., Stevenson, C. M., Francis, S. T., Barnes, G. R., Owen, J. P., Morris, P. G., & Nagarajan, S. S. (2011). Measuring functional connectivity using MEG: Methodology and comparison with fMRI. *NeuroImage*, *56*(3), 1082–1104. <https://doi.org/10.1016/j.neuroimage.2011.02.054>
- Bruns, A., & Eckhorn, R. (2004). Task-related coupling from high- to low-frequency signals among visual cortical areas in human subdural recordings. *International Journal of Psychophysiology*, *51*(2), 97–116. <https://doi.org/10.1016/j.ijpsycho.2003.07.001>
- Buzsáki, G. (2002). Theta oscillations in the hippocampus. *Neuron*, *33*(3), 325–340. [https://doi.org/10.1016/S0896-6273\(02\)00586-X](https://doi.org/10.1016/S0896-6273(02)00586-X)
- Caldani, S., Isel, F., Septier, M., Acquaviva, E., Delorme, R., & Bucci, M. P. (2020). Impairment in attention focus during the Posner cognitive task in children with ADHD: An eye tracker study. *Frontiers in Pediatrics*, *8*, 484. <https://doi.org/10.3389/fped.2020.00484>
- Canolty, R. T., & Knight, R. T. (2010). The functional role of cross-frequency coupling. *Trends in Cognitive Sciences*, *14*(11), 506–515. <https://doi.org/10.1016/j.tics.2010.09.001>
- Cantero, J. L., Atienza, M., Salas, R. M., & Gómez, C. M. (1999). Alpha EEG coherence in different brain states: An electrophysiological index of the arousal level in human subjects. *Neuroscience Letters*, *271*(3), 167–170. [https://doi.org/10.1016/S0304-3940\(99\)00565-0](https://doi.org/10.1016/S0304-3940(99)00565-0)
- Capotosto, P., Babiloni, C., Romani, G. L., & Corbetta, M. (2009). Frontoparietal cortex controls spatial attention through modulation of anticipatory alpha rhythms. *Journal of Neuroscience*, *29*(18), 5863–5872. <https://doi.org/10.1523/JNEUROSCI.0539-09.2009>
- Carrasco, M. (2011). Visual attention: The past 25 years. *Vision Research*, *51*(13), 1484–1525. <https://doi.org/10.1016/j.visres.2011.04.012>
- Caspari, N., Arsenault, J. T., Vandenberghe, R., & Vanduffel, W. (2018). Functional similarity of medial superior parietal areas for shift-selective attention signals in humans and monkeys. *Cerebral Cortex*, *28*(6), 2085–2099. <https://doi.org/10.1093/cercor/bhx114>
- Cattai, T., Colonnese, S., Corsi, M. C., Bassett, D. S., Scarano, G., & De Vico Fallani, F. (2021). Phase/amplitude synchronization of brain signals during motor imagery BCI tasks. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, *29*, 1168–1177. <https://doi.org/10.1109/TNSRE.2021.3088637>
- Cavada, C., & Goldman-Rakic, P. S. (1989). Posterior parietal cortex in rhesus monkey: II. Evidence for segregated corticocortical networks linking sensory and limbic areas with the frontal lobe. *Journal of Comparative Neurology*, *287*(4), 422–445. <https://doi.org/10.1002/cne.902870403>
- Chen, A., DeAngelis, G. C., & Angelaki, D. E. (2011). Representation of vestibular and visual cues to self-motion in ventral intraparietal cortex. *Journal of Neuroscience*, *31*(33), 12036–12052. <https://doi.org/10.1523/JNEUROSCI.0395-11.2011>
- Chen, X., Zirnsak, M., Vega, G. M., Govil, E., Lomber, S. G., & Moore, T. (2020). Parietal cortex regulates visual salience and salience-driven behavior. *Neuron*, *106*(1), 177–187.e4. <https://doi.org/10.1016/j.neuron.2020.01.016>
- Chou, Y., Sundman, M., Whitson, H. E., Gaur, P., Chu, M.-L., Weingarten, C. P., Madden, D. J., Wang, L., Kirste, I., Joliot, M., Diaz, M. T., Li, Y.-J., Song, A. W., & Chen, N. (2017). Maintenance and representation of mind wandering during resting-state fMRI. *Scientific Reports*, *7*(1), 40722. <https://doi.org/10.1038/srep40722>
- Cohen, A. L., Fair, D. A., Dosenbach, N. U. F., Miezin, F. M., Dierker, D., Van Essen, D. C., Schlaggar, B. L., & Petersen, S. E. (2008). Defining functional areas in individual human brains using resting functional connectivity MRI. *NeuroImage*, *41*(1), 45–57. <https://doi.org/10.1016/j.neuroimage.2008.01.066>
- Colby, C. L., Duhamel, J. R., & Goldberg, M. E. (1993). Ventral intraparietal area of the macaque: Anatomic location and

- visual response properties. *Journal of Neurophysiology*, 69(3), 902–914. <https://doi.org/10.1152/jn.1993.69.3.902>
- Colclough, G. L., Woolrich, M. W., Tewarie, P. K., Brookes, M. J., Quinn, A. J., & Smith, S. M. (2016). How reliable are MEG resting-state connectivity metrics? *NeuroImage*, 138, 284–293. <https://doi.org/10.1016/j.neuroimage.2016.05.070>
- Cole, M. W., Bassett, D. S., Power, J. D., Braver, T. S., & Petersen, S. E. (2014). Intrinsic and task-evoked network architectures of the human brain. *Neuron*, 83(1), 238–251. <https://doi.org/10.1016/j.neuron.2014.05.014>
- Cole, M. W., Ito, T., Bassett, D. S., & Schultz, D. H. (2016). Activity flow over resting-state networks shapes cognitive task activations. *Nature Neuroscience*, 19(12), 1718–1726. <https://doi.org/10.1038/nn.4406>
- Connolly, J. D., Andersen, R. A., & Goodale, M. A. (2003). fMRI evidence for a “parietal reach region” in the human brain. *Experimental Brain Research*, 153(2), 140–145. <https://doi.org/10.1007/s00221-003-1587-1>
- Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, 3(3), 201–215. <https://doi.org/10.1038/nrn755>
- Corbetta, M., Shulman, G. L., Miezin, F. M., & Petersen, S. E. (1995). Superior parietal cortex activation during spatial attention shifts and visual feature conjunction. *Science*, 270(5237), 802–805. <https://doi.org/10.1126/science.270.5237.802>
- Culham, J. C., Cavina-Pratesi, C., & Singhal, A. (2006). The role of parietal cortex in visuomotor control: What have we learned from neuroimaging? *Neuropsychologia*, 44(13), 2668–2684. <https://doi.org/10.1016/j.neuropsychologia.2005.11.003>
- Daffertshofer, A., Ton, R., Kringelbach, M. L., Woolrich, M., & Deco, G. (2018). Distinct criticality of phase and amplitude dynamics in the resting brain. *NeuroImage*, 180, 442–447. <https://doi.org/10.1016/j.neuroimage.2018.03.002>
- Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I. Segmentation and surface reconstruction. *NeuroImage*, 9(2), 179–194. <https://doi.org/10.1006/nimg.1998.0395>
- Dale, A. M., Liu, A. K., Fischl, B. R., Buckner, R. L., Belliveau, J. W., Lewine, J. D., & Halgren, E. (2000). Dynamic statistical parametric mapping: Combining fMRI and MEG for high-resolution imaging of cortical activity. *Neuron*, 26(1), 55–67. [https://doi.org/10.1016/s0896-6273\(00\)81138-1](https://doi.org/10.1016/s0896-6273(00)81138-1)
- de Borst, A. W., Sack, A. T., Jansma, B. M., Esposito, F., de Martino, F., Valente, G., Roebroeck, A., di Salle, F., Goebel, R., & Formisano, E. (2012). Integration of “what” and “where” in frontal cortex during visual imagery of scenes. *NeuroImage*, 60(1), 47–58. <https://doi.org/10.1016/j.neuroimage.2011.12.005>
- de Vries, E., & Baldauf, D. (2019). Attentional weighting in the face processing network: A magnetic response image-guided magnetoencephalography study using multiple cyclic entrainments. *Journal of Cognitive Neuroscience*, 31(10), 1573–1588. https://doi.org/10.1162/jocn_a_01428
- de Vries, I. E. J., Marinato, G., & Baldauf, D. (2021). Decoding object-based auditory attention from source-reconstructed MEG alpha oscillations. *Journal of Neuroscience*, 41(41), 8603–8617. <https://doi.org/10.1523/JNEUROSCI.0583-21.2021>
- de Vries, I. E. J., Savran, E., van Driel, J., & Olivers, C. N. L. (2019). Oscillatory mechanisms of preparing for visual distraction. *Journal of Cognitive Neuroscience*, 31(12), 1873–1894. https://doi.org/10.1162/jocn_a_01460
- Dentico, D., Cheung, B. L., Chang, J.-Y., Guokas, J., Boly, M., Tononi, G., & Van Veen, B. (2014). Reversal of cortical information flow during visual imagery as compared to visual perception. *NeuroImage*, 100, 237–243. <https://doi.org/10.1016/j.neuroimage.2014.05.081>
- Derrfuss, J., Brass, M., Neumann, J., & von Cramon, D. Y. (2005). Involvement of the inferior frontal junction in cognitive control: Meta-analyses of switching and Stroop studies. *Human Brain Mapping*, 25(1), 22–34. <https://doi.org/10.1002/hbm.20127>
- Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review of Neuroscience*, 18, 193–222. <https://doi.org/10.1146/annurev.ne.18.030195.001205>
- Dijkstra, N., Bosch, S. E., & van Gerven, M. A. J. (2017). Vividness of visual imagery depends on the neural overlap with perception in visual areas. *Journal of Neuroscience*, 37(5), 1367–1373. <https://doi.org/10.1523/JNEUROSCI.3022-16.2016>
- Dijkstra, N., Mostert, P., de Lange, F. P., Bosch, S., & van Gerven, M. A. (2018). Differential temporal dynamics during visual imagery and perception. *eLife*, 7, e33904. <https://doi.org/10.7554/eLife.33904>
- Donahue, C. J., Glasser, M. F., Preuss, T. M., Rilling, J. K., & Van Essen, D. C. (2018). Quantitative assessment of prefrontal cortex in humans relative to nonhuman primates. *Proceedings of the National Academy of Sciences of the United States of America*, 115(22), E5183–E5192. <https://doi.org/10.1073/pnas.1721653115>
- Duan, W., Chen, X., Wang, Y.-J., Zhao, W., Yuan, H., & Lei, X. (2021). Reproducibility of power spectrum, functional connectivity and network construction in resting-state EEG. *Journal of Neuroscience Methods*, 348, 108985. <https://doi.org/10.1016/j.jneumeth.2020.108985>
- Duhamel, J.-R., Colby, C. L., & Goldberg, M. E. (1998). Ventral intraparietal area of the macaque: Congruent visual and somatic response properties. *Journal of Neurophysiology*, 79(1), 126–136. <https://doi.org/10.1152/jn.1998.79.1.126>
- Duncan, J. (2001). An adaptive coding model of neural function in prefrontal cortex. *Nature Reviews Neuroscience*, 2(11), 820–829. <https://doi.org/10.1038/35097575>
- Engel, A. K., & Fries, P. (2010). Beta-band oscillations—Signalling the status quo? *Current Opinion in Neurobiology*, 20(2), 156–165. <https://doi.org/10.1016/j.conb.2010.02.015>
- Fecteau, J. H., & Munoz, D. P. (2006). Saliency, relevance, and firing: A priority map for target selection. *Trends in Cognitive Sciences*, 10(8), 382–390. <https://doi.org/10.1016/j.tics.2006.06.011>
- Felleman, D. J., & Van Essen, D. C. (1991). Distributed hierarchical processing in the primate cerebral cortex. *Cerebral Cortex (New York, N.Y. 1991)*, 1(1), 1–47. <https://doi.org/10.1093/cercor/1.1.1>
- Ferreira, R. A., Göbel, S. M., Hymers, M., & Ellis, A. W. (2015). The neural correlates of semantic richness: Evidence from an fMRI study of word learning. *Brain and Language*, 143, 69–80. <https://doi.org/10.1016/j.bandl.2015.02.005>
- Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *NeuroImage*, 9(2), 195–207. <https://doi.org/10.1006/nimg.1998.0396>
- Fischl, B., Sereno, M. I., Tootell, R. B., & Dale, A. M. (1999). High-resolution intersubject averaging and a coordinate system for the cortical surface. *Human Brain Mapping*, 8(4), 272–284.

- [https://doi.org/10.1002/\(sici\)1097-0193\(1999\)8:4<272::aid-hbm10>3.0.co;2-4](https://doi.org/10.1002/(sici)1097-0193(1999)8:4<272::aid-hbm10>3.0.co;2-4)
- Fox, M. D., Corbetta, M., Snyder, A. Z., Vincent, J. L., & Raichle, M. E. (2006). Spontaneous neuronal activity distinguishes human dorsal and ventral attention systems. *Proceedings of the National Academy of Sciences*, *103*(26), 10046–10051. <https://doi.org/10.1073/pnas.0604187103>
- Fraschini, M., Demuru, M., Crobe, A., Marrosu, F., Stam, C. J., & Hillebrand, A. (2016). The effect of epoch length on estimated EEG functional connectivity and brain network organisation. *Journal of Neural Engineering*, *13*(3), 036015. <https://doi.org/10.1088/1741-2560/13/3/036015>
- Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480. <https://doi.org/10.1016/j.tics.2005.08.011>
- Fries, P. (2009). Neuronal gamma-band synchronization as a fundamental process in cortical computation. *Annual Review of Neuroscience*, *32*(1), 209–224. <https://doi.org/10.1146/annurev.neuro.051508.135603>
- Fries, P. (2015). Rhythms for cognition: Communication through coherence. *Neuron*, *88*(1), 220–235. <https://doi.org/10.1016/j.neuron.2015.09.034>
- Fuster, J. M. (2001). The prefrontal cortex—An update: Time is of the essence. *Neuron*, *30*(2), 319–333. [https://doi.org/10.1016/S0896-6273\(01\)00285-9](https://doi.org/10.1016/S0896-6273(01)00285-9)
- Geng, J. J. (2014). Attentional mechanisms of distractor suppression. *Current Directions in Psychological Science*, *23*(2), 147–153. <https://doi.org/10.1177/0963721414525780>
- Giesbrecht, B., Woldorff, M. G., Song, A. W., & Mangun, G. R. (2003). Neural mechanisms of top-down control during spatial and feature attention. *NeuroImage*, *19*(3), 496–512. [https://doi.org/10.1016/S1053-8119\(03\)00162-9](https://doi.org/10.1016/S1053-8119(03)00162-9)
- Gilbertson, T., Lalo, E., Doyle, L., Lazzaro, V. D., Cioni, B., & Brown, P. (2005). Existing motor state is favored at the expense of new movement during 13–35 Hz oscillatory synchrony in the human corticospinal system. *Journal of Neuroscience*, *25*(34), 7771–7779. <https://doi.org/10.1523/JNEUROSCI.1762-05.2005>
- Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K., Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), 171–178. <https://doi.org/10.1038/nature18933>
- Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The minimal preprocessing pipelines for the human connectome project. *NeuroImage*, *80*, 105–124. <https://doi.org/10.1016/j.neuroimage.2013.04.127>
- Goldman-Rakic, P. S. (1988). Topography of cognition: Parallel distributed networks in primate association cortex. *Annual Review of Neuroscience*, *11*, 137–156. <https://doi.org/10.1146/annurev.ne.11.030188.001033>
- Goldman-Rakic, P. S. (1996). The prefrontal landscape: Implications of functional architecture for understanding human mentation and the central executive. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *351*(1346), 1445–1453. <https://doi.org/10.1098/rstb.1996.0129>
- Goodale, M. A., & Milner, A. D. (1992). Separate visual pathways for perception and action. *Trends in Neurosciences*, *15*(1), 20–25. [https://doi.org/10.1016/0166-2236\(92\)90344-8](https://doi.org/10.1016/0166-2236(92)90344-8)
- Gorno-Tempini, M. L., & Price, C. J. (2001). Identification of famous faces and buildings: A functional neuroimaging study of semantically unique items. *Brain*, *124*(10), 2087–2097. <https://doi.org/10.1093/brain/124.10.2087>
- Gottlieb, J. (2007). From thought to action: The parietal cortex as a bridge between perception, action, and cognition. *Neuron*, *53*(1), 9–16. <https://doi.org/10.1016/j.neuron.2006.12.009>
- Goutagny, R., Jackson, J., & Williams, S. (2009). Self-generated theta oscillations in the hippocampus. *Nature Neuroscience*, *12*(12), 1491–1493. <https://doi.org/10.1038/nn.2440>
- Gregoriou, G. G., Gotts, S. J., & Desimone, R. (2012). Cell-type-specific synchronization of neural activity in FEF with V4 during attention. *Neuron*, *73*(3), 581–594. <https://doi.org/10.1016/j.neuron.2011.12.019>
- Gregoriou, G. G., Gotts, S. J., Zhou, H., & Desimone, R. (2009). High-frequency, long-range coupling between prefrontal and visual cortex during attention. *Science*, *324*, 1207–1210. <https://doi.org/10.1126/science.1171402>
- Groppe, D. M. (2021). fdr_bh, MATLAB central file exchange. Retrieved from https://www.mathworks.com/matlabcentral/fileexchange/27418-fdr_bh
- Gross, J., Schmitz, F., Schnitzler, I., Kessler, K., Shapiro, K., Hommel, B., & Schnitzler, A. (2004). Modulation of long-range neural synchrony reflects temporal limitations of visual attention in humans. *Proceedings of the National Academy of Sciences*, *101*(35), 13050–13055. <https://doi.org/10.1073/pnas.0404944101>
- Güntekin, B., & Başar, E. (2016). Review of evoked and event-related delta responses in the human brain. *International Journal of Psychophysiology*, *103*, 43–52. <https://doi.org/10.1016/j.ijpsycho.2015.02.001>
- Hall, E. L., Robson, S. E., Morris, P. G., & Brookes, M. J. (2014). The relationship between MEG and fMRI. *NeuroImage*, *102*, 80–91. <https://doi.org/10.1016/j.neuroimage.2013.11.005>
- Harmony, T. (2013). The functional significance of delta oscillations in cognitive processing. *Frontiers in Integrative Neuroscience*, *7*, 83. <https://doi.org/10.3389/fnint.2013.00083>
- Harmony, T., Fernández, T., Silva, J., Bernal, J., Díaz-Comas, L., Reyes, A., Marosi, E., Rodríguez, M., & Rodríguez, M. (1996). EEG delta activity: An indicator of attention to internal processing during performance of mental tasks. *International Journal of Psychophysiology*, *24*(1), 161–171. [https://doi.org/10.1016/S0167-8760\(96\)00053-0](https://doi.org/10.1016/S0167-8760(96)00053-0)
- Heinen, K., Feredoes, E., Ruff, C. C., & Driver, J. (2017). Functional connectivity between prefrontal and parietal cortex drives visuo-spatial attention shifts. *Neuropsychologia*, *99*, 81–91. <https://doi.org/10.1016/j.neuropsychologia.2017.02.024>
- Heinen, K., Feredoes, E., Weiskopf, N., Ruff, C. C., & Driver, J. (2014). Direct evidence for attention-dependent influences of the frontal eye-fields on feature-responsive visual cortex. *Cerebral Cortex*, *24*(11), 2815–2821. <https://doi.org/10.1093/cercor/bht157>
- Heinrichs-Graham, E., Kurz, M. J., Gehringer, J. E., & Wilson, T. W. (2017). The functional role of post-movement beta oscillations in motor termination. *Brain Structure and Function*, *222*(7), 3075–3086. <https://doi.org/10.1007/s00429-017-1387-1>

- Herweg, N. A., Solomon, E. A., & Kahana, M. J. (2020). Theta oscillations in human memory. *Trends in Cognitive Sciences*, 24(3), 208–227. <https://doi.org/10.1016/j.tics.2019.12.006>
- Hincapié, A.-S., Kujala, J., Mattout, J., Pascarella, A., Daligault, S., Delpuech, C., Mery, D., Cosmelli, D., & Jerbi, K. (2017). The impact of MEG source reconstruction method on source-space connectivity estimation: A comparison between minimum-norm solution and beamforming. *NeuroImage*, 156, 29–42. <https://doi.org/10.1016/j.neuroimage.2017.04.038>
- Hipp, J. F., Hawellek, D. J., Corbetta, M., Siegel, M., & Engel, A. K. (2012). Large-scale cortical correlation structure of spontaneous oscillatory activity. *Nature Neuroscience*, 15(6), 884–890. <https://doi.org/10.1038/nn.3101>
- Hodge, M. R., Horton, W., Brown, T., Herrick, R., Olsen, T., Hileman, M. E., McKay, M., Archie, K. A., Cler, E., Harms, M. P., Burgess, G. C., Glasser, M. F., Elam, J. S., Curtiss, S. W., Barch, D. M., Oostenveld, R., Larson-Prior, L. J., Ugurbil, K., Van Essen, D. C., & Marcus, D. S. (2016). ConnectomeDB—Sharing human brain connectivity data. *NeuroImage*, 124, 1102–1107. <https://doi.org/10.1016/j.neuroimage.2015.04.046>
- Howard, M. W., Rizzuto, D. S., Caplan, J. B., Madsen, J. R., Lisman, J., Aschenbrenner-Scheibe, R., Schulze-Bonhage, A., & Kahana, M. J. (2003). Gamma oscillations correlate with working memory load in humans. *Cerebral Cortex*, 13(12), 1369–1374. <https://doi.org/10.1093/cercor/bhg084>
- Huang, M. X., Mosher, J. C., & Leahy, R. M. (1999). A sensor-weighted overlapping-sphere head model and exhaustive head model comparison for MEG. *Physics in Medicine and Biology*, 44(2), 423–440. <https://doi.org/10.1088/0031-9155/44/2/010>
- Ipata, A. E., Gee, A. L., Bisley, J. W., & Goldberg, M. E. (2009). Neurons in the lateral intraparietal area create a priority map by the combination of disparate signals. *Experimental Brain Research*, 192(3), 479–488. <https://doi.org/10.1007/s00221-008-1557-8>
- Jensen, O., & Colgin, L. L. (2007). Cross-frequency coupling between neuronal oscillations. *Trends in Cognitive Sciences*, 11(7), 267–269. <https://doi.org/10.1016/j.tics.2007.05.003>
- Jerde, T. A., Merriam, E. P., Riggall, A. C., Hedges, J. H., & Curtis, C. E. (2012). Prioritized maps of space in human frontoparietal cortex. *Journal of Neuroscience*, 32(48), 17382–17390. <https://doi.org/10.1523/JNEUROSCI.3810-12.2012>
- Kamps, F. S., Morris, E. J., & Dilks, D. D. (2019). A face is more than just the eyes, nose, and mouth: fMRI evidence that face-selective cortex represents external features. *NeuroImage*, 184, 90–100. <https://doi.org/10.1016/j.neuroimage.2018.09.027>
- Kar, K., Kubilius, J., Schmidt, K., Issa, E. B., & DiCarlo, J. J. (2019). Evidence that recurrent circuits are critical to the ventral stream's execution of core object recognition behavior. *Nature Neuroscience*, 22(6), 974–983. <https://doi.org/10.1038/s41593-019-0392-5>
- Kassebaum, P. (2021). circularGraph, GitHub repository. Retrieved from <https://github.com/paul-kassebaum-mathworks/circularGraph>
- Kelley, T. A., Serences, J. T., Giesbrecht, B., & Yantis, S. (2008). Cortical mechanisms for shifting and holding visuospatial attention. *Cerebral Cortex*, 18(1), 114–125. <https://doi.org/10.1093/cercor/bhm036>
- Kilavik, B. E., Zaepffel, M., Brovelli, A., MacKay, W. A., & Riehle, A. (2013). The ups and downs of beta oscillations in sensorimotor cortex. *Experimental Neurology*, 245, 15–26. <https://doi.org/10.1016/j.expneurol.2012.09.014>
- Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition-timing hypothesis. *Brain Research Reviews*, 53(1), 63–88. <https://doi.org/10.1016/j.brainresrev.2006.06.003>
- Kravitz, D. J., Saleem, K. S., Baker, C. I., & Mishkin, M. (2011). A new neural framework for visuospatial processing. *Nature Reviews Neuroscience*, 12(4), 217–230. <https://doi.org/10.1038/nrn3008>
- Kravitz, D. J., Saleem, K. S., Baker, C. I., Ungerleider, L. G., & Mishkin, M. (2013). The ventral visual pathway: An expanded neural framework for the processing of object quality. *Trends in Cognitive Sciences*, 17(1), 26–49. <https://doi.org/10.1016/j.tics.2012.10.011>
- Kreiman, G., Hung, C. P., Kraskov, A., Quiroga, R. Q., Poggio, T., & DiCarlo, J. J. (2006). Object selectivity of local field potentials and spikes in the macaque inferior temporal cortex. *Neuron*, 49(3), 433–445. <https://doi.org/10.1016/j.neuron.2005.12.019>
- Kühn, A. A., Williams, D., Kupsch, A., Limousin, P., Hariz, M., Schneider, G.-H., Yarrow, K., & Brown, P. (2004). Event-related beta desynchronization in human subthalamic nucleus correlates with motor performance. *Brain*, 127(4), 735–746. <https://doi.org/10.1093/brain/awh106>
- Kusunoki, M., & Goldberg, M. E. (2003). The time course of perisaccadic receptive field shifts in the lateral intraparietal area of the monkey. *Journal of Neurophysiology*, 89(3), 1519–1527. <https://doi.org/10.1152/jn.00519.2002>
- Kusunoki, M., Gottlieb, J., & Goldberg, M. E. (2000). The lateral intraparietal area as a salience map: The representation of abrupt onset, stimulus motion, and task relevance. *Vision Research*, 40(10), 1459–1468. [https://doi.org/10.1016/S0042-6989\(99\)00212-6](https://doi.org/10.1016/S0042-6989(99)00212-6)
- Landi, S. M., Viswanathan, P., Serene, S., & Freiwald, W. A. (2021). A fast link between face perception and memory in the temporal pole. *Science*, 373(6554), 581–585. <https://doi.org/10.1126/science.abi6671>
- Larson-Prior, L. J., Oostenveld, R., Della Penna, S., Michalareas, G., Prior, F., Babajani-Feremi, A., Schoffelen, J.-M., Marzetti, L., de Pasquale, F., Di Pompeo, F., Stout, J., Woolrich, M., Luo, Q., Bucholz, R., Fries, P., Pizzella, V., Romani, G. L., Corbetta, M., & Snyder, A. Z. (2013). Adding dynamics to the human connectome project with MEG. *NeuroImage*, 80, 190–201. <https://doi.org/10.1016/j.neuroimage.2013.05.056>
- Lega, B. C., Jacobs, J., & Kahana, M. (2012). Human hippocampal theta oscillations and the formation of episodic memories. *Hippocampus*, 22(4), 748–761. <https://doi.org/10.1002/hipo.20937>
- Lehky, S. R., & Tanaka, K. (2016). Neural representation for object recognition in inferotemporal cortex. *Current Opinion in Neurobiology*, 37, 23–35. <https://doi.org/10.1016/j.conb.2015.12.001>
- Liebe, S., Hoerzer, G. M., Logothetis, N. K., & Rainer, G. (2012). Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nature Neuroscience*, 15(3), 456–462. <https://doi.org/10.1038/nn.3038>
- Liu, T. (2016). Neural representation of object-specific attentional priority. *NeuroImage*, 129, 15–24. <https://doi.org/10.1016/j.neuroimage.2016.01.034>
- Liu, T., Hospadaruk, L., Zhu, D. C., & Gardner, J. L. (2011). Feature-specific attentional priority signals in human cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 31(12), 4484–4495. <https://doi.org/10.1523/JNEUROSCI.5745-10.2011>

- Lubenov, E. V., & Siapas, A. G. (2009). Hippocampal theta oscillations are travelling waves. *Nature*, *459*(7246), 534–539. <https://doi.org/10.1038/nature08010>
- Macko, K. A., Jarvis, C. D., Kennedy, C., Miyaoka, M., Shinohara, M., Sololoff, L., & Mishkin, M. (1982). Mapping the primate visual system with [2-14C]Deoxyglucose. *Science*, *218*(4570), 394–397. <https://doi.org/10.1126/science.7123241>
- Mangun, G. R. (1995). Neural mechanisms of visual selective attention. *Psychophysiology*, *32*(1), 4–18. <https://doi.org/10.1111/j.1469-8986.1995.tb03400.x>
- Mars, R. B., Jbabdi, S., Sallet, J., O'Reilly, J. X., Crosson, P. L., Olivier, E., Noonan, M. P., Bergmann, C., Mitchell, A. S., Baxter, M. G., Behrens, T. E. J., Johansen-Berg, H., Tomassini, V., Miller, K. L., & Rushworth, M. F. S. (2011). Diffusion-weighted imaging tractography-based parcellation of the human parietal cortex and comparison with human and macaque resting-state functional connectivity. *Journal of Neuroscience*, *31*(11), 4087–4100. <https://doi.org/10.1523/JNEUROSCI.5102-10.2011>
- Marshall, T. R., O'Shea, J., Jensen, O., & Bergmann, T. O. (2015). Frontal eye fields control attentional modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(4), 1638–1647. <https://doi.org/10.1523/JNEUROSCI.3116-14.2015>
- Mazaheri, A., DiQuattro, N. E., Bengson, J., & Geng, J. J. (2011). Pre-stimulus activity predicts the winner of top-down vs. bottom-up attentional selection. *PLoS ONE*, *6*(2), e16243. <https://doi.org/10.1371/journal.pone.0016243>
- McCarthy, G., Puce, A., Constable, T., Krystal, J. H., Gore, J. C., & Goldman-Rakic, P. (1996). Activation of human prefrontal cortex during spatial and nonspatial working memory tasks measured by functional MRI. *Cerebral Cortex*, *6*(4), 600–611. <https://doi.org/10.1093/cercor/6.4.600>
- Merrikhi, Y., Clark, K., Albarran, E., Parsa, M., Zirnsak, M., Moore, T., & Noudoost, B. (2017). Spatial working memory alters the efficacy of input to visual cortex. *Nature Communications*, *8*(1), 15041. <https://doi.org/10.1038/ncomms15041>
- Meyyappan, S., Rajan, A., Mangun, G. R., & Ding, M. (2021). Role of inferior frontal junction (IFJ) in the control of feature versus spatial attention. *Journal of Neuroscience*, *41*(38), 8065–8074. <https://doi.org/10.1523/JNEUROSCI.2883-20.2021>
- Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, *24*(1), 167–202. <https://doi.org/10.1146/annurev.neuro.24.1.167>
- Miller, E. K., Erickson, C. A., & Desimone, R. (1996). Neural mechanisms of visual working memory in prefrontal cortex of the macaque. *Journal of Neuroscience*, *16*(16), 5154–5167. <https://doi.org/10.1523/JNEUROSCI.16-16-05154.1996>
- Mishkin, M., Ungerleider, L. G., & Macko, K. A. (1983). Object vision and spatial vision: Two cortical pathways. *Trends in Neurosciences*, *6*, 414–417. [https://doi.org/10.1016/0166-2236\(83\)90190-X](https://doi.org/10.1016/0166-2236(83)90190-X)
- Molenberghs, P., Mesulam, M. M., Peeters, R., & Vandenberghe, R. R. C. (2007). Remapping attentional priorities: Differential contribution of superior parietal lobule and intraparietal sulcus. *Cerebral Cortex*, *17*(11), 2703–2712. <https://doi.org/10.1093/cercor/bhl179>
- Moore, T., & Armstrong, K. M. (2003). Selective gating of visual signals by microstimulation of frontal cortex. *Nature*, *421*(6921), 370–373. <https://doi.org/10.1038/nature01341>
- Moore, T., & Fallah, M. (2004). Microstimulation of the frontal eye field and its effects on covert spatial attention. *Journal of Neurophysiology*, *91*(1), 152–162. <https://doi.org/10.1152/jn.00741.2002>
- Mostofsky, S. H., Lasker, A. G., Cutting, L. E., Denckla, M. B., & Zee, D. S. (2001). Oculomotor abnormalities in attention deficit hyperactivity disorder: A preliminary study. *Neurology*, *57*(3), 423–430. <https://doi.org/10.1212/WNL.57.3.423>
- Muhle-Karbe, P. S., Andres, M., & Brass, M. (2014). Transcranial magnetic stimulation dissociates prefrontal and parietal contributions to task preparation. *Journal of Neuroscience*, *34*(37), 12481–12489. <https://doi.org/10.1523/JNEUROSCI.4931-13.2014>
- Munoz, D. P., Armstrong, I. T., Hampton, K. A., & Moore, K. D. (2003). Altered control of visual fixation and saccadic eye movements in attention-deficit hyperactivity disorder. *Journal of Neurophysiology*, *90*(1), 503–514. <https://doi.org/10.1152/jn.00192.2003>
- Munoz, D. P., & Everling, S. (2004). Look away: The anti-saccade task and the voluntary control of eye movement. *Nature Reviews Neuroscience*, *5*(3), 218–228. <https://doi.org/10.1038/nrn1345>
- Muthukrishnan, S. P., Soni, S., & Sharma, R. (2020). Brain networks communicate through theta oscillations to encode high load in a visuospatial working memory task: An EEG connectivity study. *Brain Topography*, *33*(1), 75–85. <https://doi.org/10.1007/s10548-019-00739-3>
- Noah, S., Powell, T., Khodayari, N., Olivan, D., Ding, M., & Mangun, G. R. (2020). Neural mechanisms of attentional control for objects: Decoding EEG alpha when anticipating faces, scenes, and tools. *Journal of Neuroscience*, *40*(25), 4913–4924. <https://doi.org/10.1523/JNEUROSCI.2685-19.2020>
- Nolte, G., Bai, O., Wheaton, L., Mari, Z., Vorbach, S., & Hallett, M. (2004). Identifying true brain interaction from EEG data using the imaginary part of coherency. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, *115*(10), 2292–2307. <https://doi.org/10.1016/j.clinph.2004.04.029>
- Olson, I. R., Plotzker, A., & Ezzyat, Y. (2007). The enigmatic temporal pole: A review of findings on social and emotional processing. *Brain: A Journal of Neurology*, *130*(Pt 7), 1718–1731. <https://doi.org/10.1093/brain/awm052>
- O'Neill, P.-K., Gordon, J. A., & Sigurdsson, T. (2013). Theta oscillations in the medial prefrontal cortex are modulated by spatial working memory and synchronize with the hippocampus through its ventral subregion. *Journal of Neuroscience*, *33*(35), 14211–14224. <https://doi.org/10.1523/JNEUROSCI.2378-13.2013>
- Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869. <https://doi.org/10.1155/2011/156869>
- Osher, D. E., Saxe, R. R., Koldewyn, K., Gabrieli, J. D. E., Kanwisher, N., & Saygin, Z. M. (2016). Structural connectivity fingerprints predict cortical selectivity for multiple visual categories across cortex. *cerebral cortex (New York, N.Y.: 1991)*, *26*, 1668–1683. <https://doi.org/10.1093/cercor/bhu303>
- Paneri, S., & Gregoriou, G. G. (2017). Top-down control of visual attention by the prefrontal cortex. functional specialization and long-range interactions. *Frontiers in Neuroscience*, *11*, 545. <https://doi.org/10.3389/fnins.2017.00545>

- Passingham, R. E., Stephan, K. E., & Kötter, R. (2002). The anatomical basis of functional localization in the cortex. *Nature Reviews Neuroscience*, 3(8), 606–616. <https://doi.org/10.1038/nrn893>
- Pearson, J. (2019). The human imagination: The cognitive neuroscience of visual mental imagery. *Nature Reviews Neuroscience*, 20(10), 624–634. <https://doi.org/10.1038/s41583-019-0202-9>
- Petit, L., & Pouget, P. (2019). The comparative anatomy of frontal eye fields in primates. *Cortex: a Journal Devoted to the Study of the Nervous System and Behavior*, 118, 51–64. <https://doi.org/10.1016/j.cortex.2019.02.023>
- Pfurtscheller, G., & Lopes da Silva, F. H. (1999). Event-related EEG/MEG synchronization and desynchronization: Basic principles. *Clinical Neurophysiology*, 110(11), 1842–1857. [https://doi.org/10.1016/S1388-2457\(99\)00141-8](https://doi.org/10.1016/S1388-2457(99)00141-8)
- Pfurtscheller, G., Neuper, C., Andrew, C., & Edlinger, G. (1997). Foot and hand area mu rhythms. *International Journal of Psychophysiology*, 26(1), 121–135. [https://doi.org/10.1016/S0167-8760\(97\)00760-5](https://doi.org/10.1016/S0167-8760(97)00760-5)
- Pfurtscheller, G., Stancák, A., & Neuper, C. (1996). Post-movement beta synchronization. A correlate of an idling motor area? *Electroencephalography and Clinical Neurophysiology*, 98(4), 281–293. [https://doi.org/10.1016/0013-4694\(95\)00258-8](https://doi.org/10.1016/0013-4694(95)00258-8)
- Picazio, S., Veniero, D., Ponzio, V., Caltagirone, C., Gross, J., Thut, G., & Koch, G. (2014). Prefrontal control over motor cortex cycles at beta frequency during movement inhibition. *Current Biology*, 24(24), 2940–2945. <https://doi.org/10.1016/j.cub.2014.10.043>
- Pivik, R. T., & Harman, K. (1995). A reconceptualization of EEG alpha activity as an index of arousal during sleep: All alpha activity is not equal. *Journal of Sleep Research*, 4(3), 131–137. <https://doi.org/10.1111/j.1365-2869.1995.tb00161.x>
- Pogosyan, A., Gaynor, L. D., Eusebio, A., & Brown, P. (2009). Boosting cortical activity at beta-band frequencies slows movement in humans. *Current Biology*, 19(19), 1637–1641. <https://doi.org/10.1016/j.cub.2009.07.074>
- Power, J. D., Schlaggar, B. L., & Petersen, S. E. (2014). Studying brain organization via spontaneous fMRI signal. *Neuron*, 84(4), 681–696. <https://doi.org/10.1016/j.neuron.2014.09.007>
- Ptak, R. (2012). The frontoparietal attention network of the human brain: Action, saliency, and a priority map of the environment. *The Neuroscientist*, 18(5), 502–515. <https://doi.org/10.1177/1073858411409051>
- Rajan, A., Meyyappan, S., Liu, Y., Samuel, I., Nandi, B., Mangun, G. R., & Ding, M. (2021). The microstructure of attentional control in the dorsal attention network. *Journal of Cognitive Neuroscience*, 33(6), 965–983. https://doi.org/10.1162/jocn_a_01710
- Rajan, A., Siegel, S. N., Liu, Y., Bengson, J., Mangun, G. R., & Ding, M. (2019). Theta oscillations index frontal decision-making and mediate reciprocal frontal-parietal interactions in willed attention. *Cerebral Cortex*, 29(7), 2832–2843. <https://doi.org/10.1093/cercor/bhy149>
- Rihs, T. A., Michel, C. M., & Thut, G. (2007). Mechanisms of selective inhibition in visual spatial attention are indexed by α -band EEG synchronization. *European Journal of Neuroscience*, 25(2), 603–610. <https://doi.org/10.1111/j.1460-9568.2007.05278.x>
- Romanski, L. M. (2004). Domain specificity in the primate prefrontal cortex. *Cognitive, Affective, & Behavioral Neuroscience*, 4(4), 421–429. <https://doi.org/10.3758/CABN.4.4.421>
- Rossell, S. L., Bullmore, E. T., Williams, S. C. R., & David, A. S. (2001). Brain activation during automatic and controlled processing of semantic relations: A priming experiment using lexical-decision. *Neuropsychologia*, 39(11), 1167–1176. [https://doi.org/10.1016/S0028-3932\(01\)00049-5](https://doi.org/10.1016/S0028-3932(01)00049-5)
- Rossi, A. F., Pessoa, L., Desimone, R., & Ungerleider, L. G. (2008). The prefrontal cortex and the executive control of attention. *Experimental Brain Research*, 192(3), 489–497. <https://doi.org/10.1007/s00221-008-1642-z>
- Sauseng, P., Griesmayr, B., Freunberger, R., & Klimesch, W. (2010). Control mechanisms in working memory: A possible function of EEG theta oscillations. *Neuroscience & Biobehavioral Reviews*, 34(7), 1015–1022. <https://doi.org/10.1016/j.neubiorev.2009.12.006>
- Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., Gruber, W. R., & Birbaumer, N. (2005). A shift of visual spatial attention is selectively associated with human EEG alpha activity. *European Journal of Neuroscience*, 22(11), 2917–2926. <https://doi.org/10.1111/j.1460-9568.2005.04482.x>
- Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D. E., & Saxe, R. R. (2011). Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nature Neuroscience*, 15(2), 321–327. <https://doi.org/10.1038/nn.3001>
- Scalaidhe, Ó., Wilson, F. A., & Goldman-Rakic, P. S. (1999). Face-selective neurons during passive viewing and working memory performance of rhesus monkeys: Evidence for intrinsic specialization of neuronal coding. *Cerebral Cortex*, 9(5), 459–475. <https://doi.org/10.1093/cercor/9.5.459>
- Schlack, A., Hoffmann, K.-P., & Bremmer, F. (2002). Interaction of linear vestibular and visual stimulation in the macaque ventral intraparietal area (VIP). *European Journal of Neuroscience*, 16(10), 1877–1886. <https://doi.org/10.1046/j.1460-9568.2002.02251.x>
- Schlack, A., Hoffmann, K.-P., & Bremmer, F. (2003). Selectivity of macaque ventral intraparietal area (area VIP) for smooth pursuit eye movements. *The Journal of Physiology*, 551(2), 551–561. <https://doi.org/10.1113/jphysiol.2003.042994>
- Schlack, A., Sterbing-D'Angelo, S. J., Hartung, K., Hoffmann, K.-P., & Bremmer, F. (2005). Multisensory space representations in the macaque ventral intraparietal area. *Journal of Neuroscience*, 25(18), 4616–4625. <https://doi.org/10.1523/JNEUROSCI.0455-05.2005>
- Schwedhelm, P., Baldauf, D., & Treue, S. (2020). The lateral prefrontal cortex of primates encodes stimulus colors and their behavioral relevance during a match-to-sample task. *Scientific Reports*, 10(1), 4216. <https://doi.org/10.1038/s41598-020-61171-3>
- Sereno, A. B., & Maunsell, J. H. R. (1998). Shape selectivity in primate lateral intraparietal cortex. *Nature*, 395(6701), 500–503. <https://doi.org/10.1038/26752>
- Siapas, A. G., Lubenov, E. V., & Wilson, M. A. (2005). Prefrontal phase locking to hippocampal theta oscillations. *Neuron*, 46(1), 141–151. <https://doi.org/10.1016/j.neuron.2005.02.028>
- Siegel, M., Donner, T. H., & Engel, A. K. (2012). Spectral fingerprints of large-scale neuronal interactions. *Nature Reviews Neuroscience*, 13(2), 121–134. <https://doi.org/10.1038/nrn3137>
- Siegel, M., Donner, T. H., Oostenveld, R., Fries, P., & Engel, A. K. (2008). Neuronal synchronization along the dorsal visual pathway reflects the focus of spatial attention. *Neuron*, 60(4), 709–719. <https://doi.org/10.1016/j.neuron.2008.09.010>

- Siems, M., & Siegel, M. (2020). Dissociated neuronal phase- and amplitude-coupling patterns in the human brain. *NeuroImage*, 209, 116538. <https://doi.org/10.1016/j.neuroimage.2020.116538>
- Silvanto, J., Lavie, N., & Walsh, V. (2006). Stimulation of the human frontal eye fields modulates sensitivity of extrastriate visual cortex. *Journal of Neurophysiology*, 96(2), 941–945. <https://doi.org/10.1152/jn.00015.2006>
- Siuda-Krzywicka, K., Witzel, C., Bartolomeo, P., & Cohen, L. (2021). Color naming and categorization depend on distinct functional brain networks. *Cerebral Cortex*, 31(2), 1106–1115. <https://doi.org/10.1093/cercor/bhaa278>
- Smith, R., Keramatian, K., & Christoff, K. (2007). Localizing the rostralateral prefrontal cortex at the individual level. *NeuroImage*, 36(4), 1387–1396. <https://doi.org/10.1016/j.neuroimage.2007.04.032>
- Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N., Watkins, K. E., Toro, R., Laird, A. R., & Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during activation and rest. *Proceedings of the National Academy of Sciences*, 106(31), 13040–13045. <https://doi.org/10.1073/pnas.0905267106>
- Spadone, S., Wyczesany, M., Della Penna, S., Corbetta, M., & Capotosto, P. (2021). Directed flow of beta band communication during reorienting of attention within the dorsal attention network. *Brain Connectivity*, 11(9), 717–724. <https://doi.org/10.1089/brain.2020.0885>
- Sprague, T. C., & Serences, J. T. (2013). Attention modulates spatial priority maps in the human occipital, parietal and frontal cortices. *Nature Neuroscience*, 16(12), 1879–1887. <https://doi.org/10.1038/nn.3574>
- Stam, C. J., Nolte, G., & Daffertshofer, A. (2007). Phase lag index: Assessment of functional connectivity from multi channel EEG and MEG with diminished bias from common sources. *Human Brain Mapping*, 28(11), 1178–1193. <https://doi.org/10.1002/hbm.20346>
- Stanton, G. B., Bruce, C. J., & Goldberg, M. E. (1995). Topography of projections to posterior cortical areas from the macaque frontal eye fields. *Journal of Comparative Neurology*, 353(2), 291–305. <https://doi.org/10.1002/cne.903530210>
- Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A user-friendly application for MEG/EEG analysis. *Computational Intelligence and Neuroscience*, 2011, 879716. <https://doi.org/10.1155/2011/879716>
- Tallon-Baudry, C., & Bertrand, O. (1999). Oscillatory gamma activity in humans and its role in object representation. *Trends in Cognitive Sciences*, 3(4), 151–162. [https://doi.org/10.1016/S1364-6613\(99\)01299-1](https://doi.org/10.1016/S1364-6613(99)01299-1)
- Tallon-Baudry, C., Bertrand, O., Peronnet, F., & Pernier, J. (1998). Induced γ -band activity during the delay of a visual short-term memory task in humans. *Journal of Neuroscience*, 18(11), 4244–4254. <https://doi.org/10.1523/JNEUROSCI.18-11-04244.1998>
- Tamber-Rosenau, B. J., Asplund, C. L., & Marois, R. (2018). Functional dissociation of the inferior frontal junction from the dorsal attention network in top-down attentional control. *Journal of Neurophysiology*, 120(5), 2498–2512. <https://doi.org/10.1152/jn.00506.2018>
- Tanaka, K. (1996). Inferotemporal cortex and object vision. *Annual Review of Neuroscience*, 19(1), 109–139. <https://doi.org/10.1146/annurev.ne.19.030196.000545>
- Tavor, I., Jones, O. P., Mars, R. B., Smith, S. M., Behrens, T. E., & Jbabdi, S. (2016). Task-free MRI predicts individual differences in brain activity during task performance. *Science*, 352(6282), 216–220. <https://doi.org/10.1126/science.aad8127>
- Thakral, P. P., & Slotnick, S. D. (2009). The role of parietal cortex during sustained visual spatial attention. *Brain Research*, 1302, 157–166. <https://doi.org/10.1016/j.brainres.2009.09.031>
- Thompson, K. G., & Bichot, N. P. (2005). A visual salience map in the primate frontal eye field. In *Progress in brain research* (Vol. 147) (pp. 249–262). Elsevier. [https://doi.org/10.1016/S0079-6123\(04\)47019-8](https://doi.org/10.1016/S0079-6123(04)47019-8)
- Thompson, K. G., Biscoe, K. L., & Sato, T. R. (2005). Neuronal basis of covert spatial attention in the frontal eye field. *Journal of Neuroscience*, 25(41), 9479–9487. <https://doi.org/10.1523/JNEUROSCI.0741-05.2005>
- Thut, G., Nietzel, A., Brandt, S. A., & Pascual-Leone, A. (2006). α -Band electroencephalographic activity over occipital cortex indexes visuospatial attention Bias and predicts visual target detection. *Journal of Neuroscience*, 26(37), 9494–9502. <https://doi.org/10.1523/JNEUROSCI.0875-06.2006>
- Toscani, M., Marzi, T., Righi, S., Viggiano, M. P., & Baldassi, S. (2010). Alpha waves: A neural signature of visual suppression. *Experimental Brain Research*, 207(3), 213–219. <https://doi.org/10.1007/s00221-010-2444-7>
- Treue, S., & Trujillo, J. C. M. (1999). Feature-based attention influences motion processing gain in macaque visual cortex. *Nature*, 399(6736), 575–579. <https://doi.org/10.1038/21176>
- Uğurbil, K., Xu, J., Auerbach, E. J., Moeller, S., Vu, A. T., Duarte-Carvajalino, J. M., Lenglet, C., Wu, X., Schmitter, S., Van de Moortele, P. F., Strupp, J., Sapiro, G., De Martino, F., Wang, D., Harel, N., Garwood, M., Chen, L., Feinberg, D. A., Smith, S. M., ... Yacoub, E. (2013). Pushing spatial and temporal resolution for functional and diffusion MRI in the human connectome project. *NeuroImage*, 80, 80–104. <https://doi.org/10.1016/j.neuroimage.2013.05.012>
- van Diepen, R. M., Miller, L. M., Mazaheri, A., & Geng, J. J. (2016). The role of alpha activity in spatial and feature-based attention. *ENeuro*, 3(5), ENEURO.0204-16.2016. <https://doi.org/10.1523/ENeuro.0204-16.2016>
- Van Essen, D., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., Ugurbil, K., & WU-Minn HCP Consortium. (2013). The WU-Minn human connectome project: An overview. *NeuroImage*, 80, 62–79. <https://doi.org/10.1016/j.neuroimage.2013.05.041>
- Vandenberghe, R., Gitelman, D. R., Parrish, T. B., & Mesulam, M. M. (2001). Functional specificity of superior parietal mediation of spatial shifting. *NeuroImage*, 14(3), 661–673. <https://doi.org/10.1006/nimg.2001.0860>
- Veniero, D., Gross, J., Morand, S., Duecker, F., Sack, A. T., & Thut, G. (2021). Top-down control of visual cortex by the frontal eye fields through oscillatory realignment. *Nature Communications*, 12(1), 1757. <https://doi.org/10.1038/s41467-021-21979-7>
- Vernet, M., Quentin, R., Chanes, L., Mitsumasu, A., & Valero-Cabré, A. (2014). Frontal eye field, where art thou? Anatomy, function, and non-invasive manipulation of frontal regions involved in eye movements and associated cognitive operations. *Frontiers in Integrative Neuroscience*, 8, 66. <https://doi.org/10.3389/fnint.2014.00066>

- Vinck, M., Oostenveld, R., van Wingerden, M., Battaglia, F., & Pennartz, C. M. A. (2011). An improved index of phase-synchronization for electrophysiological data in the presence of volume-conduction, noise and sample-size bias. *NeuroImage*, 55(4), 1548–1565. <https://doi.org/10.1016/j.neuroimage.2011.01.055>
- Vossel, S., Geng, J. J., & Fink, G. R. (2014). Dorsal and ventral attention systems: Distinct neural circuits but collaborative roles. *The Neuroscientist*, 20(2), 150–159. <https://doi.org/10.1177/1073858413494269>
- Walker, G. M., Schwartz, M. F., Kimberg, D. Y., Faseyitan, O., Brecher, A., Dell, G. S., & Coslett, H. B. (2011). Support for anterior temporal involvement in semantic error production in aphasia: New evidence from VLSM. *Brain and Language*, 117(3), 110–122. <https://doi.org/10.1016/j.bandl.2010.09.008>
- Wang, L., Mruczek, R. E. B., Arcaro, M. J., & Kastner, S. (2015). Probabilistic maps of visual topography in human cortex. *Cerebral Cortex*, 25(10), 3911–3931. <https://doi.org/10.1093/cercor/bhu277>
- Wardak, C., Ibos, G., Duhamel, J.-R., & Olivier, E. (2006). Contribution of the monkey frontal eye field to covert visual attention. *Journal of Neuroscience*, 26(16), 4228–4235. <https://doi.org/10.1523/JNEUROSCI.3336-05.2006>
- Wilson, F. A. W., Scalaidhe, S. P. Ó., & Goldman-Rakic, P. S. (1993). Dissociation of object and spatial processing domains in primate prefrontal cortex. *Science*, 260(5116), 1955–1958. <https://doi.org/10.1126/science.8316836>
- Xie, S., Kaiser, D., & Cichy, R. M. (2020). Visual imagery and perception share neural representations in the alpha frequency band. *Current Biology*, 30(13), 2621–2627.e5. <https://doi.org/10.1016/j.cub.2020.04.074>
- Xu, J., Wang, C., Xu, Z., Li, T., Chen, F., Chen, K., Gao, J., Wang, J., & Hu, Q. (2020). Specific functional connectivity patterns of middle temporal gyrus subregions in children and adults with autism spectrum disorder. *Autism Research*, 13(3), 410–422. <https://doi.org/10.1002/aur.2239>
- Yamane, Y., Carlson, E. T., Bowman, K. C., Wang, Z., & Connor, C. E. (2008). A neural code for three-dimensional object shape in macaque inferotemporal cortex. *Nature Neuroscience*, 11(11), 1352–1360. <https://doi.org/10.1038/nn.2202>
- Yantis, S., Schwarzbach, J., Serences, J. T., Carlson, R. L., Steinmetz, M. A., Pekar, J. J., & Courtney, S. M. (2002). Transient neural activity in human parietal cortex during spatial attention shifts. *Nature Neuroscience*, 5(10), 995–1002. <https://doi.org/10.1038/nn921>
- Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, 106(3), 1125–1165. <https://doi.org/10.1152/jn.00338.2011>
- Zanto, T. P., Rubens, M. T., Bollinger, J., & Gazzaley, A. (2010). Top-down modulation of visual feature processing: The role of the inferior frontal junction. *NeuroImage*, 53(2), 736–745. <https://doi.org/10.1016/j.neuroimage.2010.06.012>
- Zanto, T. P., Rubens, M. T., Thangavel, A., & Gazzaley, A. (2011). Causal role of the prefrontal cortex in top-down modulation of visual processing and working memory. *Nature Neuroscience*, 14(5), 656–661. <https://doi.org/10.1038/nn.2773>
- Zarjam, P., Epps, J., & Chen, F. (2011). Characterizing working memory load using EEG delta activity. *2011 19th European signal processing conference*, 1554–1558.
- Zhang, X., Mlynaryk, N., Ahmed, S., Japee, S., & Ungerleider, L. G. (2018). The role of inferior frontal junction in controlling the spatially global effect of feature-based attention in human visual areas. *PLoS Biology*, 16(6), e2005399. <https://doi.org/10.1371/journal.pbio.2005399>
- Zhang, Y., Chen, Y., Bressler, S. L., & Ding, M. (2008). Response preparation and inhibition: The role of the cortical sensorimotor beta rhythm. *Neuroscience*, 156(1), 238–246. <https://doi.org/10.1016/j.neuroscience.2008.06.061>
- Zhou, H., & Desimone, R. (2011). Feature-based attention in the frontal eye field and area V4 during visual search. *Neuron*, 70(6), 1205–1217. <https://doi.org/10.1016/j.neuron.2011.04.032>
- Zimmermann, M., Mars, R. B., de Lange, F. P., Toni, I., & Verhagen, L. (2018). Is the extrastriate body area part of the dorsal visuomotor stream? *Brain Structure & Function*, 223(1), 31–46. <https://doi.org/10.1007/s00429-017-1469-0>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Soyuhos, O., & Baldauf, D. (2023). Functional connectivity fingerprints of the frontal eye field and inferior frontal junction suggest spatial versus nonspatial processing in the prefrontal cortex. *European Journal of Neuroscience*, 57(7), 1114–1140. <https://doi.org/10.1111/ejn.15936>