



Research paper

A regularity result for the Fokker–Planck equation with non-smooth drift and diffusion

Paolo Bonicatto ^a, Gennaro Ciampa ^b, Gianluca Crippa ^{c,*}^a Dipartimento di Matematica, Università di Trento, Via Sommarive, 14, 38123 Povo (TN), Italy^b DISIM - Dipartimento di Ingegneria e Scienze dell'Informazione e Matematica, Università degli Studi dell'Aquila, Via Vetoio, 67100 L'Aquila, Italy^c Departement Mathematik und Informatik, Universität Basel, Spiegelgasse 1, CH-4051, Basel, Switzerland

ARTICLE INFO

Communicated by Enrico Valdinoci

MSC:

35A02

35B65

35K57

35Q35

35Q84

Keywords:

Fokker–planck equation

Non-smooth diffusion

Irregular vector field

Regularity and uniqueness

ABSTRACT

The goal of this paper is to study weak solutions of the Fokker–Planck equation. We first discuss existence and uniqueness of weak solutions in an irregular context, providing a unified treatment of the available literature along with some extensions. Then, we prove a regularity result for distributional solutions under suitable integrability assumptions, relying on a new, simple commutator estimate in the spirit of DiPerna–Lions' theory of renormalized solutions for the transport equation. Our result is somehow transverse to Theorem 4.3 of Figally (2008): on the diffusion matrix we relax the assumption of Lipschitz regularity in time at the price of assuming Sobolev regularity in space, and we prove the regularity (and hence the uniqueness) of distributional solutions to the Fokker–Planck equation.

1. Introduction

This paper deals with regularity and uniqueness issues for weak solutions to the Fokker–Planck equation

$$\begin{cases} \partial_t u + \operatorname{div}(bu) - \frac{1}{2} \sum_{i,j} \partial_{ij}(a_{ij}u) = 0 & \text{in } (0, T) \times \mathbb{R}^d, \\ u|_{t=0} = u_0 & \text{in } \mathbb{R}^d, \end{cases} \quad (\text{FP})$$

where $u : (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is the unknown, $b : (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a given vector field, $a = (a_{ij})_{i,j=1,\dots,d} : (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$ is a given diffusion matrix and $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}$ is the initial datum. We are interested here in positive definite fixed diffusivity and not in behaviors for small (or vanishing) diffusivities and we refer to [2,5,8,10,18,20,24] for some recent results with degenerate viscosity coefficient. The Eq. (FP) arises frequently in fluid-dynamics models where u is a passive scalar which is simultaneously advected (by the given velocity field b) and diffused by the second order operator associated with the matrix a , see [25,26]. The Fokker–Planck equation is also intimately connected to the stochastic differential equation

$$\begin{cases} dX_{t,s}(x) = b(s, X_{t,s}(x)) ds + \sigma(s, X_{t,s}(x)) dW_s, \\ X_{t,t}(x) = x, \end{cases} \quad (\text{SDE})$$

* Corresponding author.

E-mail addresses: paolo.bonicatto@unitn.it (P. Bonicatto), gennaro.ciampa@univaq.it (G. Ciampa), gianluca.crippa@unibas.ch (G. Crippa).

where $\sigma : (0, T) \times \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$ is a matrix function and W_s is a d -dimensional Brownian motion on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For sufficiently smooth b and σ , by defining the matrix

$$a(t, x) = \sigma(t, x)\sigma^T(t, x), \quad (1.1)$$

one has that the law of X satisfies Eq. (FP) and the Feynman–Kac representation formula holds

$$u(t, x) := \mathbb{E}[u_0(X_{t,0}(x))], \quad (1.2)$$

where $\mathbb{E}$ denotes the expected value on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Roughly speaking, the connection between (FP) and (SDE) is reminiscent of the one between the linear transport equation and the flow of the vector field b (which can be recovered as the end-point $\sigma = 0$) – see however [12,18,19] and references therein for a more in-depth discussion of these connections. Fokker–Planck equations are also relevant in Statistical Mechanics [13,28], Mean-Field Games theory [6,16,27], and Stochastic Analysis [1,7,12,14,29].

In this paper we are interested in the case in which both the driving vector field and the diffusion matrix are irregular, i.e. we will require only integrability and/or weak differentiability properties on b and a . Besides the purely theoretical interest, this is a common situation in fluid dynamics models: we refer the reader to [18, Section 4] for an overview of modeling polymeric fluids with SDEs and Fokker–Planck equations with irregular coefficients.

The well-posedness of the Cauchy problem in the smooth context is a classical result, see [15]. Out of the smooth setting the main issue are *uniqueness* and *regularity* of solutions. Indeed, existence results can be obtained by a simple approximation argument: under global L^p -bounds on the vector field and the diffusion matrix, one easily establishes energy estimates which allow to apply standard (weak) compactness results. The linearity of the equation ensures that the weak limit is a solution to (FP). At a closer look, however, different a priori estimates are available for (FP) and this is reflected in the presence of various notions of solutions. Understanding the relationships among different notions of solutions is one of the aims of the present work. We recall that this was the subject of study in a recent survey for constant diffusion matrix, see [3].

1.1. Distributional and parabolic solutions

Let $b \in L_t^1 L_x^p$, $a \in L_t^1 L_x^p$, and let $u_0 \in L_x^q$ for some $1 \leq p, q \leq \infty$ such that $1/p + 1/q \leq 1$. With these assumptions it is possible to introduce distributional solutions to (FP), i.e. functions $u \in L_t^\infty L_x^q$ solving the equation in the sense of distributions. Define the vector field

$$\tilde{b} := b - \frac{1}{2} \sum_j \partial_j a_{jj}. \quad (1.3)$$

Note that, formally, Eq. (FP) is equivalent to

$$\partial_t u + \operatorname{div}(\tilde{b}u) - \frac{1}{2} \sum_{i,j} \partial_i(a_{ij} \partial_j u) = 0, \quad (\text{FP-div})$$

which is also known as the Fokker–Planck equation in *divergence form*. Then, we consider the following set of assumptions:

Assumptions

(A1) the diffusion matrix a is uniformly bounded, i.e.

$$a \in L^\infty((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^{d \times d}));$$

(A2) the diffusion matrix a satisfies

$$\sum_{j=1}^d \partial_j a_{ij} \in L^\infty((0, T) \times \mathbb{R}^d), \quad \text{for all } i \in \{1, \dots, d\};$$

(A3) the vector field $\tilde{b}$ satisfies

$$(\operatorname{div} \tilde{b})^- = \left(\operatorname{div} b - \frac{1}{2} \sum_{i,j=1}^d \partial_i a_{ij} \right)^- \in L^\infty((0, T) \times \mathbb{R}^d);$$

(A4) the diffusion matrix is uniformly elliptic, i.e. there exists $\alpha > 0$ such that for every $\xi \in \mathbb{R}^d$ it holds

$$\langle \xi, a(t, x) \xi \rangle \geq \alpha |\xi|^2, \quad \text{for a.e. } (t, x) \in (0, T) \times \mathbb{R}^d.$$

One a priori estimate for smooth solutions is the following

$$\|u(t, \cdot)\|_{L^q}^q \leq \|u_0\|_{L^q}^q + (q-1) \|(\operatorname{div} \tilde{b})^-\|_\infty \int_0^t \|u(s, \cdot)\|_{L^q}^q ds, \quad (1.4)$$

and a standard approximation argument provides the existence of a distributional solution to (FP). However, distributional solutions are not, in general, unique, not even in the case of constant coefficient diffusion matrices, see [21–23]. This motivates the introduction of another notion of solution. Exploiting the presence of the second order operator in the equation, one can show another energy estimate for smooth solutions, namely

$$\|u(t, \cdot)\|_{L^2}^2 + \alpha \int_0^t \int_{\mathbb{R}^d} |\nabla u(s, x)|^2 dx ds \leq \|u_0\|_{L^2}^2 - \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \tilde{b}(s, x) |u(s, x)|^2 dx ds$$

for every $t \in [0, T]$, which suggests that one should look for solutions possessing L^2 gradient, i.e. solutions that are H^1 in the space variable. We therefore say that a distributional solution $u \in L_t^\infty L_x^q$ to (FP) is *parabolic* if it holds $u \in L_t^2 H_x^1$.

Parabolic solutions carry the exact regularity needed to establish their uniqueness under suitable integrability assumptions on b and a . In the setting of rough drift/diffusion coefficients, two uniqueness results for parabolic solutions are available under slightly different assumptions:

- in [12] it is shown that a parabolic solution $u \in L_t^\infty L_x^2 \cap L_t^2 H_x^1$ is unique provided that $u_0 \in L_x^2$ and $b \in L_{t,x}^\infty$ satisfy (A1), (A2), (A3) and (A4);
- in [18] it is proved that a parabolic solution $u \in L_t^\infty (L_x^2 \cap L_x^\infty) \cap L_t^2 H_x^1$ is unique in the set

$$\{u \in L_t^2 H_x^1 : \sigma^T \nabla u \in L_{t,x}^2\},$$

where σ is defined accordingly to (1.1), provided that

- $u_0 \in L_x^2 \cap L_x^\infty$;
- $b \in L_t^2 L_{x,\text{loc}}^\infty$ satisfy the growth assumption $\frac{b}{1+|x|} \in L_t^1 L_x^1 + L_t^1 L_x^\infty$;
- the matrix $\sigma \sigma^T$ is uniformly positive definite;
- $\sigma \in L_t^\infty L_{x,\text{loc}}^\infty$ satisfy the growth assumption $\frac{\sigma}{1+|x|} \in L_t^2 L_x^2 + L_t^2 L_x^\infty$;
- $\operatorname{div} \tilde{b} \in L_t^1 L_x^\infty$.

The main difference between the two results lies in the L^∞ assumption: [12] assumes $b, a \in L_{t,x}^\infty$, while [18] assumes $u_0 \in L_x^\infty$. Moreover, the proofs of these two results are based on different techniques: in [12, Theorem 4.3] the author uses a functional analytic approach closer to the one commonly used for elliptic/parabolic equations which make use of a Lax–Milgram type argument; on the other hand, the result in [18] is more “hyperbolic” in nature resorting to commutator estimates and on the theory of renormalized solutions as done in [9] for the linear transport equation. In this regards, the $L_t^2 H_x^1$ regularity of the solution allows to obtain a better control on the so called *commutator*, which measures the error one commits when considering smooth approximations of the solution. In particular, in [17,18] it is shown that the commutator for parabolic solutions converges strongly to 0 in $L_{t,x}^1$, but better bounds can be established for parabolic solutions $u \in L_t^2 H_x^1$ if $b \in L_t^2 L_x^2$ (see Section 2 for a comprehensive list of old and new commutator estimates).

Relying on these techniques, in Section 3 we provide a presentation of the available results on the well-posedness of (FP) together with some technical extensions.

We make use of the new commutator estimates to prove the uniqueness of parabolic solutions of (FP) in a setting that incorporates and generalizes those of [12,18], thus providing a unified treatment of the existing literature. Our theorem is the following.

Theorem 1.1. *Let $u_0 \in L^2 \cap L^q(\mathbb{R}^d)$ be a given initial datum and assume that*

- (i) $b \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ with $1/p + 1/q = 1/2$;
- (ii) b satisfies the growth condition
$$\frac{b}{1+|x|} \in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d));$$
- (iii) a satisfies (A1), (A2) and (A4);
- (iv) the vector field $\tilde{b}$ defined in (1.3) satisfies
$$(\operatorname{div} \tilde{b})^- \in L^1((0, T); L^\infty(\mathbb{R}^d)).$$

Then, there exists a unique parabolic solution u to (FP).

Note that if $p = \infty$ and $q = 2$ we precisely recover the existence and uniqueness of parabolic solutions to (FP) in [12, Theorem 4.3], while when $p = 2$ and $q = \infty$ we obtain *essentially* the corresponding result in [18] – see however Section 3.1.1 below for a detailed comparison between the two results.

1.2. Regularity and uniqueness of distributional solutions

According to our definitions, parabolic solutions cannot always be defined, but if they can, then they are always distributional. The converse implication is, in general, not true: in [21] it is shown that there exist infinitely many distributional solutions $u \in L_t^\infty L_x^2$ to (FP) with a divergence-free vector field $b \in L_t^\infty L_x^2$ and $a = \operatorname{Id}$, while the parabolic one is unique. This motivates the search for

a condition that guarantees *parabolic regularity* of a distributional solution. In addition to the uniqueness of parabolic solutions, in [12] it is proved that the condition

$$\partial_t a_{ij} \in L^\infty((0, T) \times \mathbb{R}^d)$$

for $i, j = 1, \dots, d$, implies that every solution $u \in L^2((0, T) \times \mathbb{R}^d)$ to (FP) belongs to the space

$$Y := \{u \in L^2((0, T); H^1(\mathbb{R}^d)) : \partial_t u \in L^2((0, T); H^{-1}(\mathbb{R}^d))\}.$$

The idea of the proof is simple: once the solutions in Y are proved to be unique, uniqueness in the functional space $L^2_{t,x}$ follows from a regularity argument which makes use of the condition $\partial_t a \in L^\infty_{t,x}$. Our purpose is to investigate the role of the assumption on the time derivative of a . In this regard, we show that if we replace $\partial_t a_{ij} \in L^\infty_{t,x}$ with some Sobolev regularity in space, distributional solutions are actually more regular. Our theorem reads as follows:

Theorem 1.2. *Let $p, q \geq 1$ such that $1/p + 1/q \leq 1/2$. Assume that $b \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ and $a \in L^2((0, T); W^{1,p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ satisfy (A2), (A4), and $(\operatorname{div} \tilde{b})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Let $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ be a distributional solution to (FP), then $u \in L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$.*

The proof relies again on a new commutator estimate: we show that, in the current regime, the convergence to zero of the commutators takes place in $L^2_t H^{-1}_{x,\text{loc}}$ and this, together with the energy estimate, implies the regularity of distributional solutions. Moreover, in the particular case $p = \infty, q = 2$ we obtain the uniqueness of distributional solutions in $L^2_{t,x}$ – notice that here we need the stronger assumption (A3) instead of the weaker $L^1_t L^\infty_x$ control on $(\operatorname{div} \tilde{b})^-$.

Theorem 1.3. *Assume that $b \in L^\infty((0, T); L^\infty(\mathbb{R}^d))$ and that (A1), (A2), (A3) and (A4) are fulfilled. Moreover, assume that $\nabla a \in L^\infty((0, T) \times \mathbb{R}^d)$. Then, any distributional solution $u \in L^2((0, T) \times \mathbb{R}^d)$ of (FP) belongs to Y and thus it is unique.*

The theorem above is somehow transverse to [12, Theorem 4.8], i.e. we replace the assumption on the time derivative $\partial_t a \in L^\infty_{t,x}$ with an assumption on the spatial gradient $\nabla a \in L^\infty_{t,x}$. Note that we need the L^∞_t assumption on ∇a to compensate the L^2_t assumption on u .

Furthermore, the local regularity in space H^1_{loc} provided by Theorem 1.2 suggests investigating whether uniqueness of solutions to (FP) holds in this larger functional space. We will say that a distributional solution $u \in L^\infty_t L^q_x$ to (FP) is *locally parabolic* if it holds $u \in L^2_t H^1_{x,\text{loc}}$. Assuming suitable growth conditions on b, a (similarly to [9,18]), we prove that distributional solutions are locally parabolic and unique in the class $L^\infty((0, T); L^q(\mathbb{R}^d))$:

Theorem 1.4. *Let $u_0 \in L^q \cap L^\infty(\mathbb{R}^d)$ be a given initial datum for some $q > 2$ and assume that $b \in L^2((0, T); L^p(\mathbb{R}^d))$ and $a \in L^2((0, T); W^{1,p}(\mathbb{R}^d))$ with $1/p + 1/q = 1/2$ satisfy (A2) and (A4). Moreover, we assume the following growth conditions*

$$\frac{b(t, x)}{1 + |x|}, \frac{\partial_j a_{ij}(t, x)}{1 + |x|}, \frac{a(t, x)}{1 + |x|^2} \in L^1((0, T); L^1(\mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d)). \quad (1.5)$$

Then, there exists at most one distributional solution $u \in L^\infty((0, T); L^q(\mathbb{R}^d)) \cap L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$.

In the proof we use the assumptions on p, q and the Regularity Theorem 1.2 to show the existence of a solution in $L^\infty((0, T); L^q(\mathbb{R}^d)) \cap L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$. Then we exploit the growth conditions on b and a to use a Gronwall type argument following the same strategy of Theorem 1.1. Finally, we show how we can play with the integrability exponents in the growth conditions to guarantee that a distributional solution is in $L^2((0, T); H^1(\mathbb{R}^d))$, obtaining the following result:

Theorem 1.5. *Let $u \in L^\infty((0, T); L^2 \cap L^q(\mathbb{R}^d))$ be a distributional solution of (FP), and let $b \in L^2((0, T); L^p(\mathbb{R}^d))$ and $a \in L^2((0, T); W^{1,p}(\mathbb{R}^d))$ with $1/p + 1/q = 1/2$ satisfy (A2) and (A4). Moreover, assume that $2 < q \leq \frac{2d}{d-2}$ together with the following growth conditions*

$$\frac{b(t, x)}{1 + |x|}, \frac{\partial_j a_{ij}(t, x)}{1 + |x|}, \frac{a(t, x)}{1 + |x|^2} \in L^1((0, T); L^{\frac{q}{q-2}}(\mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d)).$$

Then $u \in L^2((0, T); H^1(\mathbb{R}^d))$.

We conclude this introduction by observing that the well-posedness landscape of (FP) in the non-smooth regime is far from being completely understood. Recently, several authors have devoted their attention to the construction of *exotic* non-unique distributional solutions to Fokker–Planck type equations (and also for related PDEs in fluid-dynamics). These counterexamples are mainly provided in the case of constant diffusion matrix — we refer the reader to our recent survey [3] for a thorough analysis of the problem and for additional references. From the mathematical point of view, handling the PDE in the presence of irregular diffusions is even more delicate. At the same time, anisotropic diffusions naturally appear in the physical models and therefore the study of (FP) in this setting is of paramount importance.

1.3. Structure of the paper

In Section 2 we collect the commutator estimates which will then be the heart of all the proofs that follow. In Section 3 we recall the definitions of distributional and parabolic solutions of (FP) and we prove their existence together with the uniqueness of parabolic solutions. These results are technical improvements of [12,18], providing a unified treatment. Finally, in Section 4 we prove our Regularity Theorems 1.2 and 1.3. Moreover, we discuss the role of the growth conditions, proving Theorems 1.4 and 1.5.

In what follows, we will often omit the summation symbol and we will adopt the convention of summing over repeated indices.

2. Commutator estimates

In this section we collect commutator estimates that will play a predominant role in the entire paper. First, we recall some known results on the L^1 convergence of the commutator. We provide the proofs for completeness but they can be found in [9,18,19]. We remark that the convergence of the commutators is the key tool to prove the renormalization property (and hence the uniqueness) of distributional solutions of the linear transport equation, see [9]. To fix the notation, let $\rho \in C_c^\infty(\mathbb{R}^d)$ be a smooth convolution kernel such that

$$\int_{\mathbb{R}^d} \rho(x) dx = 1,$$

and for any $\delta > 0$ define ρ^δ as

$$\rho^\delta(x) := \frac{1}{\delta^d} \rho\left(\frac{x}{\delta}\right). \quad (2.1)$$

2.1. Commutator estimates in L^1

We start by proving some classical commutator estimates. The following lemma holds.

Lemma 2.1. *Let $\mathbf{b} \in L^2((0, T); L_{\text{loc}}^p(\mathbb{R}^d; \mathbb{R}^d))$ be a vector field and let $w \in L^\infty((0, T); L_{\text{loc}}^q(\mathbb{R}^d))$, where p, q are positive real numbers with $1/p + 1/q \leq 1$. Let $(\rho^\delta)_\delta$ be a family of smooth convolution kernels. Define the commutator of w and $\mathbf{b}$ as follows:*

$$r^\delta := \text{div}(\mathbf{b} w^\delta) - \text{div}(\mathbf{b} w) * \rho^\delta = r_1^\delta + r_2^\delta, \quad (2.2)$$

where

$$r_1^\delta := \mathbf{b} \cdot \nabla(w * \rho^\delta) - (\mathbf{b} \cdot \nabla w) * \rho^\delta, \quad r_2^\delta := w * \rho^\delta \text{div} \mathbf{b} - (w \text{div} \mathbf{b}) * \rho^\delta. \quad (2.3)$$

Then we have the following:

- if $\nabla w \in L^2((0, T); L_{\text{loc}}^q(\mathbb{R}^d; \mathbb{R}^d))$, then r_1^δ converges to 0 in $L^1((0, T); L_{\text{loc}}^1(\mathbb{R}^d))$;
- if $\text{div} \mathbf{b} \in L^1((0, T); L^\infty(\mathbb{R}^d))$, then r_2^δ converges to 0 in $L^1((0, T); L_{\text{loc}}^q(\mathbb{R}^d))$.

Proof. We start by showing the convergence of r_1^δ . Observe that, for a.e. $t \in (0, T)$ and a.e. $x \in \mathbb{R}^d$, we can explicitly write the commutator r_1^δ in the following form:

$$\begin{aligned} r_1^\delta(t, x) &= [\mathbf{b} \cdot \nabla(w * \rho^\delta)](t, x) - [(\mathbf{b} \cdot \nabla w) * \rho^\delta](t, x) \\ &= \mathbf{b}(t, x) \cdot \nabla \int_{\mathbb{R}^d} w(t, x - y) \rho^\delta(y) dy - \int_{\mathbb{R}^d} \mathbf{b}(t, x - y) \cdot \nabla w(t, x - y) \rho^\delta(y) dy \\ &= \int_{\mathbb{R}^d} \rho^\delta(y) (\mathbf{b}(t, x) - \mathbf{b}(t, x - y)) \cdot \nabla w(t, x - y) dy \\ &= \int_{\mathbb{R}^d} \rho(z) (\mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)) \cdot \nabla w(t, x - \delta z) dz. \end{aligned}$$

Let Q be any bounded subset of $\mathbb{R}^d$, we have that

$$\begin{aligned} \iint_{(0, T) \times Q} |r_1^\delta(t, x)| dt dx &= \iint_{(0, T) \times Q} \left| \int_{\mathbb{R}^d} \rho(z) (\mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)) \cdot \nabla w(t, x - \delta z) dz \right| dt dx \\ &\leq \int_{B_1} \rho(z) \int_0^T \int_Q |\mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)| |\nabla w(t, x - \delta z)| dx dz dt. \end{aligned} \quad (2.4)$$

Since $(t, x) \mapsto \mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)$ converges to 0 in measure (for every fixed z), the conclusion follows by the Dominated Convergence Theorem.

Now we consider the commutator r_2^δ : observe that $w * \rho^\delta \rightarrow w$ strongly in $L^\infty L_{\text{loc}}^q$ and similarly $(w \text{div} \mathbf{b}) * \rho^\delta \rightarrow w \text{div} \mathbf{b}$ strongly in $L^\infty L_{\text{loc}}^q$ and this concludes the proof. $\square$

Remark 2.2. It is worth observing that no assumption on the divergence of $\mathbf{b}$ is necessary for the convergence of r_1^δ .

Lemma 2.3. *Let $w \in L^2((0, T); W_{\text{loc}}^{1, q}(\mathbb{R}^d))$ and let $\mathbf{a} \in L^2((0, T); W_{\text{loc}}^{1, p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ be a matrix-valued function, where p, q are positive real numbers with $1/p + 1/q \leq 1$. Let $(\rho^\delta)_\delta$ be a family of smooth convolution kernels defined as in (2.1). Define the commutator of w and $\mathbf{a}$ as follows:*

$$s^\delta := \sum_{i, j} [\partial_{ij}(a_{ij} w)] * \rho^\delta - \partial_{ij}[a_{ij}(w * \rho^\delta)]. \quad (2.5)$$

Then s^δ converges to 0 in $L^1((0, T); L_{\text{loc}}^1(\mathbb{R}^d))$.

Proof. We can rewrite the commutator s^δ as follows

$$\begin{aligned} s^\delta &= \sum_{i,j} [\partial_{ij}(a_{ij}w)] * \rho_\delta - \partial_{ij}[a_{ij}(w * \rho_\delta)] \\ &= \sum_{i,j} [\partial_i(a_{ij}\partial_j w)] * \rho_\delta - \partial_i[a_{ij}\partial_j(w * \rho_\delta)] + \sum_{i,j} [\partial_i(\partial_j a_{ij}w)] * \rho_\delta - \partial_i[\partial_j a_{ij}(w * \rho_\delta)] \\ &:= s_1^\delta + s_2^\delta. \end{aligned}$$

We start by analyzing s_1^δ : for a.e. $(t, x) \in (0, T) \times \mathbb{R}^d$ we can write

$$\begin{aligned} s_1^\delta(t, x) &= \sum_{i,j} [\partial_i(a_{ij}\partial_j w)] * \rho_\delta(t, x) - \partial_i[a_{ij}\partial_j(w * \rho_\delta)](t, x) \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \partial_i[a_{ij}(t, x - y)\partial_j w(t, x - y)]\rho_\delta(y)dy - \partial_i \left(a_{ij}(t, x)\partial_j \int_{\mathbb{R}^d} w(t, x - y)\rho_\delta(y)dy \right) \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \partial_i[(a_{ij}(t, x - y) - a_{ij}(t, x))\partial_j w(t, x - y)]\rho_\delta(y)dy \\ &= - \sum_{i,j} \int_{\mathbb{R}^d} \frac{a_{ij}(t, x - \delta z) - a_{ij}(t, x)}{\delta} \partial_j w(t, x - \delta z)\partial_i \rho_\delta(z)dz \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \int_0^1 \partial_j w(t, x - \delta z)\nabla a_{ij}(t, x - s\delta z) \cdot z \partial_i \rho_\delta(z)dsdz, \end{aligned}$$

where in the fourth line we changed variables and integrated by parts. Then, for a.e. (t, x) the commutator s_1^δ converges to

$$\lim_{\delta \rightarrow 0} s_1^\delta(t, x) = \sum_{i,j} \partial_j w(t, x)\nabla a_{ij}(t, x) \cdot \int_{\mathbb{R}^d} z \partial_i \rho(z)dz = - \sum_{i,j} \partial_j w(t, x)\partial_i a_{ij}(t, x) \int_{\mathbb{R}^d} \rho(z)dz. \quad (2.6)$$

We now focus on s_2^δ : we have that

$$\begin{aligned} s_2^\delta(t, x) &= \sum_{i,j} [\partial_i(\partial_j a_{ij}w)] * \rho_\delta(t, x) - \partial_i[\partial_j a_{ij}(w * \rho_\delta)](t, x) \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \partial_i[\partial_j a_{ij}(t, x - y)w(t, x - y)]\rho_\delta(y)dy - \partial_i \left(\partial_j a_{ij}(t, x) \int_{\mathbb{R}^d} w(t, x - y)\rho_\delta(y)dy \right) \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \partial_i[\partial_j(a_{ij}(t, x - y) - a_{ij}(t, x))w(t, x - y)]\rho_\delta(y)dy \\ &= - \sum_{i,j} \int_{\mathbb{R}^d} \frac{\partial_j a_{ij}(t, x - \delta z) - \partial_j a_{ij}(t, x)}{\delta} w(t, x - \delta z)\partial_i \rho_\delta(z)dz \\ &= \sum_{i,j} \int_{\mathbb{R}^d} \frac{a_{ij}(t, x - \delta z) - a_{ij}(t, x)}{\delta} \partial_j w(t, x - \delta z)\partial_i \rho_\delta(z)dz + \sum_{i,j} \int_{\mathbb{R}^d} \frac{a_{ij}(t, x - \delta z) - a_{ij}(t, x)}{\delta} w(t, x - \delta z)\partial_{ij} \rho_\delta(z)dz \\ &= \underbrace{- \sum_{i,j} \int_{\mathbb{R}^d} \int_0^1 \partial_j w(t, x - \delta z)\nabla a_{ij}(t, x - s\delta z) \cdot z \partial_i \rho_\delta(z)dsdz}_{I^\delta} - \underbrace{\sum_{i,j} \int_{\mathbb{R}^d} \int_0^1 w(t, x - \delta z)\nabla a_{ij}(t, x - s\delta z) \cdot z \partial_{ij} \rho_\delta(z)dsdz}_{II^\delta}. \end{aligned}$$

Thus, one can check the following a.e. convergences

$$\lim_{\delta \rightarrow 0} I^\delta(t, x) = \sum_{i,j} \partial_j w(t, x)\partial_i a_{ij}(t, x) \int_{\mathbb{R}^d} \rho(z)dz, \quad (2.7)$$

and

$$\lim_{\delta \rightarrow 0} II^\delta(t, x) = -w(t, x) \sum_{i,j} \nabla a_{ij}(t, x) \cdot \int_{\mathbb{R}^d} z \partial_{ij} \rho(z)dsdz = 0. \quad (2.8)$$

We have thus shown that $s^\delta \rightarrow 0$ for a.e. $(t, x) \in (0, T) \times \mathbb{R}^d$. The convergence in $L^1((0, T); L^1_{\text{loc}}(\mathbb{R}^d))$ follows from the dominated convergence theorem. $\square$

2.2. Commutator estimates on negative Sobolev norms

We now show some new commutator estimates which will allow us to prove the regularity of a distributional solution within an appropriate integrability range.

Lemma 2.4. Let $b \in L^2((0, T); L^p_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d))$ be a vector field and let $w \in L^\infty((0, T); L^q_{\text{loc}}(\mathbb{R}^d))$, where p, q are positive real numbers with $1/p + 1/q \leq 1/2$. Let $(\rho^\delta)_\delta$ be a family of smooth convolutions kernels and define r^δ as in (2.2). Then r^δ converges to 0 in $L^2((0, T); H^{-1}_{\text{loc}}(\mathbb{R}^d))$.

Proof. We recall that the commutator r^δ is given by

$$r^\delta = \operatorname{div}[\mathbf{b}(w * \rho^\delta)] - [\operatorname{div}(\mathbf{b}w) * \rho^\delta] = \operatorname{div}[\mathbf{b}(w * \rho^\delta) - (\mathbf{b}w) * \rho^\delta],$$

in the sense of distributions on $(0, T) \times \mathbb{R}^d$. We can thus write

$$r^\delta(t, x) = \operatorname{div}_x \left(\int_{\mathbb{R}^d} [\mathbf{b}(t, x) - \mathbf{b}(t, x - y)] w(t, x - y) \rho^\delta(y) dy \right)$$

and, for a given bounded subset Q of $\mathbb{R}^d$ we can estimate

$$\begin{aligned} \|r^\delta\|_{L^2(H^{-1}(Q))} &= \sup_{\|\varphi\|_{L^2 H_0^1(Q)} \leq 1} \left| \iint_{(0,T) \times Q} r^\delta(t, x) \varphi(t, x) dt dx \right| \\ &= \sup_{\|\varphi\|_{L^2 H_0^1(Q)} \leq 1} \left| \iint_{(0,T) \times Q} \left(\int_{\mathbb{R}^d} [\mathbf{b}(t, x) - \mathbf{b}(t, x - y)] w(t, x - y) \rho^\delta(y) dy \right) \cdot \nabla \varphi(t, x) dt dx \right| \\ &\leq \sup_{\|\varphi\|_{L^2 H_0^1(Q)} \leq 1} \int_{B_1} \rho(z) \int_0^T \int_Q |\mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)| |w(t, x - \delta z)| |\nabla \varphi(t, x)| dx dz. \end{aligned}$$

Notice now that, as in the proof of [Lemma 2.1](#), the map $(t, x) \mapsto \mathbf{b}(t, x) - \mathbf{b}(t, x - \delta z)$ converges to 0 in measure (for every fixed z). Hölder inequality on the product space $(0, T) \times Q$ with exponents $(p, q, 2)$ (in space) and $(2, \infty, 2)$ (in time) allows us to apply Lebesgue Dominated Convergence Theorem and we can therefore conclude that $r^\delta \rightarrow 0$ in $L^2(H_{\text{loc}}^{-1})$. $\square$

Remark 2.5. Notice that with global assumptions on $\mathbf{b}$ and w , in the case $1/p + 1/q = 1/2$ the commutator r^δ converges to 0 in $L^2((0, T); H^{-1}(\mathbb{R}^d))$.

We now prove the analogous result on the diffusion matrix.

Lemma 2.6. Let $\mathbf{a} \in L^2((0, T); W_{\text{loc}}^{1,p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ be a matrix valued function and let $w \in L^\infty((0, T); L_{\text{loc}}^q(\mathbb{R}^d))$ with p, q are positive real numbers with $1/p + 1/q \leq 1/2$. Let $(\rho^\delta)_\delta$ be a family of smooth convolutions kernels and define s^δ as in [\(2.5\)](#). Then s^δ converge to 0 in $L^2((0, T); H_{\text{loc}}^{-1}(\mathbb{R}^d))$.

Proof. Let Q be any bounded subset of $\mathbb{R}^d$. Then, for any $\varphi \in L^2((0, T); H_0^1(Q))$ we compute

$$\begin{aligned} \iint_{(0,T) \times Q} s^\delta(t, x) \varphi(t, x) dt dx &= \iint_{(0,T) \times Q} \left(\sum_{i,j} [\partial_{ij}(a_{ij}w)] * \rho_\delta - \partial_{ij}[a_{ij}(w * \rho_\delta)] \right) \varphi(t, x) dt dx \\ &= \sum_{i,j} \iint_{(0,T) \times Q} \partial_{ij} \left(\int_{\mathbb{R}^d} a_{ij}(t, y) w(t, y) \rho_\delta(x - y) dy - a_{ij}(t, x) \int_{\mathbb{R}^d} w(t, y) \rho_\delta(x - y) dy \right) \varphi(t, x) dt dx \\ &= - \sum_{i,j} \iint_{(0,T) \times Q} \partial_j \left(\int_{\mathbb{R}^d} a_{ij}(t, y) w(t, y) \rho_\delta(x - y) dy - a_{ij}(t, x) \int_{\mathbb{R}^d} w(t, y) \rho_\delta(x - y) dy \right) \partial_i \varphi(t, x) dt dx \\ &= - \sum_{i,j} \int_0^T \int_{\mathbb{R}^d} \int_Q \partial_i \varphi(t, x) w(t, y) \frac{1}{\delta^{d+1}} \partial_j \rho \left(\frac{x - y}{\delta} \right) (a_{ij}(t, y) - a_{ij}(t, x)) dx dy dt \\ &\quad + \sum_i \int_0^T \int_Q \int_{\mathbb{R}^d} w(t, y) \rho_\delta(x - y) \partial_i \varphi(t, x) \left(\sum_j \partial_j a_{ij}(t, x) \right) dy dx dt, \end{aligned}$$

and by using a change of variables and Fubini to exchange integrals we finally get that

$$\begin{aligned} \iint_{(0,T) \times Q} s^\delta(t, x) \varphi(t, x) dt dx &= - \sum_i \int_0^T \int_{\mathbb{R}^d} \int_Q \partial_i \varphi(t, x) w(t, x - \delta z) \sum_j \left(\partial_j \rho(z) \left(\frac{a_{ij}(t, x - \delta z) - a_{ij}(t, x)}{\delta} \right) \right) dx dz dt \\ &\quad + \sum_i \int_0^T \int_Q \int_{\mathbb{R}^d} w(t, x - \delta z) \rho(z) \partial_i \varphi(t, x) \left(\sum_j \partial_j a_{ij}(t, x) \right) dz dx dt. \end{aligned}$$

Thus, by letting $\delta \rightarrow 0$ we obtain

$$\begin{aligned} \lim_{\delta \rightarrow 0} \iint_{(0,T) \times Q} s^\delta(t, x) \varphi(t, x) dt dx &= \sum_{i,j} \int_0^T \int_{\mathbb{R}^d} \int_Q \partial_i \varphi(t, x) w(t, x) \partial_j \rho(z) \nabla a_{ij}(t, x) \cdot z dx dz dt \\ &\quad + \sum_{ij} \int_0^T \int_Q \int_{\mathbb{R}^d} w(t, x) \rho(z) \partial_i \varphi(t, x) \partial_j a_{ij}(t, x) dz dx dt \end{aligned}$$

$$\begin{aligned}
 &= - \sum_{i,j} \int_0^T \int_{\mathbb{R}^d} \int_Q \partial_i \varphi(t, x) w(t, x) \rho(z) \nabla a_{ij}(t, x) \cdot e_j dx dz dt \\
 &\quad + \sum_{ij} \int_0^T \int_Q \int_{\mathbb{R}^d} w(t, x) \rho(z) \partial_i \varphi(t, x) \partial_j a_{ij}(t, x) dz dx dt = 0,
 \end{aligned}$$

where in the last step we have integrated by parts in the z -variable. This concludes the proof. $\square$

We conclude this section with the following commutator estimate: it will be crucial to recover the uniqueness of parabolic solutions in the setting of [12].

Lemma 2.7. *Let $\mathbf{a} \in L^\infty((0, T); L^\infty_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ be a matrix valued function and let $\nabla w \in L^2((0, T); L^2_{\text{loc}}(\mathbb{R}^d))$. Let $(\rho^\delta)_\delta$ be a family of smooth convolutions kernels and consider the commutator s_1^δ as follows*

$$s_1^\delta := \sum_i \partial_i \left(\sum_j a_{ij} \partial_j w^\delta - (a_{ij} \partial_j w) * \rho^\delta \right). \quad (2.9)$$

Then s_1^δ converges to 0 in $L^2((0, T); H^{-1}_{\text{loc}}(\mathbb{R}^d))$.

Proof. The proof is very similar to the one of Lemma 2.4 and relies on the divergence form of the commutator. Given a bounded subset Q of $\mathbb{R}^d$, we can estimate

$$\begin{aligned}
 \|s_1^\delta\|_{L^2(H^{-1}(Q))} &= \sup_{\|\varphi\|_{L^2 H^1_0(Q)} \leq 1} \left| \iint_{(0,T) \times Q} s_1^\delta(t, x) \varphi(t, x) dx dt \right| \\
 &= \sup_{\|\varphi\|_{L^2 H^1_0(Q)} \leq 1} \left| \iint_{(0,T) \times Q} \sum_i \left(\sum_j a_{ij} \partial_j w^\delta - (a_{ij} \partial_j w) * \rho^\delta \right) \partial_i \varphi(t, x) dx dt \right| \\
 &\leq \sup_{\|\varphi\|_{L^2 H^1_0(Q)} \leq 1} \int_{B_1} \rho(z) \int_0^T \int_Q |\mathbf{a}(t, x) - \mathbf{a}(t, x - \delta z)| |\nabla w(t, x - \delta z)| |\nabla \varphi(t, x)| dx dz dt.
 \end{aligned}$$

As in the proof of Lemma 2.1, the map $(t, x) \mapsto \mathbf{a}(t, x) - \mathbf{a}(t, x - \delta z)$ converges to 0 in measure (for every fixed z) and then by Hölder inequality on the product space $(0, T) \times Q$ with exponents $(\infty, 2)$ we can apply Lebesgue Dominated Convergence Theorem obtaining that $s_1^\delta \rightarrow 0$ in $L^2((0, T); H^{-1}_{\text{loc}}(\mathbb{R}^d))$. $\square$

Remark 2.8. It is worth to note that the commutator s_1^δ does not converge pointwise a.e. to 0, as shown in (2.6). However it converges in the sense of the distributions and, by density, in the functional space $L^2_t H^{-1}_{\text{loc}}$.

3. Well-posedness of the Cauchy problem

In this section we collect some results on the Cauchy problem for the Fokker–Planck equation (FP). Our aim is to give a complete presentation of the available results and to provide technical extensions. In this regard, we first analyze the Fokker–Planck equation in divergence form, providing some technical improvements of the results in [18]. Then we move to the analysis of the Fokker–Planck equation, proving existence and uniqueness in a slightly more general setting than the one considered in [12]. Theorem 3.13 somehow interpolates the results from [12, 18] using the $L^2 H^{-1}_{\text{loc}}$ convergence of the commutators.

3.1. The equation in divergence form

In this section we study the initial value problem

$$\begin{cases} \partial_t u + \operatorname{div}(\tilde{\mathbf{b}}u) - \frac{1}{2} \sum_{i,j} \partial_i (a_{ij} \partial_j u) = 0 & \text{in } (0, T) \times \mathbb{R}^d, \\ u|_{t=0} = u_0 & \text{in } \mathbb{R}^d, \end{cases} \quad (\text{FP-div})$$

which is the Cauchy problem for the Fokker–Planck equation in divergence form. Requiring only integrability assumptions on $\tilde{\mathbf{b}}$ and $\mathbf{a}$ we can give the following definition of solutions to (FP-div).

Definition 3.1. Assume that

$$\tilde{\mathbf{b}} \in L^1((0, T); L^2(\mathbb{R}^d; \mathbb{R}^d)), \quad \mathbf{a} \in L^2((0, T); L^2(\mathbb{R}^d; \mathbb{R}^{d \times d})), \quad u_0 \in L^2(\mathbb{R}^d). \quad (3.1)$$

A function $u \in L^\infty((0, T); L^2(\mathbb{R}^d)) \cap L^2((0, T); H^1(\mathbb{R}^d))$ is a *parabolic solution* to (FP-div) if for any $\varphi \in C_c^\infty([0, T) \times \mathbb{R}^d)$ the following identity holds:

$$\int_0^T \int_{\mathbb{R}^d} u \left(\partial_t \varphi + \tilde{\mathbf{b}} \cdot \nabla \varphi \right) - \frac{1}{2} \sum_{ij} a_{ij} \partial_j u \partial_i \varphi dx dt + \int_{\mathbb{R}^d} u_0 \varphi(0, \cdot) dx = 0.$$

The following theorem holds.

Theorem 3.2. Let $p, q \in [2, \infty]$ be such that $1/p + 1/q = 1/2$. Assume that:

- (i) $\tilde{\mathbf{b}} \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$;
- (ii) $\tilde{\mathbf{b}}$ satisfies the growth condition $\frac{\tilde{\mathbf{b}}}{1+|x|} := \tilde{\mathbf{b}}_1 + \tilde{\mathbf{b}}_2$ with

$$\tilde{\mathbf{b}}_1 \in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^d)), \quad \tilde{\mathbf{b}}_2 \in L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d)); \quad (3.2)$$

- (iii) $(\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$;
- (iv) $\mathbf{a}$ satisfies (A1) and (A4);
- (v) $u_0 \in L^2 \cap L^q(\mathbb{R}^d)$.

Then, there exists a unique parabolic solution u to (FP-div).

Proof. Let $(\rho^\delta)_\delta$ be a standard family of mollifiers as in (2.1) and let us define $\tilde{\mathbf{b}}^\delta := \tilde{\mathbf{b}} * \rho^\delta$, $\mathbf{a}^\delta := \mathbf{a} * \rho^\delta$, and $u_0^\delta = u_0 * \rho^\delta$. Then, we consider the approximating problem

$$\begin{cases} \partial_t u^\delta + \operatorname{div}(\tilde{\mathbf{b}}^\delta u^\delta) - \frac{1}{2} \sum_{i,j} \partial_i(a_{ij}^\delta \partial_j u^\delta) = 0, \\ u^\delta(0, \cdot) = u_0^\delta. \end{cases} \quad (3.3)$$

The Eq. (3.3) is classically well-posed, so we can consider for any fixed $\delta > 0$ its unique smooth solution u^δ . We multiply Eq. (3.3) by u^δ and we integrate over $\mathbb{R}^d$, obtaining that

$$\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 dx + \frac{1}{2} \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 \operatorname{div} \tilde{\mathbf{b}}^\delta(t, x) dx + \frac{1}{2} \sum_{i,j} \int_{\mathbb{R}^d} a_{ij}^\delta \partial_j u^\delta \partial_i u^\delta = 0. \quad (3.4)$$

Thus, we use the uniform ellipticity of $\mathbf{a}^\delta$ to obtain the inequality

$$\frac{d}{dt} \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 dx + \alpha \int_{\mathbb{R}^d} |\nabla u^\delta(t, x)|^2 dx \leq \|(\operatorname{div} \tilde{\mathbf{b}}^\delta(t, \cdot))\|_\infty \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 dx, \quad (3.5)$$

and by using Gronwall's lemma we obtain that

$$\|u^\delta\|_{L^\infty L^2} + \|u^\delta\|_{L^2 H^1} \leq C(\alpha, \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty}) \|u_0\|_{L^2}. \quad (3.6)$$

A standard compactness argument and the linearity of Eq. (FP-div) guarantee that

$$u^\delta \rightharpoonup^* u \in L^\infty L^2 \cap L^2 H^1,$$

where u is a weak solution to (FP-div). We now move to the uniqueness: assume that u is a weak solution to (FP-div) with $u_0 = 0$ and define $u^\delta := u * \rho^\delta$. Then, u^δ solves the equation

$$\begin{cases} \partial_t u^\delta + \operatorname{div}(\tilde{\mathbf{b}} u^\delta) - \frac{1}{2} \sum_{i,j} \partial_i(a_{ij} \partial_j u^\delta) = r^\delta + s_1^\delta, \\ u^\delta(0, \cdot) = 0, \end{cases} \quad (3.7)$$

where r^δ and s_1^δ are the commutators defined in (2.2) and (2.9). Consider a smooth function φ such that

$$\varphi \geq 0, \quad \operatorname{supp} \varphi \subset B_2, \quad \varphi = 1 \text{ in } B_1,$$

and define $\varphi_R(x) = \varphi(x/R)$. Let $\beta_M : \mathbb{R} \rightarrow \mathbb{R}$ be an even function with the following properties:

- $\beta_M \in C^2 \cap L^\infty(\mathbb{R})$ with $\|\beta_M'\|_\infty < CM$ and $\|\beta_M''\|_\infty < C$ for some positive constant C independent on M ;
- $\beta_M(z) = z^2$ for $|z| \leq M$ for some positive M ;
- $0 \leq \beta_M(z) \leq 2M^2$ for every $z \in \mathbb{R}$;
- $\beta_M(z) \leq z^2$ for every $z \in \mathbb{R}$;
- $\beta_M(z) = 0 \iff z = 0$;
- $|\beta_M'(z)| \leq C|z|$.

We multiply Eq. (3.7) by $\beta_M'(u^\delta) \varphi_R$ and integrating on $(0, t) \times \mathbb{R}^d$ we obtain the following: for the term involving the vector field we have that

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d} \operatorname{div}(\tilde{\mathbf{b}} u^\delta) \beta_M'(u^\delta) \varphi_R dx ds &= - \int_0^t \int_{\mathbb{R}^d} u^\delta \tilde{\mathbf{b}} \cdot \nabla \beta_M'(u^\delta) \varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds \\ &= - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds \\ &= - \iint_{\{|u^\delta| < M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) dx ds \end{aligned}$$

$$- \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds.$$

Now we use that $\beta_M(u^\delta) = |u^\delta|^2$ for $|u^\delta| \leq M$, so

$$\begin{aligned} &= -2 \iint_{\{|u^\delta| < M\}} (u^\delta \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + |u^\delta|^2 \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds - \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds \\ &= - \iint_{\{|u^\delta| < M\}} (\tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R + 2\beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds - \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds \\ &= - \iint_{(0,t) \times \mathbb{R}^d} \tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R \, dx ds - \iint_{\{|u^\delta| < M\}} 2\beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds \\ &\quad - \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R - \tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R) \, dx ds. \end{aligned}$$

Finally, by integrating by parts we get that

$$\begin{aligned} &= \iint_{(0,t) \times \mathbb{R}^d} (\operatorname{div} \tilde{\mathbf{b}} \beta_M(u^\delta) \varphi_R + \beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R) \, dx ds - \iint_{\{|u^\delta| < M\}} 2\beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds \\ &\quad - \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R - \tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R) \, dx ds \\ &= \iint_{(0,t) \times \mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}} \beta_M(u^\delta) \varphi_R \, dx ds + \iint_{\{|u^\delta| > M\}} \beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds - \iint_{\{|u^\delta| < M\}} \beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds \\ &\quad - \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R - \tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R) \, dx ds. \end{aligned}$$

On the other hand, for the term involving the diffusion we have

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d} \partial_i (a_{ij} \partial_j u^\delta) \beta_M'(u^\delta) \varphi_R \, dx ds &= - \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j u^\delta \partial_i u^\delta \beta_M''(u^\delta) \varphi_R \, dx ds - \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j u^\delta \beta_M'(u^\delta) \partial_i \varphi_R \, dx ds \\ &= - \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j u^\delta \partial_i u^\delta \beta_M''(u^\delta) \varphi_R \, dx ds - \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j \beta_M(u^\delta) \partial_i \varphi_R \, dx ds. \end{aligned} \quad (3.8)$$

Then, we use the uniform ellipticity of $\mathbf{a}$, i.e.

$$2\alpha \iint_{\{|u^\delta| < M\}} |\nabla u^\delta|^2 \varphi_R \, dx ds \leq \iint_{\{|u^\delta| < M\}} a_{ij} \partial_i u^\delta \partial_j u^\delta \beta_M''(u^\delta) \varphi_R \, dx ds \quad (3.9)$$

to obtain the following inequality

$$\begin{aligned} &\int_{\mathbb{R}^d} \beta_M(u^\delta) \varphi_R \, dx + \alpha \iint_{\{|u^\delta| < M\}} |\nabla u^\delta|^2 \varphi_R \, dx ds \\ &\leq - \iint_{(0,t) \times \mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}} \beta_M(u^\delta) \varphi_R \, dx ds - \iint_{\{|u^\delta| > M\}} \beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds + \iint_{\{|u^\delta| < M\}} \beta_M(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R \, dx ds \\ &\quad + \iint_{\{|u^\delta| > M\}} (u^\delta \beta_M''(u^\delta) \tilde{\mathbf{b}} \cdot \nabla u^\delta \varphi_R + u^\delta \beta_M'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R - \tilde{\mathbf{b}} \cdot \nabla \beta_M(u^\delta) \varphi_R) \, dx ds \\ &\quad - \frac{1}{2} \iint_{\{|u^\delta| > M\}} a_{ij} \partial_j u^\delta \partial_i u^\delta \beta_M''(u^\delta) \varphi_R \, dx ds - \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j \beta_M(u^\delta) \partial_i \varphi_R \, dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) \beta_M'(u^\delta) \varphi_R \, dx ds. \end{aligned}$$

Then, we exploit the fact that we need to account only for the negative part of the divergence in the term on the right, and using trivial estimates we obtain that

$$\begin{aligned} \int_{\mathbb{R}^d} \beta_M(u^\delta) \varphi_R \, dx &\leq \iint_{(0,t) \times \mathbb{R}^d} (\operatorname{div} \tilde{\mathbf{b}})^- \beta_M(u^\delta) \varphi_R \, dx ds + \iint_{(0,t) \times \mathbb{R}^d} \beta_M(u^\delta) |\tilde{\mathbf{b}}| |\nabla \varphi_R| \, dx ds \\ &\quad + \iint_{\{|u^\delta| > M\}} \left(|u^\delta \tilde{\mathbf{b}}| |\nabla u^\delta| |\beta_M''(u^\delta)| + |\tilde{\mathbf{b}}| |\nabla u^\delta| |\beta_M'(u^\delta)| + \frac{1}{2} |\mathbf{a}| |\nabla u^\delta|^2 |\beta_M''(u^\delta)| \right) \varphi_R \, dx ds \\ &\quad + \iint_{\{|u^\delta| > M\}} |u^\delta \tilde{\mathbf{b}}| |\beta_M'(u^\delta)| |\nabla \varphi_R| \, dx ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} |\mathbf{a}| |\beta_M'(u^\delta)| |\nabla u^\delta| |\nabla \varphi_R| \, dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) \beta_M'(u^\delta) \varphi_R \, dx ds. \end{aligned} \quad (3.10)$$

Let us estimate by Young

$$\begin{aligned} &\iint_{\{|u^\delta| > M\}} \left(|u^\delta \tilde{\mathbf{b}}| |\nabla u^\delta| |\beta_M''(u^\delta)| + |\tilde{\mathbf{b}}| |\nabla u^\delta| |\beta_M'(u^\delta)| + \frac{1}{2} |\mathbf{a}| |\nabla u^\delta|^2 |\beta_M''(u^\delta)| \right) \varphi_R \, dx ds \\ &\leq \|\beta_M''\|_\infty \iint_{\{|u^\delta| > M\}} |u^\delta \tilde{\mathbf{b}}| |\nabla u^\delta| \, dx ds + C \iint_{\{|u^\delta| > M\}} |\tilde{\mathbf{b}}| |u^\delta| |\nabla u^\delta| \, dx ds + \frac{1}{2} \|\mathbf{a}\|_\infty \|\beta_M''\|_\infty \iint_{\{|u^\delta| > M\}} |\nabla u^\delta|^2 \, dx ds \end{aligned}$$

$$\leq C(\|\beta_M''\|_\infty, \|a\|_\infty) \iint_{\{|u^\delta| > M\}} (|u^\delta \tilde{b}|^2 + |\nabla u^\delta|^2) dx ds.$$

By Chebishev's inequality we can estimate the super-level as follows

$$\mathcal{L}^{d+1}(\{(t, x) \in [0, T] \times \mathbb{R}^d : |u^\delta(t, x)| > M\}) \leq \frac{\|u^\delta\|_{L^2_{t,x}}^2}{M^2} \leq \frac{C(T, \|u_0\|_{L^2})}{M^2}, \quad (3.11)$$

which is small in M uniformly in δ . Then, since the quantity $(|u^\delta \tilde{b}|^2 + |\nabla u^\delta|^2) \in L^1_{t,x}$ uniformly in δ by the assumptions, we use the equi-integrability of the integrand to obtain the following estimate

$$\iint_{\{|u^\delta| > M\}} \left(|u^\delta \tilde{b}| |\nabla u^\delta| |\beta_M''(u^\delta)| + |\tilde{b}| |\nabla u^\delta| |\beta_M'(u^\delta)| + \frac{1}{2} |a| |\nabla u^\delta|^2 |\beta_M''(u^\delta)| \right) \varphi_R dx ds \leq F(M), \quad (3.12)$$

for some function F , depending on $\|\beta_M''\|_\infty, \|a\|_\infty, \|u_0\|_{L^2}, T$, such that $F(M) \rightarrow 0$ as $M \rightarrow \infty$. On the other hand, for the third line in (3.10) we proceed as follows: for the term involving the vector field we use Hölder inequality and (3.11) to obtain the estimate

$$\iint_{\{|u^\delta| > M\}} |u^\delta \tilde{b}| |\beta_M'(u^\delta)| |\nabla \varphi_R| dx ds \leq \frac{\|\beta_M'\|_\infty}{R} \|u^\delta \tilde{b}\|_{L^2} \frac{C(T, \|u_0\|_{L^2})}{M} := \frac{C}{R}.$$

For the term involving the diffusion we use the definition of β_M to obtain

$$\int_0^t \int_{\mathbb{R}^d} |a| |\beta_M'(u^\delta)| |\nabla u^\delta| |\nabla \varphi_R| dx ds \leq \frac{C \|a\|_\infty}{R} \int_0^t \int_{\mathbb{R}^d} |\nabla u^\delta| |u^\delta| dx ds \leq \frac{C \|a\|_\infty}{R} \|\nabla u\|_{L^2 L^2} \|u\|_{L^\infty L^2} \leq \frac{C}{R},$$

where in the penultimate inequality we used Hölder's inequality, and the constant C depends on $\|a\|_\infty, \|u_0\|_{L^2}, T$. Summarizing, we rewrite (3.10) as follows

$$\begin{aligned} \int_{\mathbb{R}^d} \beta_M(u^\delta) \varphi_R dx + \alpha \iint_{\{|u^\delta| < M\}} |\nabla u^\delta|^2 \varphi_R dx ds &\leq \iint_{(0,t) \times \mathbb{R}^d} (\operatorname{div} \tilde{b})^- \beta_M(u^\delta) \varphi_R dx ds + \iint_{(0,t) \times \mathbb{R}^d} \beta_M(u^\delta) |\tilde{b}| |\nabla \varphi_R| dx ds \\ &\quad + F(M) + \frac{C}{R} + \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) \beta_M'(u^\delta) \varphi_R dx ds. \end{aligned}$$

Since $\beta_M'(u^\delta) \varphi_R \rightarrow \beta_M'(u) \varphi_R$ strongly in $L^2((0, T); H_0^1(B_{2R}))$, we can use Lemmas 2.4 and 2.7 and, by letting $\delta \rightarrow 0$, we get that

$$\begin{aligned} \int_{\mathbb{R}^d} \beta_M(u) \varphi_R dx &\leq \iint_{(0,t) \times \mathbb{R}^d} (\operatorname{div} \tilde{b})^- \beta_M(u) \varphi_R dx ds + \iint_{(0,t) \times \mathbb{R}^d} \beta_M(u) |\tilde{b}| |\nabla \varphi_R| dx ds + \frac{C}{R} \\ &\leq \int_0^t \|(\operatorname{div} \tilde{b}(s, \cdot))^- \|_\infty \int_{\mathbb{R}^d} \beta_M(u) \varphi_R dx ds + 2M^2 \int_0^T \int_{|x| > R} |\tilde{b}_1(s, x)| dx ds \\ &\quad + \int_0^T \|\tilde{b}_2(s, \cdot)\|_\infty \int_{|x| > R} |u(s, x)|^2 dx ds + F(M) + \frac{C}{R} \\ &:= \int_0^t \|(\operatorname{div} \tilde{b}(s, \cdot))^- \|_\infty \int_{\mathbb{R}^d} \beta_M(u) \varphi_R dx ds + H(R, M) + F(M), \end{aligned}$$

where $\tilde{b}_1$ and $\tilde{b}_2$ are as in (3.2), and the function $H(R, M)$ goes to 0 as $R \rightarrow \infty$ for any fixed $M > 0$. Notice that in the last inequality we used the definition of φ_R , the growth assumption on $\tilde{b}$, and the properties of β_M . Finally, we apply Gronwall's inequality

$$\int_{\mathbb{R}^d} \beta_M(|u(t, x)|) \varphi_R dx \leq [H(R, M) + F(M)] \exp(\|(\operatorname{div} \tilde{b})^- \|_{L^1 L^\infty}),$$

and by letting $R \rightarrow \infty$, we obtain that

$$\int_{\mathbb{R}^d} \beta_M(|u(t, x)|) dx \leq F(M) \exp(\|(\operatorname{div} \tilde{b})^- \|_{L^1 L^\infty}), \quad (3.13)$$

for any $M > 0$. By Lebesgue's dominated convergence Theorem

$$\int_{\mathbb{R}^d} \beta_M(|u(t, x)|) dx \rightarrow \int_{\mathbb{R}^d} |u(t, x)|^2 dx,$$

as $M \rightarrow \infty$, and together with (3.13), this implies that $u(t, x) = 0$ for a.e. $(t, x) \in (0, T) \times \mathbb{R}^d$. $\square$

Remark 3.3. An example of a function β_M satisfying the assumptions of Theorem 3.2 is given by a regularized version of the following function:

$$\beta_M(z) := \begin{cases} z^2 & \text{if } |z| \leq M, \\ -\frac{z^2}{2} + 2Mz - \frac{M^2}{2} & \text{if } M < |z| < 2M, \\ \frac{7}{2}M^2 & \text{if } |z| > 2M. \end{cases} \quad (3.14)$$

3.1.1. Comparison with [18]

We now compare the uniqueness result of Theorem 3.2 with Proposition 4 of [18], which we report here for completeness.

Proposition 3.4. Assume that the matrix $\mathbf{a} := \sigma\sigma^T$ is uniformly positive definite and that

$$\begin{cases} \tilde{\mathbf{b}} \in L^2((0, T); L^2_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d)), & \frac{\tilde{\mathbf{b}}}{1+|x|} \in L^1((0, T); L^1 + L^\infty(\mathbb{R}^d; \mathbb{R}^d)), \\ \operatorname{div} \tilde{\mathbf{b}} \in L^1((0, T); L^\infty(\mathbb{R}^d)), \\ \sigma \in L^\infty((0, T); L^\infty_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^{d \times d})), & \frac{\sigma}{1+|x|} \in L^2((0, T); L^2 + L^\infty(\mathbb{R}^d; \mathbb{R}^{d \times d})). \end{cases} \quad (3.15)$$

Then, for each initial condition in $L^1 \cap L^\infty(\mathbb{R}^d)$, Eq. (FP-div) has a unique solution in the space

$$\{u \in L^\infty((0, T); L^1 \cap L^\infty(\mathbb{R}^d)), u \in L^2((0, T); H^1(\mathbb{R}^d)), \sigma^T \nabla u \in L^2((0, T); L^2(\mathbb{R}^d))\}. \quad (3.16)$$

Note that if we consider the assumptions of Theorem 3.2 with $p = 2, q = \infty$ LeBris-Lions' result is not immediately recovered because of local integrability assumptions. However, by making the following changes to the proof of Theorem 3.2, we can prove a slightly different result.

Corollary 3.5. Assume that the matrix $\mathbf{a} := \sigma\sigma^T$ is uniformly positive definite and that

$$\begin{cases} \tilde{\mathbf{b}} \in L^2((0, T); L^2(\mathbb{R}^d; \mathbb{R}^d)), & \frac{\tilde{\mathbf{b}}}{1+|x|} \in L^1((0, T); L^1 + L^\infty(\mathbb{R}^d; \mathbb{R}^d)), \\ (\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d)), \\ \sigma \in L^\infty((0, T); L^\infty_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^{d \times d})), & \frac{\sigma}{1+|x|} \in L^2((0, T); L^2 + L^\infty(\mathbb{R}^d; \mathbb{R}^{d \times d})). \end{cases} \quad (3.17)$$

Then, for each initial condition in $L^1 \cap L^\infty(\mathbb{R}^d)$, Eq. (FP-div) has a unique solution in the space

$$\{u \in L^\infty((0, T); L^1 \cap L^\infty(\mathbb{R}^d)), u \in L^2((0, T); H^1(\mathbb{R}^d)), \sigma^T \nabla u \in L^2((0, T); L^2(\mathbb{R}^d))\}. \quad (3.18)$$

Proof. Since the argument is very similar to that of Theorem 3.2 we only sketch the proof. First, since we are dealing with bounded solutions, we can consider the function $\beta(z) = |z|^2$ and all the integrals on the set $\{|u^\delta| > M\}$ are identically zero for M large enough. This implies that we have to use a Gronwall type argument on the following estimate

$$\begin{aligned} \int_{\mathbb{R}^d} \beta(u^\delta) \varphi_R dx + \alpha \iint_{\{|u^\delta| < M\}} |\nabla u^\delta|^2 \varphi_R dx ds &\leq - \iint_{(0,t) \times \mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}} \beta(u^\delta) \varphi_R dx ds + \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) \beta'(u^\delta) \varphi_R dx ds \\ &\quad + \iint_{(0,t) \times \mathbb{R}^d} \beta(u^\delta) |\tilde{\mathbf{b}}| |\nabla \varphi_R| dx ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j \beta(u^\delta) \partial_i \varphi_R dx ds. \end{aligned} \quad (3.19)$$

The local integrability assumptions together with the bound on $(\operatorname{div} \tilde{\mathbf{b}})^-$ are sufficient for the convergence of the commutators. On the other hand, we use that the matrix $\mathbf{a}$ is representable as $a_{ij} = \sigma_{ik} \sigma_{jk}$, to estimate the last term in (3.19) as follows

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d} a_{ij} \partial_j \beta(u^\delta) \partial_i \varphi_R dx ds &= \int_0^t \int_{\mathbb{R}^d} \sigma_{ik} \sigma_{jk} \partial_j u^\delta \beta'(u^\delta) \partial_i \varphi_R dx ds \\ &\leq \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} |\sigma_{jk} \partial_j u^\delta|^2 |\partial_i \varphi_R| dx ds + \frac{1}{2} \int_0^t \int_{\mathbb{R}^d} |\sigma_{ik}|^2 |\beta'(u^\delta)|^2 |\partial_i \varphi_R| dx ds \\ &\leq \frac{C}{R} \int_0^t \int_{\mathbb{R}^d} |\sigma_{jk} \partial_j u^\delta|^2 dx ds + C \|u_0\|_\infty^2 \int_0^t \int_{|x| > R} |\sigma_{ik}|^2 dx ds + \int_0^t \|\sigma_{ik}^2(s, \cdot)\|_\infty \int_{|x| > R} |u^\delta|^2 dx ds, \end{aligned}$$

which goes to 0 as $R \rightarrow \infty$. The conclusion follows along the same lines as in the proof of Theorem 3.2. $\square$

Notice that our corollary works under assumptions of global integrability on the vector field $\tilde{\mathbf{b}}$, but we only assume that the negative part of the divergence is bounded. Remarkably, we can use this weaker assumption thanks to the convergence of the commutators in $L^2_t H_x^{-1}$. Assuming a bound on the full divergence we would then be able to consider the case of locally integrable vector fields, as done by LeBris-Lions, relying on the convergence in L^1 of the commutator r^δ (see Lemma 2.1) and finding the very same proof.

3.2. The Fokker–Planck equation

We start by giving the following definition.

Definition 3.6 (Distributional Solution). Assume that

$$\mathbf{b} \in L^1((0, T); L^p(\mathbb{R}^d)), \quad \mathbf{a} \in L^1((0, T); L^p(\mathbb{R}^d; \mathbb{R}^{d \times d})), \quad u_0 \in L^q(\mathbb{R}^d) \quad (3.20)$$

are given, with p, q satisfying $1/p + 1/q \leq 1$. A function $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ is a *distributional solution* to (FP) if for any $\varphi \in C_c^\infty([0, T) \times \mathbb{R}^d)$ the following identity holds:

$$\int_0^T \int_{\mathbb{R}^d} u \left(\partial_t \varphi + \mathbf{b} \cdot \nabla \varphi + \frac{1}{2} \sum_{ij} a_{ij} \partial_{ij} \varphi \right) dx dt + \int_{\mathbb{R}^d} u_0 \varphi(0, \cdot) dx = 0.$$

Notice that in the definition of distributional solutions the assumption that p, q satisfy $1/p + 1/q \leq 1$ is the minimum requirement we need in order to have $ub, ua \in L^1$ so that the definition makes sense.

We now recall the following definition.

Definition 3.7. Let Ω be an open subset of $\mathbb{R}^d$. We say that a bounded family $\{\varphi_i\}_{i \in I}$ is equi-integrable if the following two conditions hold:

(i) For any $\varepsilon > 0$ there exists a borel set $A \subset \Omega$ with finite measure such that

$$\int_{\Omega \setminus A} |\varphi_i(x)| dx \leq \varepsilon,$$

for any $i \in I$.

(ii) For any $\varepsilon > 0$ there exists $\delta > 0$ such that, for any Borel set $E \subset \Omega$ with $\mathcal{L}^d(E) \leq \delta$, there holds

$$\int_E |\varphi_i(x)| dx \leq \varepsilon,$$

for any $i \in I$.

Before to prove the existence of distributional solutions we recall the following characterization of equi-integrability, see [4] (or alternatively [2] for a proof on the d -dimensional torus $\mathbb{T}^d$).

Lemma 3.8. Consider a family $\{\varphi_i\}_{i \in I} \subset L^1(\Omega)$ which is bounded in $L^1(\Omega)$. Then this family satisfies condition (ii) of Definition 3.7 if and only if for every $\varepsilon > 0$, there exists a constant $C_\varepsilon > 0$ such that for every $i \in I$ we can write

$$\varphi_i = \varphi_i^1 + \varphi_i^2,$$

with

$$\|\varphi_i^1\|_{L^1(\Omega)} \leq \varepsilon, \quad \|\varphi_i^2\|_{L^2(\Omega)} \leq C_\varepsilon, \text{ for all } i \in I. \quad (3.21)$$

The proof of existence of distributional solutions immediately follows from a classical a priori estimate.

Proposition 3.9 (Existence of Distributional Solutions). Let b, a and u_0 be as in (3.20) with p, q such that $1/p + 1/q \leq 1$ and let (A2) and (A4) hold and assume that $(\operatorname{div} b)^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Then there exists a distributional solution $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ to (FP).

Proof. Let $(\rho^\delta)_\delta$ be a standard family of mollifiers as in (2.1) and let us define $b^\delta := b * \rho^\delta$, $a^\delta := a * \rho^\delta$, and $u_0^\delta = u_0 * \rho^\delta$. Then, we consider the approximating problem

$$\begin{cases} \partial_t u^\delta + \operatorname{div}(b^\delta u^\delta) - \frac{1}{2} \sum_{i,j} \partial_{ij}(a_{ij}^\delta u^\delta) = 0, \\ u^\delta(0, \cdot) = u_0^\delta. \end{cases} \quad (3.22)$$

The existence of a unique solution of (3.22) follows from classical theory [11]. It is readily checked that the sequence u^δ is equi-bounded in $L^\infty((0, T); L^q(\mathbb{T}^d))$. Indeed, from the smoothness of the objects in (3.22), we can rewrite it in divergence form

$$\partial_t u^\delta + \operatorname{div}(\tilde{b}^\delta u^\delta) - \frac{1}{2} \sum_{i,j} \partial_{ij}(a_{ij}^\delta \partial_j u^\delta) = 0, \quad (3.23)$$

and then we can multiply the equation by $\beta'(u^\delta)$, where $\beta: \mathbb{R} \rightarrow \mathbb{R}$ is a smooth, convex function: by an easy application of the chain rule and by integrating in space we obtain (for fixed t)

$$\begin{aligned} \int_{\mathbb{R}^d} \operatorname{div}(\tilde{b}^\delta u^\delta) \beta'(u^\delta) dx &= \int_{\mathbb{R}^d} u^\delta \beta'(u^\delta) \operatorname{div} \tilde{b}^\delta dx + \int_{\mathbb{R}^d} \tilde{b}^\delta \cdot \nabla \beta(u^\delta) dx \\ &= \int_{\mathbb{R}^d} u^\delta \beta'(u^\delta) \operatorname{div} \tilde{b}^\delta dx + \int_{\mathbb{R}^d} \operatorname{div}(\tilde{b}^\delta \beta(u^\delta)) dx - \int_{\mathbb{R}^d} \operatorname{div} \tilde{b}^\delta \beta(u^\delta) dx, \end{aligned}$$

and

$$\int_{\mathbb{R}^d} \partial_{ij}(a_{ij}^\delta \partial_j u^\delta) \beta'(u^\delta) dx = - \int_{\mathbb{R}^d} a_{ij}^\delta \beta''(u^\delta) \partial_i u^\delta \partial_j u^\delta dx.$$

By considering a sequence of smooth, convex functions, uniformly convergent to $\beta(s) = |s|^q$, for $1 < q < \infty$, the following uniform bounds on the L^q -norm of the solutions u^δ :

$$\frac{d}{dt} \|u^\delta(t, \cdot)\|_{L^q}^q \leq -(q-1) \int_{\mathbb{R}^d} \operatorname{div} \tilde{b}^\delta(t, x) |u^\delta(t, x)|^q dx - \alpha \int_{\mathbb{R}^d} |\nabla u^\delta(t, x)|^2 \beta''(u^\delta(t, x)) dx \quad (3.24)$$

$$\leq (q-1) \|(\operatorname{div} \tilde{b}^\delta)^-(t, \cdot)\|_\infty \|u^\delta(t, \cdot)\|_{L^q}^q, \quad (3.25)$$

where in (3.24) we have used that, by standard properties of mollifiers, a^δ is uniformly elliptic with the same constant α as in (A4). Notice that $\|(\operatorname{div} \tilde{b}^\delta)^-\|_{L^1 L^\infty}$ is uniformly bounded (in δ) by (A2). The conclusion now follows from an application of Gronwall's

lemma: we have that

$$\sup_{t \in [0, T]} \|u^\delta(t, \cdot)\|_{L^q(\mathbb{R}^d)} \leq \|u_0\|_{L^q(\mathbb{R}^d)} e^{C(q)\|(\operatorname{div} \tilde{b})^-\|_{L^1 L^\infty}}, \quad (3.26)$$

with $C(q) = \frac{q-1}{q}$. Thus, for $q > 1$ by standard compactness arguments, we can extract a subsequence which converges weakly-* to a function $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ and it is immediate to deduce that u is a distributional solution of (FP) because of the linearity of the equation. For $q = \infty$, the estimate (3.26) still holds for every $\delta > 0$: we send $q \rightarrow \infty$ in (3.26) and then we can conclude as in the previous case. For the case $q = 1$, the estimate is not sufficient to obtain weak compactness in L^1 , as we need to show the equi-integrability of the family $(u^\delta)_{\delta > 0}$. We first discuss the equi-integrability on small sets: let $u_0 \in L^1(\mathbb{R}^d)$ and define u_0^δ as above. Since $u_0^\delta \rightarrow u_0$ strongly in $L^1(\mathbb{R}^d)$, it converges weakly and thus it is equi-integrable. Then, we apply Lemma 3.8 to the family u_0^δ : let $\varepsilon > 0$ be fixed, there exists a constant $C_\varepsilon > 0$ such that

$$u_0^\delta = u_{0,1}^\delta + u_{0,2}^\delta,$$

with

$$\|u_{0,1}^\delta\|_{L^1} \leq \varepsilon, \quad \|u_{0,2}^\delta\|_{L^2} \leq C_\varepsilon.$$

We define u_1^δ and u_2^δ to be the solutions of (3.22) with initial datum, respectively, $u_{0,1}^\delta$ and $u_{0,2}^\delta$. Thus, the estimate in (3.26) with $q = 1, 2$ implies that

$$\sup_{t \in [0, T]} \|u_1^\delta(t, \cdot)\|_{L^1(\mathbb{R}^d)} \leq \varepsilon e^{C\|(\operatorname{div} \tilde{b})^-\|_{L^1 L^\infty}}, \quad (3.27)$$

$$\sup_{t \in [0, T]} \|u_2^\delta(t, \cdot)\|_{L^2(\mathbb{R}^d)} \leq C_\varepsilon e^{C\|(\operatorname{div} \tilde{b})^-\|_{L^1 L^\infty}}. \quad (3.28)$$

The linearity of the equations implies that the function $u^\delta = u_1^\delta + u_2^\delta$ is the unique solution of (3.22) with initial datum u_0^δ and it admits a decomposition as in (3.21). Then, we can infer that the sequence u^δ satisfies (ii) of Definition 3.7.

We now verify the condition on the tails, i.e. (i) in Definition 3.7: let $r, R > 0$ such that $2r < R$ and consider a positive test function ψ_r^R such that

$$\begin{cases} 0 & \text{if } 0 < |x| < r, \\ 1 & \text{if } 2r < |x| < R, \\ 0 & \text{if } |x| > R, \end{cases} \quad (3.29)$$

such that $0 \leq \psi_r^R \leq 1$ and

$$|\nabla \psi_r^R| \leq \frac{C}{r}, \quad |\nabla^2 \psi_r^R| \leq \frac{C}{r^2}. \quad (3.30)$$

Then consider a smooth convex function β_ε converging uniformly to $\beta(s) = |s|$ (for example we can take $\beta_\varepsilon(s) = \sqrt{s^2 + \varepsilon^2} - \varepsilon$) and multiply Eq. (3.23) by $\beta'_\varepsilon(|u^\delta(t, x)|)\psi_r^R(x)$: integrating by parts we get that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^d} \beta_\varepsilon(|u^\delta(t, x)|) \psi_r^R(x) dx &+ \int_{\mathbb{R}^d} \operatorname{div} \tilde{b}^\delta(t, x) (u^\delta(t, x) \beta'_\varepsilon(|u^\delta(t, x)|) - \beta_\varepsilon(|u^\delta(t, x)|)) \psi_r^R(x) dx - \int_{\mathbb{R}^d} \beta_\varepsilon(u^\delta(t, x)) \tilde{b}(t, x) \cdot \nabla \psi_r^R(x) dx \\ &+ \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} a_{ij}(t, x) \partial_j u^\delta(t, x) \partial_i u^\delta(t, x) \beta''_\varepsilon(|u^\delta(t, x)|) \psi_r^R(x) dx + \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} a_{ij}(t, x) \partial_j \beta_\varepsilon(|u^\delta(t, x)|) \partial_i \psi_r^R(x) dx = 0. \end{aligned}$$

Then we estimate the fourth term above using the convexity of β_ε together with (A4), while we do a further integration by parts in the last term to obtain that

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^d} \beta_\varepsilon(|u^\delta(t, x)|) \psi_r^R(x) dx &\leq \int_{\mathbb{R}^d} \operatorname{div} \tilde{b}^\delta(t, x) (u^\delta(t, x) \beta'_\varepsilon(|u^\delta(t, x)|) - \beta_\varepsilon(|u^\delta(t, x)|)) \psi_r^R(x) dx + \int_{\mathbb{R}^d} |\beta_\varepsilon(u^\delta(t, x))| \tilde{b}(t, x) \cdot \nabla \psi_r^R(x) dx \\ &+ \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} |\partial_j a_{ij}(t, x)| \beta_\varepsilon(|u^\delta(t, x)|) |\partial_i \psi_r^R(x)| dx + \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} |a_{ij}(t, x)| \beta_\varepsilon(|u^\delta(t, x)|) |\partial_{ij} \psi_r^R(x)| dx. \end{aligned}$$

Let $t \in (0, T)$ be fixed, we integrate on $(0, t)$ and by letting $\varepsilon \rightarrow 0$ we have that

$$u^\delta(t, x) \beta'_\varepsilon(|u^\delta(t, x)|) - \beta_\varepsilon(|u^\delta(t, x)|) \rightarrow 0,$$

leading to the inequality

$$\begin{aligned} \int_{\mathbb{R}^d} |u^\delta(t, x)| \psi_r^R(x) dx &\leq \int_{\mathbb{R}^d} |u_0^\delta(x)| \psi_r^R(x) dx + \int_{\mathbb{R}^d} |u^\delta(t, x)| \tilde{b}(t, x) \cdot \nabla \psi_r^R(x) dx + \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} |\partial_j a_{ij}(t, x)| |u^\delta(t, x)| |\partial_i \psi_r^R(x)| dx \\ &+ \frac{1}{2} \sum_{ij} \int_{\mathbb{R}^d} |a_{ij}(t, x)| |u^\delta(t, x)| |\partial_{ij} \psi_r^R(x)| dx. \end{aligned}$$

Finally, we use the definition of ψ_r^R and by letting $R \rightarrow \infty$ we obtain

$$\int_{|x| > r} |u^\delta(t, x)| dx \leq \int_{|x| > r} |u_0^\delta(x)| dx + C \left(\frac{\|\tilde{b}\|_\infty + \sum_{ij} \|\partial_j a_{ij}\|_\infty}{r} \right) \int_{|x| > r} |u^\delta(t, x)| dx + C \frac{\sum_{ij} \|a_{ij}\|_\infty}{r^2} \int_{|x| > r} |u^\delta(t, x)| dx.$$

Note that the constant C above does not depend on δ and ι . The conclusion follows by using the equi-integrability of u_0^δ and the uniform bound on $\|u^\delta\|_{L^\infty L^1}$. So far we have shown that the sequence $\{u^\delta\}_{\delta>0} \subset L^1((0, T) \times \mathbb{R}^d)$ is uniformly integrable. This implies that there exists a function $u \in L^1((0, T) \times \mathbb{R}^d)$ such that, up to a subsequence, the following convergence holds

$$u^\delta \rightharpoonup u \quad \text{in } L^1((0, T) \times \mathbb{R}^d).$$

In order to conclude we need to check that $u^\delta \xrightarrow{*} u$ in $L^\infty L^1$. Note that the sequence u^δ satisfies the inequality (3.26) with $q = 1$, which implies that

$$u^\delta \xrightarrow{*} u \quad \text{in } L^\infty((0, T); \mathcal{M}(\mathbb{R}^d)).$$

Then, for any $\psi \in C_c^\infty([0, T])$ and $\phi \in C_0(\mathbb{R}^d)$ we have that

$$\int_0^T \psi(t) \int_{\mathbb{R}^d} \phi(x) du_t(x) dt = \int_0^T \psi(t) \int_{\mathbb{R}^d} \phi(x) u(t, x) dx dt. \quad (3.31)$$

We can conclude that for a.e. $t \in [0, T]$ we have that

$$\int_{\mathbb{R}^d} \phi(x) du_t(x) = \int_{\mathbb{R}^d} \phi(x) u(t, x) dx, \quad (3.32)$$

which implies that u_t is absolutely continuous with respect to the Lebesgue measure and

$$\|u_t\|_{\mathcal{M}} = \|u(t, \cdot)\|_{L^1}, \quad \text{for a.e. } t \in (0, T).$$

Note that is crucial that the space $C_0(\mathbb{R}^d)$ is separable, because the set of zero measure where (3.32) does not hold may depend on ϕ . Finally, we can infer that $u \in L^\infty((0, T); L^1(\mathbb{R}^d))$ and therefore together with the weak convergence in $L^1((0, T) \times \mathbb{R}^d)$ we can conclude that

$$u^\delta \xrightarrow{*} u, \quad \text{in } L^\infty((0, T); L^1(\mathbb{R}^d)),$$

and this concludes the proof. $\square$

A special sub-class of distributional solutions is given by *parabolic solutions*, which are weakly differentiable in the space variable. This notion of solution is natural for vector fields and diffusion matrices possessing enough integrability in the space variable.

Definition 3.10. Assume that

$$b \in L^1((0, T); L^2(\mathbb{R}^d; \mathbb{R}^d)), \quad a \in L^2((0, T); L^2(\mathbb{R}^d; \mathbb{R}^{d \times d})), \quad u_0 \in L^2(\mathbb{R}^d). \quad (3.33)$$

A function $u \in L^\infty((0, T); L^2(\mathbb{R}^d))$ is a *parabolic solution* to (FP) if it is a distributional solution to (FP) and furthermore $u \in L^2((0, T); H^1(\mathbb{R}^d))$.

We refer to the space $L^2((0, T); H^1(\mathbb{R}^d))$ as the *parabolic class*. We recall that the Fokker–Planck equation (FP) can be formally rewritten in divergence-form as

$$\begin{cases} \partial_t u + \operatorname{div}(\tilde{b}u) - \frac{1}{2} \sum_{i,j} \partial_i(a_{ij} \partial_j u) = 0 & \text{in } (0, T) \times \mathbb{R}^d, \\ u|_{t=0} = u_0 & \text{in } \mathbb{R}^d, \end{cases} \quad (\text{FP-div})$$

by defining the vector field $\tilde{b} := b - \frac{1}{2} \sum_j \partial_j a_{ij}$. The following result holds.

Proposition 3.11. Assume that a satisfies (A2). Then, u is a parabolic solution of (FP) if and only if u is a parabolic solution of (FP-div) with vector field $\tilde{b}$.

Proof. The proof is rather straightforward and relies on a simple integration by parts which makes use of the regularity assumption $u \in L^2((0, T); H^1(\mathbb{R}^d))$. In fact one can show that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} \left(u \tilde{b} \cdot \nabla \varphi + \frac{1}{2} \sum_{i,j} a_{ij} u \partial_{ij} \varphi \right) dx dt &= \int_0^T \int_{\mathbb{R}^d} \left(u \tilde{b} \cdot \nabla \varphi - \frac{1}{2} \sum_{i,j} (\partial_j a_{ij} u + a_{ij} \partial_j u) \partial_i \varphi \right) dx dt \\ &= \int_0^T \int_{\mathbb{R}^d} \left(u \tilde{b} \cdot \nabla \varphi - \frac{1}{2} \sum_{i,j} a_{ij} \partial_j u \partial_i \varphi \right) dx dt, \end{aligned}$$

where in the last line we used the definition of $\tilde{b}$, i.e. $\tilde{b} = b - \frac{1}{2} \sum_j \partial_j a_{ij}$. $\square$

In some situations it is convenient to use the formulation (FP-div) since, roughly speaking, we have to take into account one derivative less on a_{ij} . We will return on this point below. Under our assumptions, there exists at least one solution in the parabolic class: the result is the following.

Proposition 3.12 (Existence of Parabolic Solutions). Let a, b and u_0 be as in (3.33), let (A2), (A4) hold and assume that $(\operatorname{div} \tilde{b})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Then there exists at least one parabolic solution to (FP).

Proof. The proof follows from the same regularization procedure developed in [Proposition 3.9](#): we consider the unique solution u^δ of the regularized problem (3.22). Then, we define the function $\beta(s) = s^2/2$ and integrating in time the inequality (3.24) we get

$$\|u^\delta(t, \cdot)\|_{L^2}^2 + \alpha \int_0^t \int_{\mathbb{R}^d} |\nabla u^\delta(s, x)|^2 dx ds \leq \|u_0^\delta\|_{L^2}^2 - \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}}^\delta(s, x) |u^\delta(s, x)|^2 dx ds. \quad (3.34)$$

From [Proposition 3.9](#) we know that

$$\|u^\delta(t, \cdot)\|_{L^2}^2 \leq \|u_0^\delta\|_{L^2}^2 e^{\int_0^T \|(\operatorname{div} \tilde{\mathbf{b}})^\delta(s, \cdot)\|_\infty ds},$$

and together with (3.34) we obtain that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} |\nabla u^\delta(t, x)|^2 dx dt &\leq \frac{1}{\alpha} \left(\|u_0\|_{L^2}^2 + \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty} \|u^\delta\|_{L^\infty L^2}^2 \right) \\ &\leq \frac{1}{\alpha} \left(\|u_0\|_{L^2}^2 + \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty} \|u_0\|_{L^2}^2 e^{\int_0^T \|(\operatorname{div} \tilde{\mathbf{b}})^\delta(s, \cdot)\|_\infty ds} \right). \end{aligned} \quad (3.35)$$

Thus, we have obtained that

$$\|u^\delta\|_{L^\infty L^2} + \|\nabla u^\delta\|_{L^2 L^2} \leq C(T, \alpha, \|u_0\|_{L^2}, \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty}),$$

and, since the equation is linear, the conclusion follows from a standard compactness argument. $\square$

We can now state the uniqueness theorem for solutions in the parabolic class. The proof easily follows from [Theorem 3.2](#) and [Proposition 3.11](#), thus we omit the details.

Theorem 3.13 (Uniqueness of Parabolic Solutions). Let $u_0 \in L^2 \cap L^q(\mathbb{R}^d)$ be a given initial datum and assume that

(i) $\mathbf{b} \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ with $\frac{1}{p} + \frac{1}{q} = \frac{1}{2}$;

(ii) $\tilde{\mathbf{b}}$ satisfies the growth condition

$$\frac{\tilde{\mathbf{b}}}{1 + |\mathbf{x}|} \in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d));$$

(iii) $\mathbf{a}$ satisfies (A1), (A2) and (A4);

(iv) $(\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$.

Then, there exists a unique parabolic solution u to (FP).

Remark 3.14. Note that if $p = \infty$ and $q = 2$ we recover the existence and uniqueness of parabolic solutions to (FP) in [12, Theorem 4.3].

3.3. Further comments and remarks

In this section we collect some observations concerning the uniqueness theorem, taking into account the different forms of the equation that one can consider. We point out that in [18] there is a nice comparison of the different form of the equations, showing how commutator estimates allow to transfer well-posed results from one equation to another.

- If we consider the Fokker–Planck equation in transport form, i.e.

$$\partial_t u + \mathbf{b} \cdot \nabla u - \frac{1}{2} \sum_{ij} \partial_i (a_{ij} \partial_j u) = 0, \quad (3.36)$$

we can drop the assumption on the $L^p - L^q$ integrability of $\mathbf{b}$ and u in [Theorem 3.2](#). This means that we can consider $\mathbf{b} \in L^2((0, T); L^2(\mathbb{R}^d; \mathbb{R}^d))$ and $u_0 \in L^2(\mathbb{R}^d)$, then we use the convergence of the commutator r_i^δ in [Lemma 2.1](#) to prove the uniqueness of parabolic solutions. The same observation holds true if instead we consider the Fokker–Planck equation (FP) and we assume that $\operatorname{div} \mathbf{b} \in L^1 L^\infty$.

- We can relax the assumption on $\sum_{j=1}^d \partial_j a_{ij} \in L^\infty((0, T) \times \mathbb{R}^d; \mathbb{R}^d)$ in [Propositions 3.9](#) and [3.12](#). Note that in [12] the author considers the case of (FP) with a bounded vector field $\mathbf{b}$, thus he needs (A2) to use the formulation in divergence form with a bounded vector field $\tilde{\mathbf{b}}$. Indeed, the results in Section 3 hold true if one consider

$$\tilde{\mathbf{b}} := \mathbf{b} - \frac{1}{2} \sum_j \partial_j a_{ij} \in L^2((0, T); L^2 \cap L^p(\mathbb{R}^d; \mathbb{R}^d)),$$

and the growth condition

$$\tilde{\mathbf{b}} \in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d)).$$

- Another form of the equation that could be considered is the following

$$\partial_t u + \operatorname{div}(\mathbf{b}u) - \frac{1}{2} \sum_{ij} a_{ij} \partial_{ij} u = 0. \quad (3.37)$$

Roughly speaking, if we test the equation with a test function $\varphi \in C_c^\infty([0, T] \times \mathbb{R}^d)$ and we integrate by parts we get that

$$\int_0^T \int_{\mathbb{R}^d} u(\partial_t \varphi + \mathbf{b} \cdot \nabla \varphi) - \frac{1}{2} \sum_{ij} \partial_j(a_{ij} \varphi) \partial_i u \, dx dt + \int_{\mathbb{R}^d} u_0 \varphi(0, \cdot) dx = 0. \quad (3.38)$$

Thus, we can define weak solutions as in [Definition 3.1](#) requiring $\mathbf{a}, \sum_j \partial_j a_{ij} \in L^2_{L^q}$. Then, if one assumes (A1), (A2), (A4), and $(\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$ the existence and uniqueness of the weak solution to (3.37) can be proved similarly to [Theorem 3.2](#).

4. The regularity theorems

In this section we present some regularity results on distributional solutions of the Fokker–Planck equation. These then will be the key to achieving uniqueness in the class of distributional solutions in several contexts.

4.1. Local regularity theorem and uniqueness

Below we prove a regularity result which guarantees that distributional solutions of (FP) in the class $L^\infty((0, T); L^q(\mathbb{R}^d))$ are actually in $L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$, provided that we have enough regularity on $\mathbf{a}$ and integrability $\mathbf{b}$.

Theorem 4.1. *Let $p, q \geq 1$ such that $1/p + 1/q \leq 1/2$. Assume that $\mathbf{a} \in L^2((0, T); W^{1,p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ and $\mathbf{b} \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ satisfy (A2) and (A4) and $(\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Let $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ be a distributional solution to (FP), then $u \in L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$.*

Proof. To commence, we observe that $1/p + 1/q \leq 1/2$ clearly implies that both $p, q \geq 2$ and then any $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ belongs to $L^\infty((0, T); L^2_{\text{loc}}(\mathbb{R}^d))$. The strategy is to prove that $\nabla u \in L^2((0, T); L^2_{\text{loc}}(\mathbb{R}^d; \mathbb{R}^d))$ and this will be achieved exhibiting an approximating sequence $(u^\delta)_\delta$ enjoying uniform bounds on ∇u^δ : in turn, this estimate will be obtained as a consequence of [Lemma 2.4](#) and [Lemma 2.6](#).

Let $(\rho^\delta)_\delta$ be a standard family of mollifiers, then we know that the function $u^\delta := u * \rho^\delta$ solves Eq. (3.7) with $\tilde{\mathbf{b}}$ instead of $\mathbf{b}$. Consider a smooth function φ such that

$$\varphi \geq 0, \quad \operatorname{supp} \varphi \subset B_2, \quad \varphi = 1 \text{ in } B_1,$$

and define $\varphi_R(x) = \varphi(x/R)$. We multiply the equation in (3.7) by $u^\delta \varphi_R$ and we integrate over $(0, t) \times \mathbb{R}^d$. Then, integration by parts together with (A4) gives that

$$\begin{aligned} \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 \varphi_R dx + \alpha \int_0^t \int_{\mathbb{R}^d} |\nabla u^\delta|^2 \varphi_R dx ds &\leq \int_{\mathbb{R}^d} |u_0^\delta|^2 \varphi_R dx + \int_0^t \int_{\mathbb{R}^d} |u^\delta|^2 \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds \\ &\quad - \int_0^t \int_{\mathbb{R}^d} |u^\delta|^2 \operatorname{div} \tilde{\mathbf{b}} \varphi_R dx ds + \sum_{ij} \int_0^t \int_{\mathbb{R}^d} \partial_j a_{ij} \frac{|u^\delta|^2}{2} \partial_i \varphi_R dx ds \\ &\quad + \sum_{ij} \int_0^t \int_{\mathbb{R}^d} a_{ij} \frac{|u^\delta|^2}{2} \partial_j \partial_i \varphi_R dx ds + 2 \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) u^\delta \varphi_R dx ds. \end{aligned}$$

We now estimate separately the terms on the right hand side (the constant may change from line to line and is allowed to depend on $T, \|\varphi\|_{C^2}, \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty}, \|u\|_{L^\infty L^q}, \|u_0\|_{L^q}, \|\partial_j a_{ij}\|_\infty$):

- $\int_{\mathbb{R}^d} |u_0^\delta|^2 \varphi_R dx \leq \|u_0\|_{L^q}^2 \|\varphi_R\|_{L^{\frac{q}{q-2}}} \leq C \|u_0\|_{L^q}^2 R^{\frac{q-2}{q}}$;
- if s is such that $2/q + 1/p + 1/s = 1$, then we have

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} |u^\delta|^2 \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx dt &\leq \|u^\delta\|_{L^\infty L^q}^2 \|\tilde{\mathbf{b}}\|_{L^1 L^p} \|\nabla \varphi_R\|_{L^s} \\ &\leq C \|u\|_{L^\infty L^q}^2 \|\tilde{\mathbf{b}}\|_{L^1 L^p} \|\nabla \varphi_R\|_{L^s} \\ &\leq C R^{\frac{d}{s}-1} \|u\|_{L^\infty L^q}^2 \|\tilde{\mathbf{b}}\|_{L^1 L^p}; \end{aligned}$$

- since φ_R is positive, we can use the bound on the negative part of $\operatorname{div} \tilde{\mathbf{b}}$ to obtain that

$$\begin{aligned} - \int_0^T \int_{\mathbb{R}^d} |u^\delta|^2 \operatorname{div} \tilde{\mathbf{b}} \varphi_R dx ds &\leq \int_0^T \|(\operatorname{div} \tilde{\mathbf{b}})^-(s, \cdot)\|_\infty \int_{\mathbb{R}^d} |u^\delta|^2 \varphi_R dx ds \\ &\leq \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty} \|u^\delta\|_{L^\infty L^q}^2 \|\varphi_R\|_{L^{\frac{q}{q-2}}} \\ &\leq \|(\operatorname{div} \tilde{\mathbf{b}})^-\|_{L^1 L^\infty} \|u\|_{L^\infty L^q}^2 R^{\frac{q-2}{q}}; \end{aligned}$$

- we use (A2) to obtain that

$$\begin{aligned} \sum_{ij} \int_0^T \int_{\mathbb{R}^d} \partial_j a_{ij} \frac{|u^\delta|^2}{2} \partial_i \varphi_R dx ds &= \sum_i \int_0^T \int_{\mathbb{R}^d} \left(\sum_j \partial_j a_{ij} \right) \frac{|u^\delta|^2}{2} \partial_i \varphi_R dx ds \\ &\leq C \left\| \sum_j \partial_j a_{ij} \right\|_{L^\infty} \|u^\delta\|_{L^q}^2 \|\nabla \varphi_R\|_{L^{\frac{q}{q-2}}} \\ &\leq C \left\| \sum_j \partial_j a_{ij} \right\|_{L^\infty} \|u\|_{L^\infty L^q}^2 R^{d \frac{q-2}{q}-1}; \end{aligned}$$

- if s is such that $2/q + 1/p + 1/s = 1$, then we have

$$\int_0^T \int_{\mathbb{R}^d} a_{ij} \frac{|u^\delta|^2}{2} \partial_i \partial_j \varphi_R dx dt \leq C R^{\frac{d}{s}-2} \|u\|_{L^\infty L^q}^2 \|a_{ij}\|_{L^1 L^p}.$$

For the commutator terms, first of all notice that $u^\delta \varphi_R \in H_0^1(B_{2R})$. Then, we have that

- the transport commutator

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} r^\delta u^\delta \varphi_R dx dt &\leq C \|r^\delta\|_{L^2 H^{-1}(B_{2R})} \|u^\delta \varphi_R\|_{L^2 H^1(B_{2R})} \\ &\leq C \|r^\delta\|_{L^2 H^{-1}(B_{2R})} \left(\|u^\delta \varphi_R\|_{L^\infty L^2(B_{2R})} + \|\nabla(u^\delta \varphi_R)\|_{L^2 L^2(B_{2R})} \right); \end{aligned}$$

- the diffusion commutator

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} s^\delta u^\delta \varphi_R dx dt &\leq C \|s^\delta\|_{L^2 H^{-1}(B_{2R})} \|u^\delta \varphi_R\|_{L^2 H^1(B_{2R})} \\ &= C \|s^\delta\|_{L^2 H^{-1}(B_{2R})} \left(\|u^\delta \varphi_R\|_{L^\infty L^2(B_{2R})} + \|\nabla(u^\delta \varphi_R)\|_{L^2 L^2(B_{2R})} \right). \end{aligned}$$

We now use Young inequality to obtain the bound

$$\int_0^T \int_{\mathbb{R}^d} (r^\delta + s^\delta) u^\delta \varphi_R dx dt \leq C_\alpha (\|r^\delta\|_{L^2 H^{-1}(B_{2R})}^2 + \|s^\delta\|_{L^2 H^{-1}(B_{2R})}^2) + \frac{\alpha}{2} \|u^\delta \varphi_R\|_{L^2 H^1(B_{2R})}^2.$$

Then, $0 \leq \varphi_R \leq 1$ implies $\varphi_R^2 \leq \varphi_R$, thus

$$\|u^\delta \varphi_R\|_{L^2 H^1(B_{2R})}^2 \leq \int_0^T \int_{B_{2R}} |u^\delta|^2 \varphi_R dx ds + \int_0^T \int_{B_{2R}} |\nabla u^\delta|^2 \varphi_R dx ds + \int_0^T \int_{B_{2R}} |u^\delta|^2 |\nabla \varphi_R|^2 dx ds.$$

The first and the last summand can be estimated similarly as above and gives respectively

$$\begin{aligned} \int_0^T \int_{B_{2R}} |u^\delta|^2 \varphi_R dx ds &\leq C \|u\|_{L^\infty L^q}^2 R^{d \frac{q-2}{q}}, \\ \int_0^T \int_{B_{2R}} |u^\delta|^2 |\nabla \varphi_R|^2 dx ds &\leq C \|u\|_{L^\infty L^q}^2 R^{d \frac{q-2}{q}-2}. \end{aligned}$$

All in all,

$$\begin{aligned} \alpha \int_0^t \int_{\mathbb{R}^d} |\nabla u^\delta|^2 \varphi_R dx ds &\leq C \left(R^{d \frac{q-2}{q}} + R^{\frac{d}{s}-1} + R^{d \frac{q-2}{q}-1} \right) + C_\alpha (\|r^\delta\|_{L^2 H^{-1}(B_{2R})}^2 + \|s^\delta\|_{L^2 H^{-1}(B_{2R})}^2) \\ &\quad + \frac{\alpha}{2} \int_0^T \int_{B_{2R}} |\nabla u^\delta|^2 \varphi_R dx ds. \end{aligned}$$

Passing to the supremum in t in the left hand side we obtain that

$$\begin{aligned} \frac{\alpha}{2} \int_0^T \int_{B_R} |\nabla u^\delta|^2 dx ds &\leq \frac{\alpha}{2} \int_0^T \int_{\mathbb{R}^d} |\nabla u^\delta|^2 \varphi_R dx ds \\ &\leq C \left(R^{d \frac{q-2}{q}} + R^{\frac{d}{s}-1} + R^{d \frac{q-2}{q}-1} \right) + C_\alpha (\|r^\delta\|_{L^2 H^{-1}(B_{2R})}^2 + \|s^\delta\|_{L^2 H^{-1}(B_{2R})}^2). \end{aligned}$$

Finally, the commutators r^δ, s^δ go to 0 in $L^2 H_{\text{loc}}^{-1}$ by Lemmas 2.3 and 2.6, thus they are uniformly bounded for δ small enough (possibly depending on R), i.e.

$$\|r^\delta\|_{L^2 H^{-1}(B_{2R})}^2 + \|s^\delta\|_{L^2 H^{-1}(B_{2R})}^2 \leq \bar{C}(R),$$

for some constant $\bar{C}(R)$ which may diverge as $R \rightarrow \infty$. Therefore, we have

$$\|\nabla u^\delta\|_{L^2 L^2(B_R)}^2 \leq C \left(R^{\frac{d-2}{q}} + R^{\frac{d}{s}-1} + R^{d\frac{q-2}{q}-1} \right) + \bar{C}(R), \quad (4.1)$$

which is finite uniformly in δ for any $R > 0$. This proves that ∇u exists and belongs to $L^2((0, T); L^2(B_R))$. $\square$

Remark 4.2. Notice that in the proof of [Theorem 4.1](#) we do not use the bound

$$\sup_{t \in (0, T)} \|u(t, \cdot)\|_{L^q} \leq \|u_0\|_{L^q}, \quad (4.2)$$

since u is any given distributional solution and a priori we do not know if the energy bound (4.2) holds. However, a posteriori this is true under the assumptions of [Theorem 4.1](#): in fact, it can be easily proved using the regularity of the solution and the convergence of the commutators r^δ, s^δ to 0 in $L^1_{t,x,\text{loc}}$.

Remark 4.3. In addition to regularity, [Theorem 4.1](#) proves the existence of solutions in the class $L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$. It would be interesting to prove the existence of solutions in this class with a more direct argument, instead of relying on regularity.

Combining [Theorems 3.13](#) and [4.1](#), we obtain the result which is transverse to that of [\[12\]](#).

Theorem 4.4. Let $\mathbf{b} \in L^\infty((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d))$ and $\mathbf{a} \in L^\infty((0, T); W^{1,\infty}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ satisfy (A2), (A3), and (A4). Then there exists at most one distributional solution $u \in L^2((0, T) \times \mathbb{R}^d)$.

Proof. Note that if we consider a distributional solution $u \in L^2((0, T); L^q(\mathbb{R}^d))$, arguing as in [Theorem 4.1](#) it is easy to show that $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ if one assume that $\bar{\mathbf{b}} \in L^\infty_t L^p_x$, $\mathbf{a} \in L^\infty_t W^{1,p}_x$, and $(\text{div} \bar{\mathbf{b}})^- \in L^\infty_{t,x}$ (the estimates depend on $\|u\|_{L^2 L^q}$). Thus, under our assumptions, if we set $p = \infty, q = 2$, the solution u belongs to $L^\infty((0, T); L^2(\mathbb{R}^d))$. Then, it is enough to repeat the proof of [Theorem 4.1](#) noticing that the estimate (4.1) is now uniform in R . In fact, by [Remark 2.5](#) the commutators converge in $L^2((0, T); H^{-1}(\mathbb{R}^d))$, thus the constant $\bar{C}$ is uniformly bounded in R . Moreover, if we substitute $q = 2$ in the powers of R in (4.1), we obtain that

$$\limsup_{R \rightarrow \infty} \|\nabla u\|_{L^2 L^2(B_R)} < \infty.$$

This implies the global information $\nabla u \in L^2((0, T); L^2(\mathbb{R}^d; \mathbb{R}^d))$. Then, we can conclude that $u \in L^\infty((0, T); L^2(\mathbb{R}^d)) \cap L^2((0, T); H^1(\mathbb{R}^d))$ and the uniqueness follows from [Theorem 3.13](#). $\square$

4.2. Extensions with suitable growth conditions

We now discuss the role of the growth assumptions on $\mathbf{b}, \mathbf{a}$ to recover further uniqueness and regularity results for the Fokker–Planck equation (FP). We show that under suitable growth assumption uniqueness holds in the local parabolic class $L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$. In particular, weak solutions of (FP-div) are also unique in $L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$ as a consequence of [Proposition 3.11](#). The theorem is the following.

Theorem 4.5. Let $u_0 \in L^q \cap L^\infty(\mathbb{R}^d)$ be a given initial datum with $q > 2$ and assume that $\mathbf{b} \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ and $\mathbf{a} \in L^2((0, T); W^{1,p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ with $1/p + 1/q = 1/2$ satisfy (A2) and (A4), and $(\text{div} \bar{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Moreover, we assume the following growth conditions

$$\begin{aligned} \frac{\mathbf{b}(t, x)}{1 + |x|}, \frac{\partial_j a_{ij}(t, x)}{1 + |x|} &\in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d)), \\ \frac{\mathbf{a}(t, x)}{1 + |x|^2} &\in L^1((0, T); L^1(\mathbb{R}^d; \mathbb{R}^{d \times d})) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^{d \times d})). \end{aligned}$$

Then, there exists at most one distributional solution $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$.

Proof. The proof is similar to the one of [Theorem 3.2](#) so we just sketch it. Let $u \in L^\infty((0, T); L^q(\mathbb{R}^d))$ be a distributional solution of (FP) whose existence is proved in [Proposition 3.9](#). Let ρ^δ be a standard mollifier, we have that the function $u^\delta := u * \rho^\delta$ satisfies the equation

$$\begin{cases} \partial_t u^\delta + \text{div}(\bar{\mathbf{b}} u^\delta) - \frac{1}{2} \sum_{ij} \partial_i (a_{ij} \partial_j u^\delta) = r^\delta + s^\delta, \\ u^\delta(0, \cdot) = u_0 * \rho_\delta. \end{cases} \quad (4.3)$$

Given our assumptions on p and q , [Theorem 4.1](#) implies that the solution u belongs to the space $L^2((0, T); H^1_{\text{loc}}(\mathbb{R}^d))$. Thus, as a consequence of [Lemmas 2.4](#) and [2.6](#) we know that $r^\delta, s^\delta \rightarrow 0$ in $L^2((0, T); H^{-1}_{\text{loc}}(\mathbb{R}^d))$. Consider a smooth function φ such that

$$\varphi \geq 0, \quad \text{supp } \varphi \subset B_2, \quad \varphi = 1 \text{ in } B_1,$$

and define $\varphi_R(x) = \varphi(x/R)$. Let $\beta : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined as

$$\beta(z) = |z|^q.$$

The following argument can be made rigorous by considering a regularized version of the function β as we did in the proof of [Theorem 3.2](#). We skip this technical detail for simplicity of exposition. We multiply Eq. (4.3) by $\beta'(u^\delta)\varphi_R$ and with computations similar to those of the proof of [Theorem 3.2](#) we get that

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d} \operatorname{div}(\tilde{\mathbf{b}}u^\delta)\beta'(u^\delta)\varphi_R dx ds &= - \int_0^t \int_{\mathbb{R}^d} u^\delta \tilde{\mathbf{b}} \cdot \nabla u^\delta \beta''(u^\delta)\varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds \\ &= - (q-1) \int_0^t \int_{\mathbb{R}^d} \tilde{\mathbf{b}} \cdot \nabla \beta(u^\delta)\varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds \\ &= (q-1) \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}} \beta(u^\delta)\varphi_R dx ds + (q-1) \int_0^t \int_{\mathbb{R}^d} \beta(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} u^\delta \beta'(u^\delta) \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds. \end{aligned}$$

On the other hand, for the term involving the diffusion we use the assumption on the divergence obtaining that

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^d} \partial_i(a_{ij}\partial_j u^\delta)\beta'(u^\delta)\varphi_R dx ds &= - \int_0^t \int_{\mathbb{R}^d} a_{ij}\partial_j u^\delta \partial_i u^\delta \beta''(u^\delta)\varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} a_{ij}\partial_j u^\delta \beta'(u^\delta) \partial_i \varphi_R dx ds \\ &= - \int_0^t \int_{\mathbb{R}^d} a_{ij}\partial_j u^\delta \partial_i u^\delta \beta''(u^\delta)\varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} a_{ij}\partial_j \beta(u^\delta) \partial_i \varphi_R dx ds \\ &= - \int_0^t \int_{\mathbb{R}^d} a_{ij}\partial_j u^\delta \partial_i u^\delta \beta''(u^\delta)\varphi_R dx ds + \int_0^t \int_{\mathbb{R}^d} \partial_j a_{ij} \beta(u^\delta) \partial_i \varphi_R dx ds + \int_0^t \int_{\mathbb{R}^d} a_{ij} \beta(u^\delta) \partial_j \partial_i \varphi_R dx ds. \end{aligned}$$

Then, we use the uniform ellipticity of $\mathbf{a}$ to obtain

$$0 \leq \alpha \int_0^T \int_{\mathbb{R}^d} |\nabla u^\delta|^2 \beta''(u^\delta)\varphi_R dx ds \leq \int_0^T \int_{\mathbb{R}^d} a_{ij}\partial_i u^\delta \partial_j u^\delta \beta''(u^\delta)\varphi_R dx ds. \quad (4.4)$$

We finally have the following inequality

$$\begin{aligned} \int_{\mathbb{R}^d} \beta(u^\delta)\varphi_R dx &\leq -(q-1) \int_0^t \int_{\mathbb{R}^d} \operatorname{div} \tilde{\mathbf{b}} \beta(u^\delta)\varphi_R dx ds + C \int_0^t \int_{\mathbb{R}^d} |\tilde{\mathbf{b}}| \beta(u^\delta) |\nabla \varphi_R| dx ds + \int_0^t \int_{\mathbb{R}^d} a_{ij} \beta(u^\delta) \partial_i \partial_j \varphi_R dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d} \partial_j a_{ij} \beta(u^\delta) \partial_i \varphi_R dx ds + \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) \beta'(u^\delta)\varphi_R dx ds. \end{aligned}$$

Since $\beta'(u^\delta)\varphi_R \in H_0^1(B_{2R})$ we can let $\delta \rightarrow 0$ to obtain that

$$\begin{aligned} \int_{\mathbb{R}^d} \beta(u)\varphi_R dx &\leq -(q-1) \int_0^t \int_{\mathbb{R}^d} \operatorname{div}(\tilde{\mathbf{b}})\beta(u)\varphi_R dx ds + C \int_0^t \int_{\mathbb{R}^d} \beta(u) |\tilde{\mathbf{b}}| |\nabla \varphi_R| dx ds + \int_0^t \int_{\mathbb{R}^d} \beta(u) a_{ij} \partial_i \partial_j \varphi_R dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}^d} \beta(u) \partial_j a_{ij} \partial_i \varphi_R dx ds = I + II + III + IV. \end{aligned}$$

Then, we can estimate I and II as in the proof of [Theorem 3.2](#), while we use the growth assumptions on $\mathbf{a}$ to estimate III and IV as follows

$$\begin{aligned} I &\lesssim \int_0^t \|(\operatorname{div} \tilde{\mathbf{b}}(s, \cdot))^- \|_\infty \int_{\mathbb{R}^d} \beta(u)\varphi_R dx ds, \\ II &\lesssim \|u_0\|_\infty^q \int_0^T \int_{|x|>R} |\tilde{\mathbf{b}}_1(s, x)| dx ds + \int_0^T \|\tilde{\mathbf{b}}_2(s, \cdot)\|_\infty \int_{|x|>R} |u(s, x)|^q dx ds, \\ III &\lesssim \|u_0\|_\infty^q \int_0^T \int_{|x|>R} |\mathbf{a}_1(s, x)| dx ds + \int_0^T \|\mathbf{a}_2(s, \cdot)\|_\infty \int_{|x|>R} |u(s, x)|^q dx ds, \\ IV &\lesssim \|u_0\|_\infty^q \int_0^T \int_{|x|>R} |\partial_j a_{ij}^1(s, x)| dx ds + \int_0^T \|\partial_j a_{ij}^2(s, \cdot)\|_\infty \int_{|x|>R} |u(s, x)|^q dx ds, \end{aligned}$$

where it is worth to point out that in the estimate for III we used that $\partial_i \partial_j \varphi_R \leq \frac{C}{R^2}$ together with the growth assumption on $\mathbf{a}$. Moreover, all the implicit constants in the estimate depends on the C^2 norm of φ only. Thus, by defining $F(R) := II + III + IV$, which vanishes as $R \rightarrow \infty$, we get the following

$$\int_{\mathbb{R}^d} \beta(u)\varphi_R dx \leq \int_0^t \|(\operatorname{div} \tilde{\mathbf{b}}(s, \cdot))^- \|_\infty \int_{\mathbb{R}^d} \beta(u)\varphi_R dx ds + F(R),$$

and finally, we apply Gronwall's inequality to obtain that

$$\int_{\mathbb{R}^d} \beta(u)\varphi_R dx \leq F(R) \exp \left(\|(\operatorname{div} \tilde{\mathbf{b}})^- \|_{L^1 L^\infty} \right),$$

and by letting $R \rightarrow \infty$ we obtain that $|u| = 0$ for a.e. $(t, x) \in (0, T) \times \mathbb{R}^d$. $\square$

If we assume slightly stronger growth conditions, there is a regime where the solution is parabolic instead of just locally parabolic. The result is the following.

Theorem 4.6. Let $u \in L^\infty((0, T); L^2 \cap L^q(\mathbb{R}^d))$ be a distributional solution of (FP), and let $\mathbf{b} \in L^2((0, T); L^p(\mathbb{R}^d; \mathbb{R}^d))$ and $\mathbf{a} \in L^2((0, T); W^{1,p}(\mathbb{R}^d; \mathbb{R}^{d \times d}))$ with $1/p + 1/q = 1/2$ satisfy (A2) and (A4), and $(\operatorname{div} \tilde{\mathbf{b}})^- \in L^1((0, T); L^\infty(\mathbb{R}^d))$. Moreover, assume that $2 < q \leq \frac{2d}{d-2}$ together with the following growth conditions

$$\frac{\mathbf{b}(t, x)}{1 + |x|}, \frac{\partial_j a_{ij}(t, x)}{1 + |x|} \in L^1((0, T); L^{\frac{q}{q-2}}(\mathbb{R}^d; \mathbb{R}^d)) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^d)),$$

$$\frac{\mathbf{a}(t, x)}{1 + |x|^2} \in L^1((0, T); L^{\frac{q}{q-2}}(\mathbb{R}^d; \mathbb{R}^{d \times d})) + L^1((0, T); L^\infty(\mathbb{R}^d; \mathbb{R}^{d \times d})).$$

Then $u \in L^2((0, T); H^1(\mathbb{R}^d))$.

Proof. We can proceed as in Theorem 4.1: we consider the equation for u^δ and we multiply it by $2u^\delta \varphi_R$ where φ_R is defined as in Theorem 4.5 and we obtain the estimate

$$\begin{aligned} \int_{\mathbb{R}^d} |u^\delta(t, x)|^2 \varphi_R dx + \alpha \int_0^t \int_{\mathbb{R}^d} |\nabla u^\delta|^2 \varphi_R dx ds &\leq \int_{\mathbb{R}^d} |u_0^\delta|^2 \varphi_R dx + \int_0^t \int_{\mathbb{R}^d} |u^\delta|^2 \tilde{\mathbf{b}} \cdot \nabla \varphi_R dx ds - \int_0^t \int_{\mathbb{R}^d} |u^\delta|^2 \operatorname{div} \tilde{\mathbf{b}} \varphi_R dx ds \\ &+ \sum_{ij} \int_0^t \int_{\mathbb{R}^d} \partial_j a_{ij} \frac{|u^\delta|^2}{2} \partial_i \varphi_R dx ds + \sum_{ij} \int_0^t \int_{\mathbb{R}^d} a_{ij} \frac{|u^\delta|^2}{2} \partial_j \partial_i \varphi_R dx ds \\ &+ 2 \int_0^t \int_{\mathbb{R}^d} (r^\delta + s^\delta) u^\delta \varphi_R dx ds. \end{aligned}$$

The difference with respect to the proof of Theorem 4.1 is in the estimates of the second, fourth and fifth term on the right hand side above. For example, for the second term we use the growth assumption on $\mathbf{b}$ to get that

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^d} |u^\delta|^2 \mathbf{b} \cdot \nabla \varphi_R dx dt &\leq C \int_0^T \int_{R < |x| < 2R} |u^\delta|^2 \frac{|\mathbf{b}|}{1 + |x|} dx dt \\ &\leq C \|u_0\|_{L^2}^2 \left\| \frac{|\mathbf{b}_2|}{1 + |x|} \right\|_{L^\infty} + C \|u_0\|_{L^q}^2 \underbrace{\int_0^T \left(\int_{|x| > R} \left(\frac{|\mathbf{b}_1|}{1 + |x|} \right)^{\frac{q}{q-2}} dx \right)^{\frac{q-2}{q}} dt}_{(*) := f(R)} \end{aligned}$$

where $f(R) \rightarrow 0$ as $R \rightarrow \infty$. Similarly, we can estimate the other two terms using the growth assumptions on $\mathbf{a}$. Then, by using that the commutators are uniformly bounded in δ , we proceed as in Theorem 4.1 and we obtain that

$$\|\nabla u^\delta\|_{L^2 L^2(B_R)}^2 \leq C \left(1 + F(R) + R^{\frac{q-2}{q}-2} \right),$$

for some function $F(R)$ which vanishes as $R \rightarrow \infty$. Thus we have obtained that $\|\nabla u^\delta\|_{L^2 L^2(B_R)}^2$ is uniformly bounded (in δ and R) and this concludes the proof. $\square$

Combining Theorem 4.6 with the uniqueness result for parabolic solutions Theorem 3.13 we obtain the following:

Corollary 4.7. Assume that the hypothesis of Theorem 4.6 are fulfilled. In addition assume that $\mathbf{a}$ satisfies (A1). Then, any distributional solution $u \in L^\infty((0, T); L^2 \cap L^q(\mathbb{R}^d))$ of (FP) is parabolic and therefore unique.

Acknowledgments

During the preparation of this manuscript, G. Ciampa has been supported by the ERC Starting Grant 101039762 HamDyWWa. He is currently supported by IndAM-GNAMPa, and by the projects PRIN2020 “Nonlinear evolution PDEs, fluid dynamics and transport equations: theoretical foundations and applications” and PRIN2022 “Classical equations of compressible fluids mechanics: existence and properties of non-classical solutions”. G. Crippa is supported by SNF Project 212573 FLUTURA – Fluids, Turbulence, Advection. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

Data availability

No data was used for the research described in the article.

References

- [1] V. Barbu, M. Röckner, From nonlinear Fokker–Planck equations to solutions of distribution dependent SDE, *Ann. Probab.* 48 (4) (2020) 1902–1920.
- [2] P. Bonicatto, G. Ciampa, G. Crippa, Advection-diffusion equation with rough coefficients: weak solutions and vanishing viscosity, *J. Math. Pures Appl.* 167 (2022) 204–224.
- [3] P. Bonicatto, G. Ciampa, G. Crippa, Weak and parabolic solutions of advection-diffusion equations with rough velocity field, *J. Evol. Equ.* 24 (1) (2024) 1.

- [4] F. Bouchut, G. Crippa, Lagrangian flows for vector fields with gradient given by a singular integral, *J. Hyperbolic Differ. Equ.* 10 (2013) 235–282.
- [5] E. Brué, Q.-H. Nguyen, Advection diffusion equation with Sobolev velocity fields, *Comm. Math. Phys.* 383 (2021) 465–487.
- [6] P. Cardaliaguet, P. Jameson Graber, A. Porretta, D. Tonon, Second order mean field games with degenerate diffusion and local coupling, *Nonlinear Diff. Eq. Appl.* 22 (2015) 1287–1317.
- [7] N. Champagnat, P. Jabin, Strong solutions to stochastic differential equations with rough coefficients, *Ann. Probab.* 46 (3) (2018) 1498–1541.
- [8] G. Ciampa, F. Rossi, Vanishing viscosity in mean-field optimal control, *ESAIM:COCV* 29 (2023) 29.
- [9] R.J. DiPerna, P.-L. Lions, Ordinary differential equations, transport theory and Sobolev spaces, *Invent. Math.* 98 (1989) 511–547.
- [10] T.D. Drivas, T.M. Elgindi, G. Iyer, I.J. Jeong, Anomalous dissipation in passive scalar transport, *Arch. Ration. Mech. Anal.* 243 (2022) 1151–1180.
- [11] L.C. Evans, *Partial Differential Equations*, second ed., in: Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 2010.
- [12] A. Figalli, Existence and uniqueness of martingale solutions for SDEs with rough or degenerate coefficients, *J. Funct. Anal.* 254 (1) (2008) 109–153.
- [13] T.D. Frank, *Nonlinear Fokker–Planck Equations*, Springer, Berlin, Heidelberg, New York, 2005.
- [14] N.V. Krylov, M. Röckner, Strong solutions of stochastic equations with singular time dependent drift, *Probab. Theory Related Fields* 131 (2) (2005) 154–196.
- [15] O.A. Ladyženskaja, V.A. Solonnikov, N.N. Ural'ceva, *Linear and Quasilinear Equations of Parabolic Type*, in: Translations of Mathematical Monographs, vol. 23, American Mathematical Society, Providence, R.I., 1968.
- [16] J.M. Lasry, P.L. Lions, Mean field games, *Jpn. J. Math.* 2 (2007) 229–260.
- [17] C. Le Bris, P.-L. Lions, Renormalized solutions of some transport equations with partially $W^{1,1}$ velocities and applications, *Ann. di Mat.* 183 (2004) 97–130.
- [18] C. Le Bris, P.-L. Lions, Existence and uniqueness of solutions to Fokker–Planck type equations with irregular coefficients, *Comm. Partial Differential Equations* 33 (7) (2008) 1272–1317.
- [19] C. Le Bris, P.-L. Lions, *Parabolic Equations with Irregular Data and Related Issues*, De Gruyter, 2019.
- [20] H. Li, D. Luo, Quantitative stability estimates for Fokker–Planck equations, *J. Math. Pures Appl.* 122 (9) (2019) 125–163.
- [21] S. Modena, G. Sattig, Convex integration solutions to the transport equation with full dimensional concentration, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 37 (5) (2020) 1075–1108.
- [22] S. Modena, L. Székelyhidi Jr., Non-uniqueness for the transport equation with Sobolev vector fields, *Ann. PDE* 4 (2) (2018) 18.
- [23] S. Modena, L. Székelyhidi Jr., Non-renormalized solutions to the continuity equation, *Calc. Var.* 58 (2019) 208.
- [24] V. Navarro-Fernández, A. Schlichting, C. Seis, Optimal stability estimates and a new uniqueness result for advection-diffusion equations, *Pure Appl. Anal.* 4 (3) (2022) 571–596.
- [25] H.-C. Ottinger, *Stochastic Processes in Polymeric Fluids*, Springer, 1996.
- [26] R.G. Owens, T.N. Phillips, *Computational Rheology*, Imperial College Press, 2002.
- [27] A. Porretta, Weak solutions to Fokker–Planck equations and mean field games, *Arch. Ration. Mech. Anal.* 216 (2015) 1–62.
- [28] V. Schwämmle, F.D. Nobre, E.M.F. Curado, Consequences of the H theorem from nonlinear Fokker–Planck equations, *Phys. Rev. E* 76, 041123.
- [29] D.W. Stroock, S.R.S. Varadhan, Multidimensional diffusion processes, in: Grundlehren Math. Wiss., vol. 233, Springer-Verlag, Berlin, 1979.