















# Germline Findings From Tumor-Only Comprehensive Genomic Profiling in the RATIONAL Study: A Missed Opportunity?

Riziero Esposito Abate, PhD<sup>1</sup>; Alessandro Morabito, MD<sup>2</sup>; Michele Milella, MD<sup>3</sup> ; Fabrizio Tabbò, MD<sup>4</sup>; Valentina Guarneri, MD<sup>5,6</sup> ; Giacomo Pelizzari, MD<sup>7</sup>; Ilario G. Rapposelli, MD<sup>8</sup> ; Rossana Berardi, MD<sup>9</sup> ; Lucio Buffoni, MD<sup>10</sup>; Elisa Bennicelli, MD<sup>11</sup>; Francesca Zanelli, MD<sup>12</sup>; Carlo Tondini, MD<sup>13</sup>; Laura Attademo, MD<sup>14</sup>; Tiziana P. Latiano, MD<sup>15</sup> ; Salvatore Corallo, MD<sup>16,17</sup> ; Giancarlo Pruneri, MD<sup>18</sup> ; Federica Marmorino, MD<sup>19</sup> ; Orazio Caffo, MD<sup>20</sup> ; Lorenzo Antonuzzo, MD<sup>21,22</sup>; Simona Tessitore, PhD<sup>1</sup>; Silvia Novello, MD<sup>23</sup> ; Giuseppe Curigliano, MD<sup>24,25</sup> ; Carmine Pinto, MD<sup>12</sup> ; Nicola Normanno, MD<sup>26</sup> ; and Antonella De Luca, PhD<sup>1</sup> 

DOI <https://doi.org/10.1200/PO-25-00399>

## ABSTRACT

**PURPOSE** Tumor comprehensive genomic profiling (CGP) may detect potential germline pathogenic/likely pathogenic (P/LP) alterations as secondary findings. We analyzed the frequency of potentially germline variants and large rearrangements (LRs) in the RATIONAL study, an Italian multicenter, observational clinical trial that collects next-generation sequencing–based tumor profiling data, and evaluated how these findings were managed by the enrolling centers.

**PATIENTS AND METHODS** Patients prospectively enrolled in the pathway-B of the RATIONAL study and undergoing CGP with the FoundationOne CDx assays were included in the analysis. Potentially germline variants detected in 40 cancer susceptibility genes (CSGs) were classified in three classes with different actionability, most (MA), high (HA), and standard (SA), on the basis of penetrance, mutational spectrum, and intervention for prevention/early detection.

**RESULTS** On the basis of the European Society of Medical Oncology recommendations, we identified 225 potentially germline P/LP variants in 193/1,339 (14.4%) enrolled patients. In particular, 62/225 (27.5%) variants were detected in genes classified as MA-CSG class, 53/225 (23.6%) in genes belonging to the HA-CSG class, and 110/225 (48.9%) in the SA-CSG class. In addition, we detected 58 LRs in the 16/40 CSGs in 53/1,339 (3.95%) patients. Information about germline-focused analysis and follow-up was available for 99 patients with potentially germline variants. Surprisingly, 95/99 (96%) patients were not referred to oncogenetic consultation and follow-up, including 30/32 (93.75%) patients with variants in the MA-CSG class.

**CONCLUSION** Our data confirm the utility of CGP for the identification of potentially germline variants in CSGs, highlighting the importance of reporting LRs in addition to single-nucleotide variants and insertions/deletions. However, our findings also demonstrate a relative lack of knowledge of the implications of germline findings detected on tumor-only sequencing among oncologists and underline the need for specific training in this area.

## ACCOMPANYING CONTENT

-  [Data Supplement](#)
-  [Visual Abstract](#)

Accepted July 30, 2025

Published September 10, 2025

JCO Precis Oncol 9:e2500399

© 2025 by American Society of Clinical Oncology

Creative Commons Attribution  
Non-Commercial No Derivatives  
4.0 License

## INTRODUCTION

Tumor comprehensive genomic profiling (CGP) with panels that allow for a broad molecular characterization, including the detection of mutational signatures and complex biomarkers (ie, tumor mutational burden [TMB], microsatellite instability, etc), can reveal actionable genomic alterations (GAs) for targeted therapeutic intervention in patients with solid tumors.<sup>1-6</sup> Although CGP has been implemented to identify somatic variants in tumor tissue or plasma samples,

it may also detect potential germline pathogenic/likely pathogenic (P/LP) alterations as secondary findings.<sup>5,7,8</sup> The identification of germline alterations associated with increased cancer risk might have important clinical implications for patients and their relatives, in terms of cancer prevention and long-term follow-up.<sup>1,9,10</sup>

As CGP on tumor-only and plasma-only samples is not able to discriminate somatic from germline alterations, a confirmatory germline test should be performed for presumed

## CONTEXT

### Key Objective

To analyze the frequency of potential germline variants and large rearrangements (LRs) in cancer susceptibility genes (CSGs) in Italian patients with cancer undergoing tumor- or plasma-only sequencing with a validated next-generation sequencing test, and to evaluate the management of this information by the oncologists.

### Knowledge Generated

A significant fraction of patients with advanced cancer carried potentially germline variants and LRs in CSGs. Genomic alterations in CSGs were detected also in tumors that do not usually undergo genetic surveillance, such as cancer of unknown primary, cholangiocarcinoma, and lung cancer. However, only 4% of patients carrying variants in CSGs were referred to genetic counseling.

### Relevance

Our findings support the testing of CSGs in patients with cancer, to identify carriers in which health surveillance programs and interventions can lead to early diagnosis and prevention. Our data also highlight the need for specific training of oncologists, to ensure the correct interpretation of complex genomic data.

germline variants.<sup>9</sup> The European Society of Medical Oncology (ESMO) developed recommendations for confirmatory germline analysis and follow-up of tumor-only CGP.<sup>11,12</sup> The most recent recommendations suggest germline follow-up for single-nucleotide variants (SNVs) and small insertions/deletions (indels) identified in 40 cancer susceptibility genes (CSGs),<sup>11</sup> categorized into three classes on the basis of evidence of risk (Data Supplement, Table S1).

We analyzed the frequency of potential germline P/LP alterations detected in the RATIONAL study, an Italian multicenter, observational clinical trial that collects next-generation sequencing (NGS)-based tumor profiling data from patients with more of 40 tumor types. In addition, we evaluated how the discovery of possible germline variants was managed by the centers participating in the study.

## PATIENTS AND METHODS

The RATIONAL study (ClinicalTrials.gov identifier: [NCT05918666](https://clinicaltrials.gov/ct2/show/study/NCT05918666)) is a multicentric, observational study approved by the Ethics Committee of the Istituto Nazionale Tumori Fondazione G. Pascale (Prot. 6/17 OSS) and by the ethics committees of other 48 participating institutions. All patients provided a specific written informed consent to participate to the study. Patients are informed about all GAs, including those that could affect family risk, unless patients opt out.

Between October 2018 and July 2024, 1,986 patients were enrolled, 390 in pathway-A and 1,596 in pathway-B. In pathway-A, patients with advanced solid tumor who already received a NGS-based tumor profiling are registered. In pathway-B, patients for whom no access to NGS was available and with specific clinical and pathologic features are offered access to the FoundationOne CDx assays

(Foundation Medicine Inc, Cambridge, MA) on either tumor tissue or blood samples. Additional information is provided in the Data Supplement.

## RESULTS

### Patients

Between October 2018 and July 2024, FoundationOne CDx reports were successfully released for 1,341 patients enrolled in pathway-B. Two patients did not meet the inclusion criteria (age <18 years) and they were excluded from further analyses. Therefore, 1,339 patients were included in the study.

The clinical and pathologic characteristics of the whole cohort are described in [Table 1](#).

CGP was performed on tissue with the FoundationOne CDx assay for 944 (70.5%) patients and on circulating tumor DNA (ctDNA) with the FoundationOne Liquid CDx test for 395 (29.5%) patients ([Fig 1A](#)).

### Identification of Potentially Germline P/LP Variants by Tumor-Only CGP

Sequencing analysis revealed the presence of at least one GA with known significance in 1,297/1,339 (96.9%) patients. Overall, 6,807 GAs in 291 genes were identified, with a mean of 5.25 GAs per patient.

We evaluated the frequency of potentially germline P/LP GAs in the 40 CSGs of the ESMO recommendations, including SNVs, indels, and large rearrangements (LRs). We next applied the suggested variant allelic frequency (VAF) thresholds (VAF > 30% for SNVs and >20% for indels).<sup>11</sup> By

**TABLE 1.** Clinicopathologic Characteristics of Patients Enrolled in the RATIONAL Study Included in the Analysis

| Characteristic                    | All (N = 1,339) |
|-----------------------------------|-----------------|
| Age, years, median (range)        | 62 (18-87)      |
| Age <30 years, patients, No. (%)  | 26 (1.94)       |
| Sex, No. (%)                      |                 |
| Female                            | 624 (46.6)      |
| Male                              | 715 (53.4)      |
| Ancestry, %                       |                 |
| Caucasian                         | >99             |
| Family history of cancer, No. (%) |                 |
| Yes                               | 262 (19.57)     |
| No                                | 389 (29.05)     |
| NA                                | 688 (51.38)     |
| Tumor type, No. (%)               |                 |
| Lung cancer                       | 342 (25.54)     |
| Biliary tract cancer              | 271 (20.24)     |
| Pancreatic cancer                 | 165 (12.32)     |
| Colorectal cancer                 | 112 (8.36)      |
| Esophagogastric cancer            | 107 (7.99)      |
| CUP                               | 85 (6.35)       |
| Breast cancer                     | 54 (4.03)       |
| Soft tissue cancer                | 35 (2.61)       |
| Head and neck cancer              | 34 (2.54)       |
| Ovarian cancer                    | 15 (1.12)       |
| Renal cancer                      | 15 (1.12)       |
| Small bowel cancer                | 12 (0.90)       |
| Uterine cancer                    | 12 (0.90)       |
| Bladder cancer                    | 12 (0.90)       |
| CNS cancer                        | 11 (0.82)       |
| Liver cancer                      | 10 (0.75)       |
| Thymus cancer                     | 9 (0.67)        |
| Pleural mesothelioma              | 8 (0.60)        |
| Melanoma                          | 8 (0.60)        |
| Thyroid cancer                    | 6 (0.45)        |
| Skin cancer                       | 5 (0.37)        |
| Prostate cancer                   | 3 (0.22)        |
| Adrenal cancer                    | 2 (0.15)        |
| Carcinoma of the vulva/vagina     | 2 (0.15)        |
| Penile cancer                     | 2 (0.15)        |
| Anal cancer                       | 1 (0.07)        |
| Testicular cancer                 | 1 (0.07)        |

Abbreviations: CUP, cancer of unknown primary; NA, not applicable.

applying these criteria, one or more potentially germline P/LP variants above the VAF threshold were observed in 193/1,339 (14.4%) patients, including 23/193 patients (11.9%) with at least two potentially germline variants (Fig 1A). Collectively, we classified 225 variants in 30 genes (124 SNVs and 101 indels) as potentially germline. The most common genes in which potentially germline variants were observed pan-cancer were ATM (15.1% of potentially germline

variants), MUTYH (10.7%), NF1 (10.7%), BAP1 (9.6%), BRCA1 (9.6%), BRCA2 (6.7%), and CHEK2 (5.3%; Fig 1B). For genes with more than 10 potentially germline mutations, we verified whether variants clustered in a specific region. For BRCA1, BRCA2, ATM, NF1, and BAP1, the variants were scattered over the multiple exons (data not shown). By contrast, for MUTYH and CHEK2, the variants clustered in specific hotspot regions (Data Supplement, Fig S1).

Potentially germline variants were mostly detected in lung (22.3% of patients carrying potentially germline variants), biliary tract (21.8% of patients), pancreatic (13% of patients), gastric (8.8%), breast (7.3%), and colorectal cancers (6.7%; Fig 1C). However, this distribution was largely influenced by differences in the number of patients with different histology enrolled in this study. In fact, the fraction of patients carrying potential germline mutations was higher in ovarian, breast, gastric, and pancreatic cancers, followed by biliary tract, colorectal, and lung cancers (Data Supplement, Fig S2 and Table S2).


The cohort of patients with potentially germline variants was younger than patients without potentially germline variants ( $P = .0033$ ; Table 2). Males were more numerous than females, although the difference was not statistically significant. Median TMB was significantly higher in patients with potentially germline variants compared with patients without potentially germline variants ( $P = .0001$ ). No statistical differences were observed between the cohort of patients with or without germline variants in terms of familial history or cancer, although the information was available for only 50% of patients (Table 2).

The majority of potentially germline mutations was detected by tissue testing (154/193; 79.8%), compared with ctDNA testing (39/193; 20.2%).

### Classification of Potentially Germline Variants Into CSG Classes

We categorized the 225 potentially germline variants into CSG classes, accordingly with the ESMO classification<sup>11</sup> (Data Supplement, Table S1). Collectively, 62/225 (27.5%) variants were detected in genes classified as most actionable (MA)-CSG class, 53/225 (23.6%) in genes belonging to the high actionable (HA)-CSG class, and 110/225 (48.9%) in the standard actionable (SA)-CSG class.

Sixty-two potentially germline mutations (26 SNVs and 36 indels) were detected in seven MA-CSGs in 59 patients (Fig 2A). Potentially germline variants were most frequently detected in BRCA1 (22/62 mutations, 35.5%), BRCA2 (15/62 mutations, 24.2%), and MSH6 (8/62 mutations, 12.9%; Fig 2A). The higher number of potentially germline variants belonging to the MA-CSG class was detected in non-small cell lung cancer (11/59, 18.6% of patients with MA-CSG class variants), breast cancer (10/59, 16.9%) and biliary tract cancer (9/59, 15.2%; Fig 2A and Data Supplement, Figs S3A



**FIG 1.** PGVs in the cohort of patients enrolled in the RATIONAL study. (A) Flowchart of the study. A total of 1,339 patients from the RATIONAL study were included in the analysis. Filters on the basis of the ESMO recommendations applied for identifying patients with potentially germline genomic alterations were shown. The percentage and the number of patients at each filtering step were indicated. <sup>a</sup>Eleven patients had both potentially germline PGVs and LRs. (B) Two hundred and twenty-five potentially germline genomic variants were identified in 40 CSGs recommended by ESMO for germline analysis and follow-up. (C) Distribution of patients (n = 193) with potentially germline variants accordingly to tumor types. Other: anal cancer (n = 1 patient); adrenal cancer (n = 1 patient); liver cancer (n = 1 patient); melanoma (n = 1 patient); and skin cancer (n = 1 patient). (continued on following page)

**FIG 1.** (Continued). CGP, comprehensive genomic profiling; CSG, cancer susceptibility gene; ESMO, European Society of Medical Oncology; FMI, Foundation Medicine; LR, large rearrangement; P/LP, pathogenic/likely pathogenic; PGV, potentially germline P/LP variant; SNV, single-nucleotide variant; VAF, variant allelic frequency.

and S3B). However, breast cancer was the tumor with the highest probability to detect MA-CSG class variants, that were found in 10/54 (18.5%) patients with breast cancer (Data Supplement, Figs S3A and S3B).

Familial history of cancer was reported for 12/59 (20.3%) patients, including four patients with breast cancer, three patients with pancreatic cancer, three patients with ovarian cancer, one patient with gastric cancer, and one patient with cholangiocarcinoma. For 21/59 (35.6%) patients, no familial history of tumor was reported. For 26/59 (44.1%) patients, the information was not available (Data Supplement, Table S3).

We classified 45 potentially germline mutations (39 SNVs and 14 indels) in nine genes in the HA-CSG class for which germline analysis and follow-up is suggested independently from age, and additional eight SNVs in three genes with follow-up recommended only for patients with tumors arising at age <30 years (Fig 2B). Overall, the 53 potentially germline variants were identified in 48 patients, including six patients with age <30 years. The majority of potentially germline mutations were observed in the *MUTYH* gene (24/53 variants; 45.3%) in 23 patients (Fig 2B). *MUTYH* variants were the only mutation in CSGs in 21 patients, whereas two patients carried also a *BRCA1* and a *PALB2* variant, respectively. Genomic variants were more frequently observed in pancreatic cancer (14/48, 29.2% of patients with HA-CSG class variants; 14/165, 8.5% of patients with pancreatic cancer), followed by colorectal and lung cancers (Fig 2B and

Data Supplement, Figs S3A and S3C). Variants in genes belonging to the HA-CSG class were also detected in different tumors at low frequency (Fig 2B and Data Supplement, Figs S3A and S3C). Familial history of cancer was reported in 12/48 (25%) patients (Data Supplement, Table S3).

Eight SNVs in *APC* (n = 1), *PTEN* (n = 1), and *TP53* (n = 6) genes were detected in 6/26 (23.1%) patients with age <30 years (Fig 2B). All patients had a *TP53* mutation. Two patients had two co-occurring variants in HA-CSG genes, and 3/6 (50%) had at least one potentially germline variant belonging to the other classes (MA/SA). Two patients had breast cancer, two patients had CNS cancer, one patient had a GI stromal tumor, and one patient had colorectal cancer (Data Supplement, Fig S3C). Two of six (33.3%) patients reported familial history of tumor, whereas 2/6 (33.3%) did not. For two patients, the information was not available (Data Supplement, Table S3).


Finally, we categorized 110 mutations in 11 genes (51 Indels and 59 SNVs) in 103 patients in the SA-CSGs class (Fig 2C). *ATM* (34/110, 30.9%), *NF1* (24/110, 21.8%), and *BAP1* (22/110, 20%) were the most common mutated genes (Fig 2C).

Mutations belonging to the SA-CSG class were most frequent in biliary tract cancer (30/103, 29.1% of patients with SA-CSGs class variants; 30/271, 11.1% of the biliary tract cancer cohort). SA-CSG class variants were found in 27/342 (7.9%) patients with lung cancer and 11/107 (10.3%) patients with esophagogastric cancer (Fig 2C and Data Supplement, Figs

**TABLE 2.** Clinicogenomic Characteristics of Patients With or Without Potentially Germline Variants

| Characteristic                     | All (N = 1,339)  | No PGVs (n = 1,146) | PGVs (n = 193)   | P       |
|------------------------------------|------------------|---------------------|------------------|---------|
| Age, years, median (range)         | 62 (18-89)       | 63 (18-89)          | 59 (18-82)       | .0033   |
| Patients with age <30 years        |                  | 21                  | 10               | .3547   |
| Sex, No. (%)                       |                  |                     |                  | .536376 |
| Male                               | 715 (53.4)       | 616 (53.7)          | 99 (51.3)        |         |
| Female                             | 624 (46.6)       | 530 (46.3)          | 94 (48.7)        |         |
| Tumor mutational burden, No. (%)   |                  |                     |                  | .333081 |
| Determined                         | 1,225 (91.5)     | 1,045 (91.2)        | 180 (93.3)       |         |
| Not determined                     | 114 (8.5)        | 101 (8.8)           | 13 (6.7)         |         |
| Median (range)                     | 4 Mut/Mb (0-216) | 3 Mut/Mb (0-100)    | 4 Mut/Mb (0-216) | .0001   |
| Family history for cancer, No. (%) |                  |                     |                  | .3764   |
| Yes                                | 262 (19.6)       | 223 (19.4)          | 39 (20.2)        |         |
| No                                 | 389 (29.05)      | 341 (29.7)          | 48 (24.9)        |         |
| NA                                 | 688 (51.4)       | 582 (50.8)          | 106 (54.9)       |         |

Abbreviations: NA, not applicable; PGV, potentially germline P/LP variant.



**FIG 2.** Categorization of potentially germline variants in CSG classes. (A) Potentially germline variants ( $n = 62$ ) in genes classified in the MA-CSG class. Distribution of 59 patients with variants classified in the MA-CSG class across the tumor types. (B) Potentially germline variants ( $n = 53$ ) in genes classified in the HA-CSG class. Distribution of 48 patients with variants classified in the HA-CSG class across the tumor types. (C) Potentially germline variants ( $n = 110$ ) in genes classified in the SA-CSG class. Distribution of 103 patients with variants classified in the SA-CSG class across the tumor types. Other: anal cancer ( $n = 1$  patient); adrenal cancer ( $n = 1$  patient); liver cancer ( $n = 1$  patient); CNS cancer ( $n = 1$  patient); melanoma ( $n = 1$  patient); ovarian cancer ( $n = 1$  patient); and thyroid cancer ( $n = 1$  patient). (D) LRs detected in the 16/40 CSGs in patients enrolled in the RATIONAL study. Distribution of patients with LRs across the tumor types. CSG, cancer susceptibility gene; HA, high actionable; LR, large rearrangement; MA, most actionable; SA, standard actionable.

S3A and S3D). Familial history of cancer was reported in 19/103 (18.4%) patients (Data Supplement, Table S3).

### Analysis of LRs in CSGs

CGP allows for the detection of LRs, including deletions of the whole or partial gene, duplications, triplications, and inversions, in addition to SNVs and indels. In the whole cohort of patients, 58 LRs in 16 CSGs were detected, including 29 gene rearrangements (duplications, triplications, and inversions), 10 deletions of the whole gene deletions, and 19 partial gene deletions, in 53/1,339 (3.95%) patients. GAs in *NF1* were most frequently observed (12/58; 20.7%), followed by alterations in *BRCA2* (10/58, 17.2%) and *BRCA1* (8/58, 13.8%; Fig 2D).

The majority of LRs were detected in genes classified in the SA-CSG class (28/58; 48.3%), followed by genes in the MA-CSG class (26/58; 44.8%; Data Supplement, Fig S4). Among 53 patients with LRs, 33.9% of patients had familial history of tumor, 22.6% of patients did not, whereas for 43.4% of patients, the information was not available. Patients with pancreatic cancer had the higher number of potentially germline LRs (8/53, 15.1% of patients with LRs; 8/165, 4.8% of patients with pancreatic cancer), followed by patients with biliary tract cancer (7/53, 13.2% of patients with LRs; 8/271, 2.9% of patients with biliary tract cancer; Fig 2D). Eight patients (8/53; 15.1%) had concomitant potentially germline variants in genes belonging to other CSG classes. The majority of LRs were identified by tumor tissue testing (37/53; 69.8%).

### Patients Referred for Genetic Counseling

The 225 potentially germline variants were identified in patients enrolled in 26 centers. Only 13/26 (50%) centers were able to provide data on germline testing and follow-up for 99/193 (51.3%) patients (Fig 3). Only 4/99 (4%) patients were referred to genetic counseling: two patients with biliary tract cancer in whom CGP detected variants in two genes of the MA-CSG class (*MLH1* and *BRCA1*), one patient with gastric cancer and familial history of tumor with a *NF1* variant and one patient with colorectal cancer carrying an *ATM* mutation.

Among the 95/99 (96%) patients not referred to oncogenetic consultation and follow-up, 30 carried variants in the MA-CSG class. Poor conditions or death within 3 months of report issuing (18/95; 18.9%), genetic counseling before the test (5/95; 5.3%), the lack of a geneticist in the center (5/95; 5.3%), and the lack of follow-up (5/95; 5.3%) are the reasons for which germline analysis and follow-up was not performed. However, reasons were not specified for 61/95 patients (64.2%; Fig 3).

## DISCUSSION


Tumor sequencing using large panels is rapidly becoming an integral part of care of patients with metastatic cancers for

identifying actionable mutations with therapeutic relevance. However, tumor-only sequencing also allows the identification of potentially germline variants associated with increased cancer risk.<sup>8,9,13-16</sup> Although these findings may not have implications for the treatment of the patient undergoing testing, the identification of potentially germline variants has extreme relevance for family members. Identifying a genetic risk of cancer allows the carrier to follow health surveillance programs and possibly undergo interventions for risk reduction.<sup>17,18</sup>

Obviously, there is a risk of the overinterpretation of tumor sequencing results. For this reason, ESMO has released recommendations to limit the impact of tumor-only findings, defining both the genes of interest and a VAF threshold associated with the probability that the identified variants may actually be germline.<sup>11</sup> When applying these rules, we found that about 14% of patients with cancer who received CGP had putative germline variants. Studies of tumor-only sequencing reported the presence of potentially germline variants in percentages of patients ranging from 3.6% to 40%.<sup>11,13-15,19,20</sup> Different factors, including the patients tested, the genes selected and the filters applied, and the approaches used for sequencing, may have a relevant impact on the rates of patients with potential germline variants. In this regard, ESMO guidelines recommends germline follow-up of *MUTYH* only for biallelic pathogenic variants. The inclusion of all *MUTYH* variants in our study may lead to overestimate the rate of patients with variants in CSGs. However, recent studies suggested that the presence of monoallelic *MUTYH* variants associated with somatic loss of heterozygosity might increase the susceptibility to some cancer types.<sup>21</sup> In addition, the American College of Medical Genetics and Genomics guidelines acknowledge that some *CHEK2* missense variants, including the I157T, should not be routinely reported on tumor or germline tests.<sup>22</sup> In our study, we found the *CHEK2* I157T missense variant in seven patients, one of whom also had a *BRCA2* variant. Germline follow-up is not recommended in carriers of this variant and, more generally, the management of *CHEK2* heterozygotes should be guided by personalized risk estimates.

Only two studies reported data obtained from the same CGP tests used in our study.<sup>14,20</sup> A study in 710 Japanese patients with advanced cancers reported a 40% rate of potential germline findings, whereas in a large cohort (n = 125,128) of patients with cancer, potential germline variants were detected in 9.7% of patients. However, in the first study, potentially germline variants in 46 recommended CSGs were described,<sup>20</sup> whereas in the latter manuscript, only 24 CSGs were considered using as cutoff VAFs > 10% for tissue CGP and >30% for liquid CGP. The high variability in the frequency of germline variants is likely due to the heterogeneous criteria adopted in these studies.

In agreement with previous findings,<sup>14</sup> the detection rate of GAs in CSGs was higher in patients tested on tissue compared with ctDNA. We do not expect that tissue and ctDNA testing



**FIG 3.** Diagram of referrals for germline testing of patients (n = 193) with potentially germline variants identified by tumor-only sequencing.

might show difference for the detection of germline variants. However, in cases with very low circulating cell-free DNA input, the sensitivity of the test could be affected. In addition, we noted that the majority of patients with breast (74%) and ovarian (86%) cancers were tested on tissue. Since these tumors have the highest frequency of mutations in CSGs, this imbalance may have contributed to the reported differences.

Alterations in CSGs, including MA-CSG group genes, have been identified both in tumors that are routinely subjected to genetic testing, such as ovarian and breast cancers, and in tumors that do not usually undergo genetic surveillance, such as cancer of unknown primary, cholangiocarcinoma, and lung cancer. Although our data are incomplete and may weaken our conclusions, many patients with CSG alterations did not have a family history of cancer, and therefore would never have undergone genetic testing. This observation highlights the potential of CGP, which could allow the identification of families with a genetic risk of developing cancer.

The off-target identification of CSG alterations has also been described in previous studies and raises an important question about their possible pathogenic role.<sup>14,23</sup> There is no evidence to support a role of off-target mutations in CSGs, such as *BRCA* variants identified in lung cancer or cholangiocarcinoma, in the development of these tumors. However, this possible correlation is worth investigating to clarify any pathogenic mechanisms not yet identified.<sup>24</sup>

Most previous studies on CSGs have only analyzed SNVs and indels. However, inactivation of CSGs can also be determined by gene rearrangements.<sup>25-27</sup> LRs can be difficult to detect

and may be underreported as a cause for hereditary cancer risk. A previous study described that LRs account for approximately 7% of pathogenic variants identified by testing germinal DNA with a pan-cancer NGS panel.<sup>28</sup> Our study shows that LRs in CSGs can be detected by tumor tissue testing at a frequency of almost 4% in patients with cancer. Importantly, 44.8% of LRs occurred in genes of the MA-CSG class, thus confirming the relevance of detecting and reporting LRs.

The identification of individuals with a genetic risk of developing cancer could represent an important added value of the CGP.<sup>29,30</sup> However, we found that the majority of Italian oncologists are not trained to adequately manage this information. Only 4% of patients with potentially germline GAs were referred to genetic counseling, highlighting a significant lack of awareness of the relevance of this information among the Italian oncology community. The inclusion of *MUTYH* and low-risk *CHECK2* variants in our study might lead to overestimate this phenomenon. The ESMO recommendations indicate that genetic counseling is strongly recommended for all patients with alterations of the MA-CSG class, leaving the decision for the other classes to the oncologist on the basis of the organization of the national health system.<sup>11</sup> However, even considering only patients with MA-CSG alterations, 93.75% of patients did not receive genetic counseling. The RATIONAL study provides a snapshot of the Italian clinical practice, providing recommendations on the interpretation of the test result, but leaving management at the local level.<sup>31</sup> Our findings suggest that the results of CGP tests should always be discussed within molecular tumor boards or within multidisciplinary teams, where all the professional skills needed to interpret complex genomic data are available, including molecular

biologists and geneticists. Several evidences suggest that oncologists not only in Italy are trained in recognizing actionable mutations for therapeutic intervention and generally have a low ability to interpret complex genomic analyses.<sup>32</sup> In this respect, structured pathways for the management of germinal findings to support oncologists are not present in the majority of Italian oncology centers. Automated pipelines for predicting somatic versus germline origin and homozygous versus heterozygous or subclonal state of variants identified by tumor-only CGP have been recently developed that might help physicians by highlighting potentially germline variants.<sup>33</sup> A specific

training of the oncologists to manage the complex implications of potentially genomic findings and a close collaboration with geneticists and molecular biologists are definitely required.

In conclusion, the data from the RATIONAL study confirm the utility of CGP also for the identification of alterations in CSGs and highlight the importance of reporting and paying attention to LRs in addition to SNVs and Indels. However, our findings also demonstrate a relative lack of knowledge of the implications of CGP tests among oncologists and underline the need for specific training in this area.

## AFFILIATIONS

<sup>1</sup>Cell Biology and Biotherapy Unit, Istituto Nazionale Tumori IRCCS Fondazione G. Pascale, Napoli, Italy

<sup>2</sup>Thoracic Department, Istituto Nazionale Tumori IRCCS Fondazione G. Pascale, Napoli, Italy

<sup>3</sup>U.O.C. Oncology, Azienda Ospedaliera Universitaria Integrata, University and Hospital Trust of Verona, Verona, Italy

<sup>4</sup>Medical Oncology Unit, ASLCN2 Alba e Bra, Michele e Pietro Ferrero Hospital, Verduno, Italy

<sup>5</sup>Department of Surgery, Oncology and Gastroenterology, University of Padova, Padova, Italy

<sup>6</sup>Oncology 2, Istituto Oncologico Veneto (IOV) IRCCS, Padova, Italy

<sup>7</sup>Department of Oncology, Azienda Sanitaria Universitaria Friuli Centrale (ASUFC), Udine, Italy

<sup>8</sup>Department of Medical Oncology, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) “Dino Amadori”, Meldola, Italy

<sup>9</sup>Department of Oncology, Università Politecnica delle Marche, AOU delle Marche, Ancona, Italy

<sup>10</sup>Medical Oncology, Humanitas Gradenigo Hospital, Torino, Italy

<sup>11</sup>U.O. Medical Oncology 2, IRCCS Ospedale Policlinico San Martino, Genova, Italy

<sup>12</sup>Medical Oncology, Comprehensive Cancer Centre IRCCS—AUSL Reggio Emilia, Reggio Emilia, Italy

<sup>13</sup>Oncology Unit, ASST Papa Giovanni XXIII, Bergamo, Italy

<sup>14</sup>Oncology Unit, Ospedale del Mare, Napoli, Italy

<sup>15</sup>Medical Oncology Unit, Fondazione IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo, Italy

<sup>16</sup>Department of Internal Medicine and Medical Therapy, University of Pavia, Pavia, Italy

<sup>17</sup>Comprehensive Cancer Center, IRCCS Policlinico San Matteo Foundation, Pavia, Italy

<sup>18</sup>School of Medicine and Department of Diagnostic Innovation, Fondazione IRCCS, Istituto Nazionale Tumori, University of Milan, Milano, Italy

<sup>19</sup>Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy

<sup>20</sup>Department of Medical Oncology, Santa Chiara Hospital—APSS, Trento, Italy

<sup>21</sup>Department of Experimental and Clinical Medicine, University of Florence, Firenze, Italy

<sup>22</sup>Oncology Unit, Careggi University Hospital, Firenze, Italy

<sup>23</sup>Department of Oncology, University of Turin, S. Luigi Gonzaga Hospital, Torino, Italy

<sup>24</sup>European Institute of Oncology, IRCCS, Milano, Italy

<sup>25</sup>Department of Oncology and Hemato-Oncology, University of Milano, Milano, Italy

<sup>26</sup>Scientific Directorate, IRCCS Istituto Romagnolo per lo Studio dei Tumori (IRST) “Dino Amadori”, Meldola, Italy

## CORRESPONDING AUTHOR

Nicola Normanno, MD; e-mail: Nicola.normanno@irst.emr.it.

## SUPPORT

Supported by unconditional grants from Roche, AstraZeneca, Bayer, BMS, Janssen, Takeda, Amgen, Daiichi Sankyo, Eli Lilly, and Novartis. A.D.L. was supported by grants of the Italian Minister of Health (Ricerca Corrente). N.N. was supported by the Italian Ministry of Health (Project T3-AN-06 “Sviluppo di una piattaforma per la implementazione clinica della oncologia di precisione nelle regioni del centro-sud Italia”).

## AUTHOR CONTRIBUTIONS

**Conception and design:** Riziero Esposito Abate, Carmine Pinto, Nicola Normanno, Antonella De Luca

**Provision of study materials or patients:** Michele Milella, Fabrizio Tabbò, Giacomo Pelizzari, Rossana Berardi, Lucio Buffoni, Tiziana P. Latiano, Salvatore Corallo, Giancarlo Pruneri, Orazio Caffo, Silvia Novello, Giuseppe Curigliano

**Collection and assembly of data:** Alessandro Morabito, Michele Milella, Fabrizio Tabbò, Valentina Guarneri, Giacomo Pelizzari, Ilario G. Rapposelli, Rossana Berardi, Lucio Buffoni, Elisa Bennicelli, Francesca Zanelli, Carlo Tondini, Laura Attademo, Tiziana P. Latiano, Salvatore Corallo, Giancarlo Pruneri, Federica Marmorino, Orazio Caffo, Lorenzo Antonuzzo, Simona Tessitore, Silvia Novello, Giuseppe Curigliano, Carmine Pinto, Nicola Normanno

**Data analysis and interpretation:** Riziero Esposito Abate, Carmine Pinto, Nicola Normanno, Antonella De Luca

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/po/author-center](http://ascopubs.org/po/author-center).

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://OpenPayments.org)).

**Alessandro Morabito**

**Consulting or Advisory Role:** MSD, Boehringer Ingelheim, Takeda  
**Speakers' Bureau:** AstraZeneca/MedImmune, Bristol Myers Squibb/Celgene, MSD, Novartis, Roche, Boehringer Ingelheim, Lilly

**Michele Milella**

**Honoraria:** MSD, AstraZeneca, Roche, Ipsen, Janssen Oncology  
**Consulting or Advisory Role:** Novartis  
**Research Funding:** Roche (Inst)

**Fabrizio Tabbò**

**Speakers' Bureau:** AstraZeneca, Roche, Novartis, Takeda, Pfizer, Pierre Fabre  
**Research Funding:** Roche

**Valentina Guarneri**

**Consulting or Advisory Role:** Lilly, Novartis, MSD, Gilead Sciences, Olema Oncology, Exact Sciences, Pierre Fabre, Daiichi Sankyo, AstraZeneca, Menarini Group, Roche, Pfizer  
**Speakers' Bureau:** Novartis, Lilly, AstraZeneca, Daiichi Sankyo, Menarini Stemline, Roche, Gilead Sciences, Pfizer  
**Research Funding:** AstraZeneca (Inst), Roche (Inst), Novartis (Inst), Merck (Inst), BMS (Inst), Lilly (Inst), Merck KGaA (Inst), Daiichi Sankyo/AstraZeneca (Inst), Nerviano Medical Sciences (Inst), GlaxoSmithKline (Inst), Synthon (Inst), Pfizer (Inst), Seagen (Inst)  
**Travel, Accommodations, Expenses:** Gilead/Forty Seven, AstraZeneca, Menarini Group, Roche

**Giacomo Pelizzari**

**Consulting or Advisory Role:** Amgen, Johnson & Johnson/Janssen, Roche, Novartis, Lilly, AstraZeneca, MSD  
**Speakers' Bureau:** Amgen, AstraZeneca  
**Travel, Accommodations, Expenses:** Sanofi, Roche, Novartis, Pfizer

**Rossana Berardi**

**Consulting or Advisory Role:** Lilly, Boehringer Ingelheim, Roche (Inst), GlaxoSmithKline, Eisai, Menarini, Bayer  
**Research Funding:** AstraZeneca (Inst), Pfizer (Inst), Roche (Inst)

**Elisa Bennicelli**

**Honoraria:** Servier

**Salvatore Corallo**

**Consulting or Advisory Role:** Daiichi Sankyo Europe GmbH, Merck Serono, AstraZeneca, Amgen  
**Travel, Accommodations, Expenses:** Amgen, Roche, Daiichi Sankyo Europe GmbH

**Giancarlo Pruneri**

**Honoraria:** Novartis, Roche, Genomic Health, AstraZeneca, Menarini, Exact Sciences  
**Consulting or Advisory Role:** ADS Biotec  
**Research Funding:** Roche, Roche Molecular Diagnostics

**Orazio Caffo**

**Honoraria:** Astellas Pharma, Bayer, AstraZeneca, Pfizer, Ipsen, Recordati  
**Consulting or Advisory Role:** Astellas Medivation, MSD Oncology, Advanced Accelerator Applications, Bayer, Pfizer, Novartis

**Lorenzo Antonuzzo**

**Honoraria:** Amgen, Roche, AstraZeneca, Novartis  
**Consulting or Advisory Role:** Merck Serono, MSD Oncology

**Silvia Novello**

**Honoraria:** BMS GmbH & Co KG, Lilly, Takeda, Roche, Pfizer, AstraZeneca, Boehringer Ingelheim, MSD, AbbVie, Sanofi, PharmaMar, BeiGene, Accord Healthcare, Novartis  
**Consulting or Advisory Role:** AstraZeneca, Daiichi Sankyo Europe GmbH, Lilly, AbbVie, GlaxoSmithKline  
**Travel, Accommodations, Expenses:** Sanofi

**Giuseppe Curigliano**

**Leadership:** ESMO, European Society of Breast Cancer Specialists (EUSOMA = ESMO Open, ESMO)  
**Honoraria:** Ellipses Pharma  
**Consulting or Advisory Role:** Roche/Genentech, Pfizer, Novartis, Lilly, Foundation Medicine, Bristol Myers Squibb, Samsung, AstraZeneca, Daiichi Sankyo, Boehringer Ingelheim, GlaxoSmithKline, Seagen, Guardant Health, Veracyte, Celcuity, Hengrui Therapeutics, Menarini, Merck, Exact Sciences, Blueprint Medicines, Gilead Sciences  
**Speakers' Bureau:** Roche/Genentech, Novartis, Pfizer, Lilly, Foundation Medicine, Samsung, Daiichi Sankyo, Seagen, Menarini, Gilead Sciences, AstraZeneca, Exact Sciences  
**Research Funding:** Merck (Inst)  
**Travel, Accommodations, Expenses:** Roche/Genentech, Pfizer, Daiichi Sankyo, AstraZeneca

**Carmine Pinto**

**Consulting or Advisory Role:** Amgen, Astellas Pharma, AstraZeneca, Bristol Myers Squibb, Ipsen, Janssen, Incyte, Merck Serono, MSD, Novartis, Roche, Servier  
**Speakers' Bureau:** AstraZeneca, Bristol Myers Squibb, Ipsen, Janssen, Incyte, Merck Serono, MSD, Novartis, Pierre Fabre, Roche, Servier  
**Research Funding:** Daiichi Sankyo, AstraZeneca, Bayer, Bristol Myers Squibb, Lilly, Roche

**Nicola Normanno**

**Honoraria:** Thermo Fisher Scientific, Lilly, MSD, Illumina, Merck Serono, Incyte, Biocartis, AstraZeneca, Amgen, Servier, Novartis, SOPHiA Genetics  
**Consulting or Advisory Role:** Biocartis, AstraZeneca, Bayer, Incyte, Novartis, Roche  
**Research Funding:** AstraZeneca (Inst), Biocartis (Inst), Illumina (Inst), Incyte (Inst), Merck Serono (Inst), Qiagen (Inst), Roche (Inst), Thermo Fisher Scientific (Inst), SOPHiA Genetics (Inst)  
**Travel, Accommodations, Expenses:** Merck Serono

No other potential conflicts of interest were reported.

**ACKNOWLEDGMENT**

The authors thank Jaime Huayhua Ccanto and Salvatore Tufano for technical support and all clinicians and data managers of the medical and academic centers participating in the study.

**REFERENCES**

- Casolino R, Beer PA, Chakravarty D, et al: Interpreting and integrating genomic tests results in clinical cancer care: Overview and practical guidance. *CA Cancer J Clin* 74:264-285, 2024
- Mateo J, Steuten L, Aftimos P, et al: Delivering precision oncology to patients with cancer. *Nat Med* 28:658-665, 2022
- Normanno N, Apostolidis K, de Lorenzo F, et al: Cancer biomarkers in the era of precision oncology: Addressing the needs of patients and health systems. *Semin Cancer Biol* 84:293-301, 2022
- Chakravarty D, Solit DB: Clinical cancer genomic profiling. *Nat Rev Genet* 22:483-501, 2021
- Meric-Bernstam F, Brusco L, Shaw K, et al: Feasibility of large-scale genomic testing to facilitate enrollment onto genomically matched clinical trials. *J Clin Oncol* 33:2753-2762, 2015
- Pinto C, Biffoni M, Popoli P, et al: Molecular tests and target therapies in oncology: Recommendations from the Italian workshop. *Future Oncol* 17:3529-3539, 2021
- Mandelker D, Zhang L: The emerging significance of secondary germline testing in cancer genomics. *J Pathol* 244:610-615, 2018
- Terraf P, Pareja F, Brown DN, et al: Comprehensive assessment of germline pathogenic variant detection in tumor-only sequencing. *Ann Oncol* 33:426-433, 2022
- Raymond VM, Gray SW, Roychowdhury S, et al: Germline findings in tumor-only sequencing: Points to consider for clinicians and laboratories. *J Natl Cancer Inst* 108:djv351, 2016
- Stadler ZK, Schrader KA, Vijai J, et al: Cancer genomics and inherited risk. *J Clin Oncol* 32:687-698, 2014
- Kuzbari Z, Bandlamudi C, Loveday C, et al: Germline-focused analysis of tumour-detected variants in 49,264 cancer patients: ESMO Precision Medicine Working Group recommendations. *Ann Oncol* 34:215-227, 2023
- Mandelker D, Donoghue M, Talukdar S, et al: Germline-focussed analysis of tumour-only sequencing: Recommendations from the ESMO Precision Medicine Working Group. *Ann Oncol* 30:1221-1231, 2019

13. Clark DF, Maxwell KN, Powers J, et al: Identification and confirmation of potentially actionable germline mutations in tumor-only genomic sequencing. *JCO Precis Oncol* 3:1-11, 2019
  14. Tung N, Dougherty KC, Gatof ES, et al: Potential pathogenic germline variant reporting from tumor comprehensive genomic profiling complements classic approaches to germline testing. *NPJ Precis Oncol* 7:76, 2023
  15. Sanabria-Salas MC, Anggala NC, Gillies B, et al: Clinical integration of germline findings from a tumor testing precision medicine program. *BMC Cancer* 25:176, 2025
  16. Meric-Bernstam F, Brusco L, Daniels M, et al: Incidental germline variants in 1000 advanced cancers on a prospective somatic genomic profiling protocol. *Ann Oncol* 27:795-800, 2016
  17. Dullens B, de Putter R, Lambertini M, et al: Cancer surveillance in healthy carriers of germline pathogenic variants in BRCA1/2: A review of secondary prevention guidelines. *J Oncol* 2020:9873954, 2020
  18. McCarthy RL, Copson E, Tapper W, et al: Risk-reducing surgery for individuals with cancer-predisposing germline pathogenic variants and no personal cancer history: A review of current UK guidelines. *Br J Cancer* 129:383-392, 2023
  19. Bychkovsky BL, Li T, Sotelo J, et al: Identification and management of pathogenic variants in BRCA1, BRCA2, and PALB2 in a tumor-only genomic testing program. *Clin Cancer Res* 28:2349-2360, 2022
  20. Kawamura M, Shiota H, Niihori T, et al: Management of patients with presumed germline pathogenic variant from tumor-only genomic sequencing: A retrospective analysis at a single facility. *J Hum Genet* 68:399-408, 2023
  21. Paller CJ, Tukachinsky H, Maertens A, et al: Pan-cancer interrogation of MUTYH variants reveals biallelic inactivation and defective base excision repair across a spectrum of solid tumors. *JCO Precis Oncol* 8:e2300251, 2024
  22. Hanson H, Astiazaran-Symonds E, Amendola LM, et al: Management of individuals with germline pathogenic/likely pathogenic variants in CHEK2: A clinical practice resource of the American College of Medical Genetics and Genomics (ACMG). *Genet Med* 25:100870, 2023
  23. Vitale A, Mastrandoni L, Russo J, et al: Impact of comprehensive genome profiling on the management of advanced non-small cell lung cancer: Preliminary results from the lung cancer cohort of the FPG500 program. *JCO Precis Oncol* 8:e2400297, 2024
  24. Passiglia F, Righi L, Bironzo P, et al: Niraparib plus dostarlimab in pleural mesothelioma or non-small cell lung cancer harboring HRR mutations: Interim results of the UNITO-001 phase II prospective trial. *Clin Cancer Res* 30:959-964, 2024
  25. Sun Z, Bai C, Su M, et al: Comprehensive profiling of pathogenic germline large genomic rearrangements in a pan-cancer analysis. *Mol Oncol* 17:1917-1929, 2023
  26. Leite Rocha D, Ashton-Prolla P, Rosset C: Reviewing the occurrence of large genomic rearrangements in patients with inherited cancer predisposing syndromes: Importance of a comprehensive molecular diagnosis. *Expert Rev Mol Diagn* 22:319-346, 2022
  27. Baudhuin LM, Ferber MJ, Winters JL, et al: Characterization of hMLH1 and hMSH2 gene dosage alterations in Lynch syndrome patients. *Gastroenterology* 129:846-854, 2005
  28. Mancini-DiNardo D, Judkins T, Kidd J, et al: Detection of large rearrangements in a hereditary pan-cancer panel using next-generation sequencing. *BMC Med Genomics* 12:138, 2019
  29. Armon S, Hollander A, Segev Y, et al: Presymptomatic awareness of BRCA1/BRCA2 status and outcomes in women with ovarian cancer. *JAMA Netw Open* 8:e251435, 2025
  30. Lambertini M, Blondeaux E, Tomasello LM, et al: Clinical behavior of breast cancer in young BRCA carriers and pre-diagnostic awareness of germline BRCA status. *J Clin Oncol* 43:1706-1719, 2025
  31. Normanno N, De Luca A, Abate RE, et al: Current practice of genomic profiling of patients with advanced solid tumours in Italy: The Italian Register of Actionable Mutations (RATIONAL) study. *Eur J Cancer* 187:174-184, 2023
  32. de Moor JS, Gray SW, Mitchell SA, et al: Oncologist confidence in genomic testing and implications for using multimarker tumor panel tests in practice. *JCO Precis Oncol* 4:620-631, 2020
  33. Sun JX, He Y, Sanford E, et al: A computational approach to distinguish somatic vs. germline origin of genomic alterations from deep sequencing of cancer specimens without a matched normal. *PLoS Comput Biol* 14:e1005965, 2018
-