



RESEARCH ARTICLE

10.1002/2014WR016555

The role of stratification on lakes' thermal response: The case of Lake Superior

Sebastiano Piccolroaz¹, Marco Toffolon¹, and Bruno Majone¹

¹Department of Civil, Environmental, and Mechanical Engineering, University of Trento, Trento, Italy

Key Points:

- Prediction of lake surface temperature (LST) using the lumped model *air2water*
- The feedback between LST and stratification is investigated and discussed
- Investigation of the exceptional warming of Lake Superior in summer 1998

Supporting Information:

- Supporting Information S1

Correspondence to:

S. Piccolroaz,
s.piccolroaz@unitn.it

Citation:

Piccolroaz, S., M. Toffolon, and B. Majone (2015), The role of stratification on lakes' thermal response: The case of Lake Superior, *Water Resour. Res.*, 51, 7878–7894, doi:10.1002/2014WR016555.

Received 14 OCT 2014

Accepted 6 SEP 2015

Accepted article online 10 SEP 2015

Published online 3 OCT 2015

Abstract During the last several decades, the Great Lakes region has been experiencing a significant rise in temperatures, with the extraordinary summer warming that affected Lake Superior in 1998 as an example of the marked response of the lake to increasingly warmer atmospheric conditions. In this work, we combine the analysis of this exceptional event with some synthetic scenarios, to achieve a deeper understanding of the main processes driving the thermal dynamics of surface water temperature in Lake Superior. The analysis is performed by means of the lumped model *air2water*, which simulates lake surface temperature as a function of air temperature alone. The model provides information about the seasonal stratification dynamics, suggesting that unusual warming events can result from two factors: anomalously high summer air temperatures, and increased strength of stratification resulting from a warm spring. The relative contribution of the two factors is quantified using the model by means of synthetic scenarios, which provide a simple but effective description of the positive feedback between the thermal behavior and the stratification dynamics of the lake.

1. Introduction

Recent studies have demonstrated that lakes are highly sensitive to changes in environmental conditions, thus representing a valuable proxy to evaluate the effects of a changing climate [Quayle *et al.*, 2002; Adrian *et al.*, 2009; Williamson *et al.*, 2009]. As a consequence, many climate change studies have focused on long-term trends observed in lakes, with particular emphasis on the analysis of water temperature dynamics of large lakes [e.g., Livingstone, 2003; Verburg *et al.*, 2003; Vollmer *et al.*, 2005; Coats *et al.*, 2006; Hampton *et al.*, 2008]. Observational evidences revealed that inland water bodies are rapidly warming throughout the world, with Lake Surface water Temperature (LST) increasing at rates up to an order of magnitude higher than those found for the global ocean [Schneider *et al.*, 2009]. Furthermore, it has been shown that lakes warming is in some case larger than that observed for the surrounding air temperature [Austin and Colman, 2007; Lenters *et al.*, 2012], especially at midlatitudes including North America and North Europe [Schneider and Hook, 2010].

Of particular interest is the warming trend that the Laurentian Great Lakes have been experiencing in the last century [e.g., McCormick and Fahnenstiel, 1999]. For example, by using a 100 year long-time series of water temperature measured at Lake Superior outlet as a proxy of the offshore LST, Austin and Colman [2008] estimated a mean rate of warming of about $0.027^{\circ}\text{C yr}^{-1}$ over the last century (i.e., data covering the period 1906–2005), with a dramatic increase up to $0.11^{\circ}\text{C yr}^{-1}$ taking place starting in the 1980s. This recent trend has been also confirmed by analyzing in situ measurements from offshore buoys [Austin and Colman, 2007]. Notice that these values refer to the 3 month summer period July, August, and September (JAS), which are the months with the largest LST increases [Lenters, 2004; Austin and Colman, 2008]. During the last several decades, similar summer warming trends have been also observed for other lakes in North America, as is the case of Lake Michigan and Lake Huron [Austin and Colman, 2007], and some smaller lakes between California and Nevada [Schneider *et al.*, 2009].

In 1998, uncommonly high air temperatures throughout the year determined an exceptional warming of Lake Superior, with summer LST difference between 1998 and 1997 being higher than the corresponding air temperature difference. This extraordinary event can be explained as a combined consequence of a particularly mild, nearly ice-free winter in part due to a significantly strong El Niño event [Assel *et al.*, 2000; Van Cleave *et al.*, 2014], and an anomalously warm summer season [Austin and Colman, 2007]. More generally, it




Figure 1. Lake Superior with the location of the air temperature station (STD4—Stannard Rock) and the moored thermistor chain [Titze and Austin, 2014] used in this work.

can be seen as a remarkable example of the amplified response of a lake to increasing air temperature, and developing a phenomenological understanding is of paramount importance for a comprehensive description of lake behavior under evolving climate conditions. In addition, as water temperature plays a primary role in controlling a wide range of geochemical and ecological processes, an in-depth analysis of LST dynamics can also provide significant indirect information concerning possible influences on lake water quality and ecosystem functioning [e.g., Wetzel, 2001; Winder and Sommer, 2012; De Senerpont Domis et al., 2013]. This is even more relevant considering that an enduring and rapid rising of Lake Superior water temperature could substantially affect the lake ecosystem [e.g., Magnuson et al., 1997; Cline et al., 2013].

Recognizing the main processes affecting the thermal dynamics of LST during an exceptional event like the 1998 summer warming of Lake Superior can provide a deeper understanding of the thermal response in more general contexts. A fundamental question, in this perspective, concerns the contribution to LST variation which is attributable to the increased heat flux in the summer period as opposed to the influence of previous conditions of the lake. To address such an issue, we exploited *air2water* [Piccolroaz et al., 2013; Toffolon et al., 2014; Piccolroaz and Toffolon, 2015], a simple open-source model that simulates LST relying solely on air temperature, to isolate the relative contributions of two factors: external forcing and thermal structure of the lake. After introducing the model in the next section, we show that these factors are sufficient to suitably reproduce the LST dynamics, and then we analyze some synthetic scenarios to isolate their relative importance. Wider implications of the results and conclusions are finally presented in the last two sections.

2. Material and Methods

2.1. Study Site and Available Data

Lake Superior (Figure 1) is the largest of the five Great Lakes of North America, and the largest freshwater lake on Earth by surface area (surface area 82,103 km²; volume: 12,000 km³; maximum depth: 406 m). Long-term data of air and surface water temperature are available and are freely distributed by the National Oceanic and Atmospheric Administration (NOAA).

Two different sources of data have been used in this work: daily in situ measurements of air temperature provided by the National Oceanic and Atmospheric Administration's (NOAA), National Data Buoy Center (NDBC, webpage: <http://www.ndbc.noaa.gov/>), and daily LST retrieved from satellite imagery provided by NOAA Great Lakes Environmental Research Laboratory (GLERL, webpage: <http://www.glerl.noaa.gov/>). Air

temperature data adopted in this study are measured at the STDM4—Stannard Rock station, which belongs to the NOAA NDBC Coastal Marine Automated Network (C-MAN) and is installed at about 35 m above the lake level on a lighthouse located in the South-Eastern part of the lake (see Figure 1). The values of LST freely downloadable by the GLERL website refer to satellite-derived LST averaged over the whole lake, do not present significant gaps, and provide information during the entire year. Local LST values of the GLERL data set are in overall good agreement with LST measured by offshore buoys (NDBC network) [Schwab *et al.*, 1999], making the first data set preferable compared to in situ NDBC measurements, which are characterized by systematic gaps during winter when devices are removed to prevent damage from ice. We remark that for the purpose of the present work, the availability of LST data during winter time is fundamental as they provide the information to investigate the role of ice in controlling the timing and intensity of LST warming in summer. Both air temperature and LST data have been downloaded for the period between 1994 and 2011.

2.2. *air2water*: A Simple Model to Predict LST

The analysis presented in this work has been performed by means of *air2water* [Piccolroaz *et al.*, 2013], a simple lumped model that allows for estimating LST using air temperature as the only meteorological forcing. *air2water* is derived from the volume-integrated heat equation applied to the upper layer of the lake

$$\rho c_p V_s \frac{dT_w}{dt} = A \Phi_{net}, \quad (1)$$

where ρ is water density, c_p is the specific heat capacity, V_s is the surface volume of water that is involved in the heat exchange with the atmosphere, T_w is LST, t is time, A is the surface area of the lake, and Φ_{net} is the net heat flux into the upper water volume (accounting for the main fluxes entering and exiting V_s : short-wave and longwave radiation, sensible, and latent heat fluxes). After introducing appropriate simplifications, which are summarized in Appendix A (for a thorough discussion we refer to Piccolroaz *et al.* [2013] and Toffolon *et al.* [2014]), the equations of the model in its full (eight parameters, from a_1 to a_8) version reads as follows:

$$\frac{dT_w}{dt} = \frac{1}{\delta} \left\{ a_1 + a_2 T_a - a_3 T_w + a_5 \cos \left[2\pi \left(\frac{t}{t_y} - a_6 \right) \right] \right\}, \quad (2)$$

$$\begin{cases} \delta = \exp \left(-\frac{T_w - T_h}{a_4} \right) & \text{for } T_w \geq T_h \\ \delta = \exp \left(-\frac{T_h - T_w}{a_7} \right) + \exp \left(-\frac{T_w}{a_8} \right) & \text{for } T_w < T_h \end{cases} \quad (3)$$

where T_a is air temperature, δ is a dimensionless number given by the ratio between the volume V_s of the surface layer introduced in equation (1) and a reference volume V_r , and T_h is a reference value of the deep water temperature, which is approximately 4°C for deep dimictic lakes. We note that V_s (hereafter referred to as the reactive volume) varies in time due to thermal stratification, while V_r is the maximum volume affected by the surface heat flux when the lake experiences the weakest stratification conditions. In our formulation, these two volumes and the surface heat flux are not estimated separately, and the parameter δ is used to implicitly account for temporary reduction or enhanced efficiency of the heat exchange with the atmosphere (for instance, in the case of ice cover), as will be discussed later on.

The model parameters a_1 – a_8 account for a series of different processes, are defined within a physically reasonable range of variation, and are obtained through calibration against LST measurements. The ordinary differential equation (2) is solved numerically using the Runge-Kutta fourth-order scheme, with a daily time step. In spite of the simple formulation and the limited number of parameters, *air2water* is able to satisfactorily capture seasonal variations and interannual dynamics in LST [see Piccolroaz *et al.*, 2013; Toffolon *et al.*, 2014, for application to different lakes], thus representing an appealing tool for both conceptual studies and real case analyses. The fact that the model is data-driven, while being physically based, allows for the direct acquisition of information about the studied system during the calibration phase, which is performed via an automatic optimization procedure. Besides predicting LST, the model also estimates the seasonal evolution of the upper volume of water affected by the surface heat budget, through the evaluation of the volume ratio δ (well-mixed $\delta \rightarrow 1$, stratified $\delta \rightarrow 0$, ice covered $\delta > 1$, Piccolroaz *et al.* [2013]). The model has




Figure 2. Comparison between air temperature T_a (observed) and lake surface temperature T_w (observed and simulated): (a) long-term trends of mean summer values (JAS) within the period 1994–2011; and (b) daily values in 1997 and 1998. LST is retrieved by satellite imagery (GLERL data set). For representation purposes, air temperature has been filtered with a 30 days moving average in Figure 2b.

been successfully applied using different sources of data (i.e., LST measured at buoys or retrieved from satellite) and considering different case studies [Piccolroaz et al., 2013; Toffolon et al., 2014].

3. Results

3.1. Model Performances

Model parameters have been calibrated considering the historical series of air temperature and LST available for the period 1994–2011. A general good agreement between measured and simulated series has been obtained, with a Kling Gupta Efficiency (KGE) index [Gupta et al., 2009] of 0.98 and a root-mean-square error of about $1.1\text{ }^\circ\text{C}$. We recall that KGE ranges from $-\infty$ to 1: the closer to 1, the better the model performances. The mean deviation between simulated and observed LST is equal to $-0.0004\text{ }^\circ\text{C}$, suggesting the absence of a significant bias. We notice that the performance achieved by *air2water* is fully comparable to that obtained with more complex process-based models [e.g., Thiery et al., 2014], which however require a significantly larger amount of input data.

Figure 2a shows air temperature and LST records for the summer season (averages over JAS). Detecting long-term trends by means of linear regressions is difficult and may lead to results that are not statistically significant. Anyway, it is interesting to note that a significant warming of about $0.098\text{ }^\circ\text{C yr}^{-1}$ ($R^2 = 0.17$, p value = 0.09) and $0.107\text{ }^\circ\text{C yr}^{-1}$ ($R^2 = 0.08$, p value = 0.27) can be observed for measured air temperature and LST, respectively. This result differs from previous analyses [Austin and Colman, 2007; Schneider and Hook, 2010], which found that LST increased faster than air temperature based on meteorological stations located within a 500 km radius from the lake. Notice that in the present work we refer to air temperature data retrieved at about 35 m above the lake surface from a single station at an offshore location (i.e., STDM4 C-MAN station), which is more representative of the real lake conditions, and at the same time is




Figure 3. Comparison between δ simulated by *air2water* (δ_{sim}) and estimated from the analysis of water temperature profiles measured at the mooring station indicated in Figure 1 (δ_{est}). For representation purposes, δ_{est} has been filtered with a 7 days moving average.

fully representative of the typical seasonal thermal pattern of the atmosphere. In fact, the marked time lag between the temporal variations of air and water temperatures is clearly demonstrated in Figure 2b for the two years 1997 and 1998. Figure 2 also shows the performance of the model, which behaves properly both in detecting the long-term LST trend ($0.100^{\circ}\text{C yr}^{-1}$, $R^2 = 0.13$, p value = 0.14) and the associated intra-annual and interannual variations. In particular, the two years have been characterized by substantial differences (Figure 2b): 1997 being particularly cold, especially in summer, and 1998 showing unusually warm temperatures during the whole year (as already discussed above). These marked differences are also evident in Figure 2a, where both JAS air temperature and JAS LST are positioned below and above the corresponding trend lines, respectively, for 1997 and 1998. In this perspective, this 2 year period constitutes an interesting example of interannual climate variability, which is worth being examined to characterize the general thermal dynamics of the lake.

In Figure 3, we show the capability of *air2water* to reliably reproduce the temporal evolution of the reactive surface volume. This is done by comparing the simulated dimensionless volume ratio (δ_{sim}) (see section 2.2) with an independent estimate of the same variable (δ_{est}) obtained from the analysis of water temperature profiles measured at a mooring station deployed in the central part of the lake [Titze and Austin, 2014]. Water temperature data cover the period 2008–2011 with hourly resolution, and are available at different depths, from surface down to 250 m depth. Following the same procedure adopted in Toffolon *et al.* [2014] for the case of Lake Constance, we used the measured vertical profiles of temperature to estimate the volume of the surface well-mixed layer, assuming that this is a reasonable approximation of the reactive volume V_s . We first identified the thickness of the surface well-mixed layer as the smallest depth where water temperature difference with respect to surface is lower than a threshold of 1°C . Then we converted this depth into the corresponding volume of water on the basis of the hypsometric curve of the lake. Finally, we calculated δ_{est} by normalization to a reference volume, V_r , here assumed to be the entire volume of the lake. Despite the temperature threshold and the reference volume are arbitrary and may influence the evaluation of δ_{est} , the overall comparison between simulated and estimated δ (Figure 3, see also the supporting information for the comparison with δ_{est} evaluated in terms of dimensionless depth ratio instead of dimensionless volume ratio) clearly shows that the simple parameterization (3) is able to correctly reproduce seasonal and interannual (see, e.g., the anticipated stratification in summer 2010) patterns of stratification, thus indicating its suitability to be used for the purposes of this work. In order to make the comparison between δ_{sim} and δ_{est} fair, we have calibrated the model locally using LST retrieved from the closest NDBC buoy (i.e., station 45001, located at 2 km from the mooring station, and for which a 27 year long data set is available during the period 1985–2011) instead of the lake-averaged, satellite-retrieved LST. We note that this longer LST series was required to achieve a more robust calibration of the model than the one that could be obtained relying only on the 4 year data measured at the thermistor chain. Model calibration over the 27 year period yielded $\text{KGE} = 0.94$, a root-mean-square error of 1.6°C , and mean deviation of -0.0086°C . The performances are not significantly affected by the absence of LST data in winter, when the buoy is removed to prevent ice damage, with efficiency indexes being almost equal to those obtained adopting GLERL satellite-based temperature as observational data. Vertical temperature profiles are not available in winter, as well, so the comparison between δ_{sim} and δ_{est} is limited only to the “warm” period. Finally, δ_{sim} in




Figure 4. Annual cycle of (a) observed T_a (thin line) and simulated T_w (thick line), and (b) simulated volume ratio δ , considering 1997, 1998, and the mean year 1994–2011. For representation purposes, all temperature series have been filtered with a 30 days moving average.

Figure 3 has been calculated through equation (3) with the same surface T_w used to evaluate δ_{est} , and the values of the parameters a_4 , a_7 , and a_8 derived from point calibration of the model.

Although δ_{sim} is defined as the dimensionless reactive volume, it implicitly accounts for possible increase or decrease of heat fluxes due to complex processes that are not explicitly included in the simple formulation of the model. For instance, this is the case of the insulating effect of ice cover that results in a fictitious larger reactive volume (i.e., $\delta > 1$ as discussed in Piccolroaz *et al.* [2013]), or the increased effective heat fluxes due to unstable atmospheric boundary layer in late summer [Blanken *et al.*, 2011; Lofgren and Zhu, 1999] that can be partially explained with the lower values of δ_{sim} compared to δ_{est} in Figure 3. These results indicate that *air2water* is able to accurately simulate thermal and stratification dynamics of the lake, without the need to introduce a complex description of the air-water interface processes based on the quantification of the single heat flux components. This should be seen as a major advantage of the proposed formulation rather than a limitation, as it is generally difficult to find complete data sets of all meteorological variables (e.g., solar radiation, cloudiness, humidity, etc.) covering long-term periods, compared to the relatively larger availability of air and water temperature measurements.

3.2. Assessing the Roles of Air Temperature and Stratification

The analysis of the heat balance equation is a good starting point for understanding the influence of stratification, in general, and for pointing out the significant differences between LST in 1997 and 1998 (Figure 2), as a particular case. Looking at equation (1), it is evident that the warming rate of the lake is directly proportional to Φ_{net} and inversely proportional to V_s . The lake warms faster if the heat flux is high and the volume that directly participates in the heat exchange with the atmosphere is small (i.e., the lake is stratified). This is a central feature of the thermal dynamics of lakes.

In order to understand the role of the main factors controlling LST, in Figure 4 we compare the model results obtained for 1997, 1998, and for the mean year (corresponding to the overall period 1994–2011).

Table 1. Synthetic Scenarios Obtained Combining Different Periods of Air Temperature and Choosing Different Procedures for Estimating the Volume Ratio δ

Scenario	T_a	δ
Test 1 (T_a 1998 and δ 1997)	1998	1997
Test 2 (T_a 1997 and δ 1998)	1997	1998
Test 3 (steeper T_a)	1997: Oct–Jun 1998: Jul–Aug	Model
Test 4 (flatter T_a)	1998: Oct–Jun 1997: Jul–Aug	Model
Test 5 (ensemble of 365 cases) (lasting warm winter)	$T_{my} + 2^\circ\text{C}$: DOY = 1,i ^a T_{my} : DOY = i + 1365	Model
Test 6 (ensemble of 365 cases) (earlier warm summer)	T_{my} : DOY = 1,i $T_{my} + 2^\circ\text{C}$: DOY = 1 + 1365	Model

^aDOY stands for day of the year.

The analysis of temperature as a function of the day of the year starting from 1 January (Figure 4a) shows again that 1997 and 1998 were, respectively, colder and warmer than the mean year both concerning air (thin lines) and water (thick lines) temperatures. Figure 4b shows the variation of the volume ratio δ calculated for the same three annual cycles. The nearly ice-free winter that occurred in 1998 is reproduced by the model with values of δ always ~ 1 (we recall that $\delta > 1$ accounts for the insulating effect due to the presence of ice). This is confirmed in Figure 4a, where simulated LST is shown to be consistently greater than 0°C in winter 1998 and did not drop beneath $\sim 2^\circ\text{C}$ from the end of January to the end of April (contrary to the mean year). Thus, direct thermal stratification in 1998 started about 30 days earlier (start of May) than in 1997 (start of June), and about 20 days earlier compared to the mean year (end of May). Moreover, a significantly faster increase of LST characterized the lake in summer 1998: starting from June (in 1997 during the same period the lake was still inversely stratified, i.e., $\text{LST} < 4^\circ\text{C}$) the lake reached a sufficiently strong thermal stratification that caused the surface mixed layer to get significantly shallower at mid-July (i.e., $\delta \ll 1$). Strong stratification conditions, much stronger than those usually characterizing the lake (i.e., $\delta_{1998} < \delta_{my}$, where the subscript “my” stands for mean year), lasted for about 2.5 months, until the start of October. Later on, stratification progressively weakened, but remaining always slightly stronger than in 1997 and in the mean year. As a consequence of the smaller surface volume in September 1998, the decrease of LST was faster in 1998 than in 1997.

As a whole, the combination of an exceptionally high air temperature and a significantly longer period of thermal stratification in summer 1998 resulted in a higher heat input to the lake acting on a smaller water volume, and eventually in the significant warming of LST. The onset of stratified conditions was anticipated, consistently with the observed tendency toward a longer ice-free season in North American lakes (earlier occurrence of ice-departure dates, see, e.g., Anderson et al. [1996]; Schindler et al. [1996]; McCormick and Fahnenstiel [1999]). This fact established a positive feedback to LST warming through an earlier reduction of the surface volume (lower values of δ) and hence a faster increase of LST, which in turn contributed to further decrease δ during summer months.

Aimed at understanding the feedback between net heat flux and thermal stratification and evaluating their distinct contributions to LST dynamics, we applied *air2water* under six different synthetic conditions, which are summarized in Table 1. In all cases, we considered the same model parameter set, as identified by calibration over the period 1994–2011. The six cases are described in the following paragraphs. In the analysis of the results, we always refer to JAS as summer period to be consistent with previous analyses [Austin and Colman, 2008; Van Cleave et al., 2014], although we will discuss the distinct role of the different months, and especially of July, on the average value.

Test 1 is obtained by combining air temperature series measured in 1998 and δ as simulated for 1997 (i.e., δ is not calculated by the model but externally imposed). The resulting annual cycle of LST is shown in Figure 5a. Although the external forcing (i.e., air temperature) is the same as in 1998, LST exhibits a significantly different behavior compared to that simulated for 1998, solely as a consequence of the different stratification conditions that have been imposed. The timing is similar to that simulated for 1997 (e.g., summer peak around the 20th of August) because it is controlled by the value of δ ; moreover, until the end of August LST is generally much lower than in 1998. The mean LST during JAS is about 1.7°C colder than in




Figure 5. Annual cycle of (a) observed T_a (thin line) and simulated T_w (thick line), and (b) volume ratio δ , considering 1997 and 1998 (dashed lines), and Test 1 and Test 2 (solid lines). The value of δ is imposed in the two tests (see Table 1). For representation purposes, all temperature series have been filtered with a 30 days moving average.

1998, with most of the difference occurring in July (almost 5°C, see Table 2). In winter, LST for Test 1 is similar to 1998, and the two series start to diverge only when the stratification begins to play a role. In fact, the lower temperatures in summer for Test 1 can be fully attributed to the slower warming caused by the larger volume of water involved in the heat exchanges, and hence to a weaker and postponed thermal stratification of the lake (i.e., larger values of δ) compared to 1998. Finally, Test 1 shows a slower cooling of LST in autumn with respect to 1998. This indicates a higher thermal inertia of the lake, coherently with the occurrence of larger values of δ , thus larger volumes of water participating to the heat exchanges with the atmosphere.

Test 2 is the reciprocal of Test 1 (see Table 1), being obtained by combining air temperature series measured in 1997 with δ simulated for 1998. The analysis of results, which are shown in Figure 5a and summarized in Table 2, leads to similar, and in some cases complementary, considerations as those made for Test 1: (i) the timing of LST is driven by δ , thus is consistent with that simulated for 1998; (ii) the mean LST during the JAS

Table 2. Simulated Annual Maximum, Mean Summer (JAS), and Mean Monthly (June, July, August, and September, Respectively) LST for Years 1997 and 1998, and for the Synthetic Scenarios Tests 1–4 Reported in Table 1

Period	T_w (°C)					
	1997	1998	Test 1	Test 2	Test 3	Test 4
Cases						
Maximum	15.17	19.14	17.61	17.25	18.76	16.55
JAS	11.97	16.19	14.48	13.61	14.58	13.63
Jul	8.65	15.50	10.76	12.86	10.62	12.67
Aug	14.02	17.69	16.81	14.77	17.55	14.88
Sep	13.28	15.35	15.73	13.10	15.35	13.30




Figure 6. Annual cycle of (a) reconstructed T_a (thin line, see Table 1) and simulated T_w (thick line), and (b) simulated volume ratio δ , considering 1997 and 1998 (dashed lines), and Test 3 and Test 4 (solid lines). The value of δ is calculated in all cases. For representation purposes, all temperature series have been filtered with a 30 days moving average.

period is about 1.6°C warmer than in 1997, but it is colder than in 1998 and Test 1; (iii) July is the month characterized by the largest difference with respect to thermal conditions in 1997; (iv) from June to September the mean monthly LST is always colder than in Test 1 with the exception of July, when the presence of a strong stratification (i.e., small values of δ) determines a faster warming of the lake; and (v) in autumn LST decreases faster than in 1997 due to the reduced thermal inertia of the system (lower values of δ).

The combined analysis of Test 1 (T_a 1998 and δ 1997) and Test 2 (T_a 1997 and δ 1998) provides interesting elements for an approximate quantification of the specific role that net heat flux and stratification play in controlling LST dynamics. Results in Table 2 suggest that the difference between mean summer (JAS) LST in 1998 and 1997 (i.e., about 4.2°C) is attributable to a warmer T_a for about 60% (the difference between mean summer LST in Test 1 and 1997 is about 2.5°C), and to a stronger stratification for the remaining 40% (the difference between mean summer LST in Test 2 and 1997 is about 1.6°C). We notice that this proportion is valid only for the comparison between 1997 and 1998, and may change in other cases due to the inherent interdependence of the two effects and the high nonlinearity of the processes involved. However, the results of this simple analysis provide a clear indication that air temperature and thermal stratification play a nearly balanced role in regulating LST behavior.

3.3. Assessing the Role of Seasonal Dynamics

Two additional synthetic tests have been built as combinations of T_a observed in 1997 and in 1998 (see Table 1), aimed at understanding the relative contribution of winter/spring and summer seasons to the 1998 summer warming. In particular, Test 3 has been defined assuming T_a from 1997 for the period January–June and October–December, and T_a from 1998 for the remaining period (i.e., JAS). Thus, this synthetic year is characterized by a warm summer and cold conditions during the rest of the year, with steeper rising and falling limbs during June and October, respectively (see Figure 6a). The second synthetic year (Test 4)

has been constructed as the opposite to Test 3, thus is characterized by a cold summer (T_a from 1997) and warm conditions during the rest of the year (T_a from 1998), with a flatter transition during later spring and early autumn (see Figure 6a). Unlike the previous tests, δ is calculated by the model rather than being externally imposed.

Results for Test 3 (Steeper T_a , Figure 6) show that water temperature remains low from January to June (when the forcing coincides with that observed in 1997), but the lake starts warming as soon as air temperature rises in summer, and eventually in August and September it reaches temperatures that coincide with those obtained in 1998 (see Table 2). This is a consequence of the fact that the late summer/early autumn period is characterized by a strong thermal stratification, thus very small values of δ , which determines a fast adaptation of LST to air temperature [Toffolon *et al.*, 2014], fully overcoming the low temperatures resulting from the colder winter months. On the contrary, the results of the second synthetic year (Test 4, Flatter T_a , Figure 6) shows that, although LST coincides with that obtained in 1998 from January to June because the external forcing is the same, it starts deviating from the trend observed in 1998 as soon as the colder summer begins. The abrupt warming of the lake is absent, as is suggested by the fact that water temperature does not reach the same maximum of 1998, not even that of the mean year (see Figure 4). Nevertheless, LST in summer is always higher than that obtained in 1997 (with the exception of September, when LST coincides in the two cases, see Table 2), coherently with the occurrence of lower (or at least equal) values of δ throughout the whole year.

The comparison between the mean summer values of LST reported in Table 2 shows that the lake in Test 3 (Steeper T_a) is about 2.6°C warmer than in 1997, which allows one to conclude that the exceptionally high air temperatures registered during the JAS period in 1998 alone contributed to about 60% of the whole LST difference between 1998 and 1997 (i.e., nearly 4.2°C). Analogously, from Test 4 (Flatter T_a) we can infer that the occurrence of a warm winter/spring period in 1998 alone explained about 40% (i.e., about 1.7°C) of the total mean summer LST difference between the two years. It is necessary to notice that although the two contributions sum to almost 100%, this is purely due to how scenarios have been constructed, and should not be taken as general rule.

The thermal response of lakes is strictly dependent on the sequence of meteorological and climatic conditions, thus making difficult to analyze the independent contributions of single periods of the year. This is especially true for deep lakes during nonstratified or weakly stratified conditions, when, thanks to a high thermal inertia (large volume of the surface layer, i.e., large δ in our model), the system is able to retain a historical memory of the past conditions, with a significant influence on the ensuing thermal dynamics. Conversely, during the stratification period, the response time of LST to changes in air temperature is shortened considerably due to the low thermal inertia [Toffolon *et al.*, 2014]. Hence, the adaptation of LST to air temperature is more rapid and its variation more intense, with the inertial effect being significantly reduced during strong stratification conditions.

The behavior discussed above is quantitatively analyzed in the last two synthetic tests (Figure 7), which rely on an ensemble of 365 air temperature cycles dynamically reconstructed on the basis of two reference years: a warm year and a cold year. The air temperature of the colder year is that of the mean year 1994–2011 (T_{my}), while the warmer year has been defined as $T_{my} + \Delta T_a$. In this case, the increment ΔT_a has been assumed equal to the average difference between 1998 and the mean year, which is approximately 2°C, so that the warm year can be considered as a good proxy of the conditions in 1998.

The 365 annual cycles of air temperature used in Test 5 have been obtained by combining the first part of the warm year with the remaining part of the cold year, progressively delaying by 1 day the transition from the warm to the cold conditions, from 1 January to 31 December (see Figure 7a for a schematic illustration). Test 5 is thus aimed at quantifying the effect that a lasting warm winter may produce on the annual evolution of LST. Test 6 is the opposite case of Test 5 (see Figure 7b) and is aimed at evaluating the possible effect of an earlier warm summer. Therefore, Tests 5 and 6, in addition to Tests 3 (Steeper T_a) and 4 (Flatter T_a), contribute to a comprehensive overview of the possible effects that different previous air temperature conditions, occurring at different times of the year, may have on LST dynamics through the year. In particular, we assessed the effects on the average LST of five target periods: June, July, August, September, and the summer period JAS. For each target period extending from time t_1 to time t_2 , we evaluated the difference ΔT_w between the period-averaged LST of the test and of the mean year, and we plotted it as a




Figure 7. Analysis of Test 5 and Test 6: (a) reconstructed T_a for Test 5 and (b) for Test 6, for a given day i ; (c–g) period-averaged LST difference (ΔT_w) between the test and the mean year as a function of i and for different target periods: June, July, August, September, and JAS (respectively, from subplot c–g). Vertical lines denote the limits (t_1 and t_2 in equation (4)) of the target periods, while numbers indicate the contribution (expressed as cumulative percentage) of different parts of the year to the total ΔT_w .

function of the i th day of the year when the transition between the two reference years takes place (Figures 7c–7g):

$$\Delta T_w|_{t_1}^{t_2}(i) = \frac{1}{(t_2 - t_1)} \left[\int_{t_1}^{t_2} T_w^{test(i)} dt - \int_{t_1}^{t_2} T_w^{my} dt \right], \quad (4)$$

where T_w^{my} is the LST of the mean year, and $T_w^{test(i)}$ is the LST calculated for each case i of the set of 365 annual cycles of air temperature constituting the two tests (Figures 7a and 7b). Continuous ascending and descending curves in Figures 7c–7g refer to Test 5 and Test 6, respectively. In order to correctly interpret the figure, we suggest to read the ascending curve from left to right (i.e., LST increases for a progressively lasting warm winter), and the descending curve from right to left (i.e., LST increases for a progressively

Table 3. Relative Contributions of Different Parts of the Year (Contributing Periods) to the Total Period-Averaged LST Difference (ΔT_w) Between the Synthetic Year of Tests 5 and 6 and the Mean Year, Considering Different Target Periods: June, July, August, September, and JAS^a

Target Period	Test	Contributing Period			Total
		Winter (JFM ^b)	Spring (AMJ ^c)	Target Period	
Jun	5	0.67 (30.4%)	1.05 ^d (48.1%)	0.47 (21.5%)	2.19
	6	0.88 (40.1%)	1.00 ^d (45.9%)	0.31 (14.0%)	
Jul	5	0.96 (22.2%)	2.10 (48.8%)	1.25 (29.0%)	4.31
	6	0.80 (18.7%)	2.37 (55.0%)	1.13 (26.3%)	
Aug	5	0.08 (5.1%)	0.07 (4.1%)	1.18 (73.0%)	1.62
	6	<0.01 (0.1%)	0.03 (2.0%)	1.18 (72.9%)	
Sep	5	0.00 (0.0%)	0.00 (0.0%)	0.96 (75.3%)	1.27
	6	0.00 (0.0%)	0.00 (0.0%)	0.95 (75.0%)	
JAS	5	0.34 (14.3%)	0.70 (29.4%)	1.34 (56.3%)	2.38
	6	0.26 (10.8%)	0.78 (32.9%)	1.34 (56.3%)	

^aRelative contributions are given in terms of both ΔT_w and percentage of ΔT_w , so that, for all pairs "test"-target period," the sum of the first is the total ΔT_w and the sum of the latter gives 100%.

^bJanuary, February, and March.

^cApril, May, and June.

^dFor the case of June spring months are only April and May (AM).

earlier warm summer). Table 3 supports Figure 7 presenting the total ΔT_w associated with each target period (i.e., the water temperature difference between the warm and the cold year averaged during the target period), as well as the relative contributions to the total ΔT_w (in terms of both ΔT_w and percentage of ΔT_w) of different parts of the year: winter, spring, and the target period itself. The same information is graphically presented in Figures 7c–7g in terms of cumulative contributions.

The analysis of the results highlights interesting differences between the five target periods, mainly due to the stratification conditions of the lake, which vary significantly during the year. Notice that the stratification during the warm year can be approximated by the evolution of δ during 1998 shown in Figure 4 (see also the following discussion). The mean LST in June and July (Figures 7c and 7d) is strongly affected by the conditions that the lake experienced during the first half of the year, as suggested by the steep slope of ΔT_w from January onward. During the first half of the year, in fact, the thermal inertia in both the cold (mean year) and warm (mean year + 2°C) years is always relatively large (i.e., not so small values of δ , see Figure 4), and the system keeps memory of the past conditions to the extent that the previous state controls the mean LST in June and July. The role played by previous lake conditions is clear: a lasting winter (Test 5) with warmer air temperature during the period January to June explains around 71% (nearly 22% in winter and 49% in spring, see Table 3) of the total simulated ΔT_w in July (Figure 7d). Similar considerations can be made for June, but the memory effect is even higher: warmer air temperature during the period January to May explains around 79% of total simulated ΔT_w for Test 5 (Figure 7c). The same analysis can be repeated for Test 6, showing that an earlier warm period explains 86% and 74% of ΔT_w in June and July, respectively (which corresponds to the complement to 100% of the values at the beginning of the target period in Figures 7d and 7e). Interestingly, in both tests, an amplified warming of LST is observed with respect to air temperature, with the total ΔT_w being equal to 2.19°C and 4.31°C in June and July (Table 3), corresponding to the 110% and 215% of ΔT_a (2°C), respectively. This dramatic effect in July results from an earlier onset of strong stratification (Figure 4b), which makes LST warming much faster in the warm year than in the cold year (compare the rising limbs of 1998 and the mean year in Figure 4a).

Differently, in August and September (Figures 7e and 7f) the dependence on past conditions is completely negligible, with the mean LST being essentially controlled by the values of air temperature observed during the target periods (73% and 75% of the total variation, respectively, for the 2 months). The main reason is that the lake is always strongly stratified from August to September, thermal inertia is reduced (causing a short historical memory of the system), and hence LST responds much faster to current air temperature modifications, thus being substantially disconnected from past thermal conditions. A faster response of LST during the start of the fall season could also be attributed to the instability of the atmospheric boundary layer over the lake and to increased heat fluxes through the lake surface, which however in our formulation are implicitly accounted for by a smaller value of the parameter δ . The short-memory effect is more pronounced in the warm year because of the smaller thermal inertia (i.e., stronger stratification): in this case

the decrease of LST in late summer occurs at a similar rate as air temperature (which begins to decline in August), and is faster than in the cold year. Indeed, after a cold summer the lake retains more heat because of the larger thermal inertia, and the decrease of LST is consequently slower. The overall resulting effect is a weaker increase of LST compared to air temperature, being ΔT_w equal to 80% of ΔT_a in August, and 63% of ΔT_a in September, respectively (see Table 3). Analogous considerations were already discussed in section 3.2 when presenting results of Tests 1 (T_a 1998 and δ 1997) and 2 (T_a 1997 and δ 1998).

Results for the JAS period are coherent with those of Tests 3 (Steeper T_a) and 4 (Flatter T_a), with the two periods January–June and JAS contributing almost equally to LST changes in summer (JAS air temperature explains 56% of total ΔT_w for both Tests 5 and 6, see Table 3 and Figure 7g). In the light of the analysis of the individual target months July, August, and September discussed above, it is interesting to note that the JAS-averaged LST is affected by winter and spring conditions only because of July.

Finally, the simultaneous comparison of Test 5 and 6 allows us to identify an intersection point. This represents the day of the year in which the effect of a warm winter with a cold summer is equal to that of a cold winter with a warm summer. Considering June and July, the intersection time is located about 1 month before the start of the target period, while for August and September it is within the period, suggesting again that in these months LST dynamics are chiefly controlled by the concurrent air temperature signal.

4. Discussion

Prediction of LST is crucial because it exerts a strong control on lake's biogeochemistry and ecology, influencing water quality parameters, chemical reaction rates, presence of pathogens, photosynthesis by algae and aquatic plants, and the habitat for macroinvertebrates and fishes [e.g., Wetzel, 2001]. For instance, higher temperatures increase algal photosynthesis and the metabolic rates of most aquatic animals, thus requiring more food and oxygen, which is contrasting with the reduced oxygen flux from the atmosphere to the lake [Winder and Sommer, 2012]. The surface layer of lakes is indeed rich in biodiversity, and at the same is very sensitive to thermal alterations, whereas the deep waters are substantially not affected during the periods of strong stratification, which dramatically reduces the vertical fluxes across the thermocline [e.g., Imberger, 1998]. This isolation of the surface layer from below strengthens with stratification (larger temperature gradients), therefore the reactive volume shrinks and responds more rapidly to the heat exchanged with the atmosphere. The process is governed by complex mechanisms, which are retained in our formulation only in a simplified, yet effective, way. In particular, *air2water* does not need to reconstruct the individual terms of the heat budget, nor the vertical mixing process in the water body. In its lumped formulation, it directly provides the relevant information that is LST and a metric for thermal stratification. Through the calibration phase, the model assimilates the dominant features of the examined lake, with the formulation remaining general and the parameters being specific of the case study. Interestingly, a few parameters capture the thermal behavior (for instance, the single parameter a_4 controls summer stratification), thus leading to a synthetic description and, potentially, to a classification of lake thermal properties. In this respect, a preliminary attempt was presented by Toffolon *et al.* [2014], which analyzed 14 temperate lakes characterized by different morphology. Particularly attractive is the possibility to expand this analysis, exploiting the growing availability of collections of lakes' observational data at the global scale (e.g., Global Lake Temperature Collaboration—GLTC) [Sharma *et al.*, 2015] (Global Lake Ecological Observatory Network—GLEON) [Weathers *et al.*, 2013]. An extensive application of *air2water* would certainly contribute to achieve a more exhaustive overview of lakes' thermal functioning, and a deeper understanding of lakes' thermal response as a function of their morphometric characteristics and climatic conditions.

The simplicity and robustness of *air2water* suggests its possible use in long-term predictions. This is particularly relevant in a future perspective, since lake temperatures are expected to be affected by warming trends as a result of climate change [e.g., Mortsch and Quinn, 1996; Stefan *et al.*, 1998]. Some studies investigated the effect of varying meteorological forcing (including air temperature) on lake thermal dynamics by means of process-based one-dimensional, two-dimensional, and three-dimensional numerical models [Leon *et al.*, 2005; Yamashiki *et al.*, 2010; Wahl and Peeters, 2014]. Three-dimensional (3-D) models are especially designed to describe the individual processes, but require large computational times and are usually applied for short-term simulations. Moreover, they require detailed time series of meteorological data as

input (e.g., wind speed, humidity, cloudiness, etc., in addition to air temperature), which are often not available or not provided with the needed time resolution.

In order to overcome these limitations, regression models have been widely used in climate change impact assessment [e.g., Dokulil, 2014], extrapolating LST from air temperature measurements on the basis of linear or nonlinear regression relationships. These models are indeed attractive because of their simplicity and limited requirement of meteorological data, but their use may be questionable in a climate change context especially when it is necessary to extrapolate temperature values beyond the limits of the measured time series.

In this perspective, *air2water* represents a valuable alternative tool to regression models, which require the same data in input but are not able to address some fundamental processes (e.g., the hysteresis cycle between air and water temperature). Furthermore, it can be used in place of process-based models when meteorological data are not sufficient for a proper calibration during a reference climate scenario. In this respect, we note that downscaling climate projections from the coarse resolution of the climate models to a finer scale suitable for the predictive lake model is a complex issue. In order to apply the downscaling procedure, a significantly large amount of historical data is required for all the meteorological forcing, with the drawback that the downscaling of some variables (especially precipitation, cloudiness, wind, and radiative fluxes) is usually associated with large uncertainties [e.g., Dettinger, 2013]. Differently, *air2water* requires as an input variable a quantity whose downscaling procedure is very robust, i.e., air temperature, and thus can be seen as a valuable tool in climate change impact studies, allowing for predictions of future trends of lake surface water temperature. Finally, the possibility of coupling *air2water*, as a lumped lake model, in atmospheric circulation and weather prediction models has also been evidenced by Tofflon *et al.* [2014].

5. Conclusions

The present analysis contributes to the understanding of the conditions leading to extreme warming of lake surface temperature (LST). These events typically occur for a positive combination of two factors: (i) warm winter and spring lead to the early onset of stratification, thus reducing the volume (and hence the thermal inertia) of the lake surface layer; (ii) with a relatively thin layer reacting to the surface heat flux, a warm summer season can determine a pronounced warming of LST, which rapidly adapts to air temperature conditions in July, August, and September (JAS). Such a description is confirmed by the analysis of the exceptional warming of Lake Superior in summer 1998, which was determined by the contribution of both factors.

The *air2water* model was shown to be able to capture the positive feedback between LST and stratification. Therefore, we analyzed some synthetic scenarios in order to isolate the relative contributions of the different factors, showing that the increase of LST during summer 1998 is almost equally ascribable to what happened in January–June (and hence to the stratification conditions at the beginning of the considered period) and to air temperature in JAS. This result is in line with Austin and Colman [2007] who based their analysis on a different method relying on long-term trends. The model also suggests that July is the month where the increase of LST with respect to a warmer air temperature is maximum, and strongly dependent upon the previous months, while LST in August and September is affected only by concurrent air temperature. Given that the JAS period is often considered in statistical analyses of Lake Superior, our results suggest the opportunity to use a more appropriate index to characterize the changes in the thermal behavior of the lake in summer. In this respect, we recommend future analyses to separate the contribution of the different summer months on the averaged warming, since the mechanisms that control LST in July are inherently different from those in August and September.

As a whole, the results point out that the dynamics of thermal stratification of the lake are crucial for the seasonal evolution of surface water temperature, and should be carefully considered in this kind of analyses. Despite the simplicity of the model, we demonstrated that *air2water* is able to satisfactorily reproduce the main processes controlling the response of LST, and can be effectively used to perform sensitivity analyses aimed at evaluating the role exerted by air temperature in controlling the seasonal behavior of stratification and LST. This possibility is especially relevant in climate change studies, where the scenarios for water temperature (when available) contain much more uncertainties than those for air temperature. Thus, *air2water* may contribute to reconstruct first approximations of the future dynamics of LST as one of the main drivers of ecology and biogeochemistry in lentic waters.

Appendix A: Net Heat Flux at the Lake-Atmosphere Interface

The net heat flux per unit surface Φ_{net} in equation (1) can be decomposed in the following terms (defined as positive when directed to the lake surface layer):

$$\Phi_{net} = \Phi_S + \Phi_A + \Phi_W + \Phi_L + \Phi_C + \Phi_P + \Phi_I + \Phi_D, \quad (A1)$$

where Φ_S is the net shortwave solar radiation actually absorbed by the water volume, Φ_A is the net longwave radiation emitted from the atmosphere toward the river, Φ_W is the longwave radiation emitted from the water, Φ_L is the latent heat flux due to evaporation/condensation, Φ_C is the sensible heat flux due to convection, Φ_P is the heat flux due to precipitation, Φ_I is the effect of the throughflow by inlets and outlets, and Φ_D is the heat flux exchanged with deep water.

Following Piccolroaz *et al.* [2013], which the reader is referred to for further details, we linearize all the terms composing Φ_{net} as a function of T_w and T_a as follows:

$$\Phi = \Phi_0 + \left. \frac{\partial \Phi}{\partial T_a} \right|_{T_{a0}} (T_a - T_{a0}) + \left. \frac{\partial \Phi}{\partial T_w} \right|_{T_{w0}} (T_w - T_{w0}), \quad (A2)$$

where Φ is a generic flux term present in (A1), T_{w0} and T_{a0} are reference values (e.g., long-term averages) of LST and air temperature, respectively, and Φ_0 is the part of the heat flux that is independent of variation of air and water temperatures, but that can vary in time. We further assume that, for the purposes of the present analysis, the term $\Phi_0 + \left. \frac{\partial \Phi}{\partial T_a} \right|_{T_{a0}} + \left. \frac{\partial \Phi}{\partial T_w} \right|_{T_{w0}}$ can be approximately described by the sum of a constant value and a sinusoidal function of time with a period of 1 year.

Hence, substituting (A1) and (A2) into equation (1) we obtain:

$$\frac{dT_w}{dt} = \frac{A}{\rho c_p V_s} (\hat{a}_1 + \hat{a}_2 T_a - \hat{a}_3 T_w + \hat{a}_t), \quad (A3)$$

where \hat{a}_1 , \hat{a}_2 , and \hat{a}_3 are coefficients that can be directly derived by the heat flux terms once suitable empirical relationships are adopted [e.g., Martin and McCutcheon, 1998], and \hat{a}_t is the sinusoidal term. We also introduce the dimensionless ratio $\delta = V_s/V_r$, where the reference volume V_r is left unspecified. Thus, equation (A3) can be rewritten as:

$$\frac{dT_w}{dt} = \frac{V_r}{V_s} \left(\frac{A \hat{a}_1}{\rho c_p V_r} + \frac{A \hat{a}_2}{\rho c_p V_r} T_a - \frac{A \hat{a}_3}{\rho c_p V_r} T_w + \frac{A \hat{a}_t}{\rho c_p V_r} \right), \quad (A4)$$

which it is straightforward to reformulate as in equation (2) by introducing δ and the parameters a_1 , a_2 , a_3 , a_5 , and a_6 .

In order to account for the strong seasonal variations of the reactive layer volume, we assume that δ is a function of the difference between LST and the deep water temperature T_h as specified in equation (3), with the introduction of three additional parameters: a_4 to quantify the effect of direct stratification, and a_7 and a_8 to account for inverse stratification and the effect of ice cover (we impose $T_w \geq 0$ as a lower limit for simulated LST).

Finally, we note that the last two terms of equation (A1) are not explicitly considered in the analysis, but their effect is implicitly retained by the other parameters through the calibration procedure. We refer to Toffolon *et al.* [2014] for a more extensive discussion of the physical interpretation of the parameters.

References

- Adrian, R., *et al.* (2009), Lakes as sentinels of climate change, *Limnol. Oceanogr. Methods*, 54(6 part 2), 2283–2297, doi:10.4319/lo.2009.54.6_part_2.2283.
- Anderson, W. L., D. M. Robertson, and J. J. Magnuson (1996), Evidence of recent warming and El Niño-related variation in ice breakup of Wisconsin lakes, *Limnol. Oceanogr.*, 41(5), 815–821.
- Assel, R. A., J. E. Janowiak, D. Boyce, C. O'Connors, F. H. Quinn, and D. C. Norton (2000), Laurentian Great Lakes ice and weather conditions for the 1998 El Niño Winter, *Bull. Am. Meteorol. Soc.*, 81, 703–717, doi:10.1175/1520-0477(2000)081<0703:LGLIAW>2.3.CO;2.

Acknowledgments

The authors are grateful to NOAA (National Oceanic and Atmospheric Administration) for air temperature and LST data used in this work (data can be downloaded from <http://www.ndbc.noaa.gov/>). Authors are indebted to Jay Austin for kindly providing vertical temperature profiles (National Science Foundation grant OCE-0825633), and for the helpful comments and contribution. We also thank Michael Notaro for the discussion during the early stage of the work, and John Lenters and other three anonymous Reviewers for their valuable suggestions. The source code (written in Fortran 90/95) of *air2water*, the precompiled executable files (Linux/Windows), a readme file, and an example application are available at the following link <https://zenodo.org/record/30231#.VeccbvnFblU> [Piccolroaz and Toffolon, 2015]. The code is published under the Creative Commons Attribution-ShareAlike 3.0 license.

- Austin, J. A., and S. M. Colman (2007), Lake Superior summer water temperatures are increasing more rapidly than regional air temperatures: A positive ice-albedo feedback, *Geophys. Res. Lett.*, *34*, L06604, doi:10.1029/2006GL029021.
- Austin, J. A., and S. M. Colman (2008), A century of temperature variability in Lake Superior, *Limnol. Oceanogr. Methods*, *53*(6), 2724–2730, doi:10.4319/lo.2008.53.6.2724.
- Blanken, P. D., C. Spence, N. Hedstrom, and J. D. Lenters (2011), Evaporation from Lake Superior. 1: Physical controls and processes, *J. Great Lakes Res.*, *37*(4), 707–716, doi:10.1016/j.jglr.2011.08.009.
- Cline T. J., V. Bennington, and J. F. Kitchell (2013), Climate change expands the spatial extent and duration of preferred thermal habitat for Lake Superior fishes, *PLoS ONE*, *8*(4), e62279, doi:10.1371/journal.pone.0062279.
- Coats, R., J. Perez-Losada, G. Schladow, R. Richards, and C. Goldman (2006), The warming of Lake Tahoe, *Clim. Change*, *76*(1–2), 121–148, doi:10.1007/s10584-005-9006-1.
- De Senerpont Domis, L. N., et al. (2013), Plankton dynamics under different climatic conditions in space and time, *Freshwater Biol.*, *58*, 463–482, doi:10.1111/fwb.12053.
- Dettinger, M. (2013), Projections and downscaling of 21st century temperatures, precipitation, radiative fluxes and winds for the Southwestern US, with focus on Lake Tahoe, *Clim. Change*, *116*, 17–33.
- Dokulil, M. T. (2014), Predicting summer surface water temperatures for large Austrian lakes in 2050 under climate change scenarios, *Hydrobiologia*, *731*, 19–29, doi:10.1007/s10750-013-1550-5.
- Gupta H. V., H. Kling, K. K. Yilmaz, and G. F. Martinez (2009), Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *J. Hydrol.*, *377*(1–2), 80–91, doi:10.1016/j.jhydrol.2009.08.003.
- Hampton, S. E., L. R. Izmet'eva, M. V. Moore, S. L. Katz, B. Dennis, and E. A. Silow (2008), Sixty years of environmental change in the world's largest freshwater lake—Lake Baikal, Siberia, *Global Change Biol.*, *14*(8), 1947–1958, doi:10.1111/j.1365-2486.2008.01616.x.
- Imberger, J., (Ed.) (1998), Physical processes in lakes and oceans, *Coastal Estuarine Stud.*, vol. 54, AGU, Washington, D. C.
- Lenters, J. D. (2004), Trends in the Lake Superior water budget since 1948: A weakening seasonal cycle, *J. Great Lakes Res.*, *30*, suppl. 1, 20–40, doi:10.1016/S0380-1330(04)70375-5.
- Lenters, J. D., S. J. Hook, and P. B. McIntyre (2012), Workshop examines warming of lakes worldwide, *Eos Trans. AGU*, *93*(43), 427, doi:10.1029/2012EO430004.
- Leon, L. F., D. C. L. Lam, W. Schertzer, and D. Swayne (2005), Lake and climate models linkage: A 3-D hydrodynamic contribution, *Adv. Geosci.* *4*, 57–62, doi:10.5194/adgeo-4-57-2005.
- Livingstone, D. M. (2003), Impact of secular climate change on the thermal structure of a large temperate central European lake, *Clim. Change*, *57*(1–2), 205–225, doi:10.1023/A:1022119503144.
- Lofgren, B. M., and Y. Zhu (1999), Seasonal climatology of surface energy fluxes on the Great Lakes, *National Oceanic and Atmospheric Administration (NOAA) Tech. Memo. ERL GLERL-112*, NOAA, Ann Arbor, Mich.
- Magnuson, J. J., et al. (1997), Potential effects of climate changes on aquatic systems: Laurentian Great Lakes and Precambrian Shield region, *Hydrol. Processes*, *11*(8), 825–871, doi:10.1002/(SICI)1099-1085(19970630)11:8 < 825::AID-HYP509 > 3.0.CO;2-G.
- Martin, J. L., and S. McCutcheon (1998), *Hydrodynamics and Transport for Water Quality Modeling*, CRC Press, Boca Raton, Fla.
- McCormick, M. J., and G. L. Fahnenstiel (1999), Recent climatic trends in nearshore water temperatures in the St. Lawrence Great Lakes, *Limnol. Oceanogr.*, *44*(3), 530–540, doi:10.4319/lo.1999.44.3.0530.
- Mortsch, L. D., and F. H. Quinn (1996), Climate change scenarios for Great Lakes Basin ecosystem studies, *Limnol. Oceanogr.*, *41*(5), 903–911.
- Piccolroaz, S., and M. Toffolon (2015), air2water (v1.0.0): A model to predict lake surface temperature using air temperature, Zenodo, doi:10.5281/zenodo.30231, Last accessed on 31 August 2015.
- Piccolroaz S., M. Toffolon, and B. Majone (2013), A simple lumped model to convert air temperature into surface water temperature in lakes, *Hydrol. Earth Syst. Sci.*, *17*, 3323–3338, doi:10.5194/hess-17-3323-2013.
- Quayle, W., L. Peck, H. Peat, J. Ellis-Evans, and P. Harrigan (2002), Extreme responses to climate change in Antarctic lakes, *Science*, *295*(5555), 645, doi:10.1126/science.106407.
- Schindler, D. W., S. E. Bayley, B. R. Parker, K. G. Beaty, D. R. Cruikshank, E. J. Fee, E. U. Schindler, and M. P. Stainton (1996), The effects of climatic warming on the properties of boreal lakes and streams at the Experimental Lakes Area, northwestern Ontario, *Limnol. Oceanogr.*, *41*(5), 1004–1017.
- Schneider, P., and S. J. Hook (2010), Space observations of inland water bodies show rapid surface warming since 1985, *Geophys. Res. Lett.*, *37*, L22405, doi:10.1029/2010GL045059.
- Schneider, P., S. J. Hook, R. G. Radocinski, G. K. Corlett, G. C. Hulley, S. G. Schladow, and T. E. Steissberg (2009), Satellite observations indicate rapid warming trend for lakes in California and Nevada, *Geophys. Res. Lett.*, *36*, L22402, doi:10.1029/2009GL040846.
- Schwab, B. J., G. A. Leshkevich, and G. C. Muhr (1999), Automated mapping of surface water temperature in the Great Lakes, *J. Great Lakes Res.*, *25*(3), 468–481, doi:10.1016/S0380-1330(99)70755-0.
- Sharma, S., et al. (2015), A global database of lake surface temperatures collected by in situ and satellite methods from 1985–2009, *Sci. Data*, *2*, 150008, 19 pp., doi:10.1038/sdata.2015.8.
- Stefan, H. G., X. Fang, and M. Hondzo (1998), Simulated climate change effects on year-round water temperatures in temperate zone lakes, *Clim. Change*, *40*, 547–576.
- Thiery, W., et al. (2014), LakeMIP Kivu: Evaluating the representation of a large, deep tropical lake by a set of 1-dimensional lake models, *Tellus, Ser. A*, *66*, 21,390, doi:10.3402/tellusa.v66.21390.
- Titze, D. D., and J. A. Austin (2014), Winter thermal structure of Lake Superior, *Limnol. Oceanogr. Methods*, *59*(4), 1336–1348, doi:10.4319/lo.2014.59.4.1336.
- Toffolon, M., S. Piccolroaz, B. Majone, A.-M. Soja, F. Peeters, M. Schmid, and A. Wüest (2014), Prediction of surface temperature in lakes with different morphology using air temperature, *Limnol. Oceanogr. Methods*, *59*(6), 2185–2202, doi:10.4319/lo.2014.59.6.2185.
- Van Cleave, K., J. D. Lenters, J. Wang, and E. M. Verhamme (2014), A regime shift in Lake Superior ice cover, evaporation, and water temperature following the warm El Niño winter of 1997–1998, *Limnol. Oceanogr. Methods*, *59*(6), 1889–1898, doi:10.4319/lo.2014.59.6.1889.
- Verburg, P., R. E. Hecky, and H. Kling (2003), Ecological consequences of a century of warming in Lake Tanganyika, *Science*, *301*(5632), 505–507, doi:10.1126/science.1084846.
- Vollmer, M. J., H. A. Bootsma, R. E. Hecky, G. Patterson, J. D. Halfman, J. M. Edmond, D. H. Eccles, and R. F. Weiss (2005), Deep-water warming trend in Lake Malawi, East Africa, *Limnol. Oceanogr. Methods*, *50*(2), 727–732, doi:10.4319/lo.2005.50.2.0727.
- Wahl, B., and F. Peeters (2014), Effect of climatic changes on stratification and deep-water renewal in Lake Constance assessed by sensitivity studies with a 3D hydrodynamic model, *Limnol. Oceanogr. Methods*, *59*, 1035–1052, doi:10.4319/lo.2014.59.3.1035.

- Weathers K. C., et al. (2013), The Global Lake Ecological Observatory Network (GLEON): The evolution of grassroots network science, *Bull. Limnol. Oceanogr.*, 22(3), 71–73.
- Wetzel, R. G. (2001), *Limnology: Lake and River Ecosystems*, 3rd ed., Academic, San Diego.
- Williamson, C. E., J. E. Saros, W. F. Vincent, and J. P. Smol (2009), Lakes and reservoirs as sentinels, integrators, and regulators of climate change, *Limnol. Oceanogr. Methods*, 54(6 part 2), 2273–2282, doi:10.4319/lo.2009.54.6_part_2.2273.
- Winder, M., and U. Sommer (2012), Phytoplankton response to a changing climate, *Hydrobiologia*, 698, 5–16, doi:10.1007/s10750-012-1149-2.
- Yamashiki, Y., M. Kato, K. Takara, E. Nakakita, and M. Kumagai (2010), Sensitivity Analysis on Lake Biwa under the A1B SRES climate change scenario using Biwa-3D Integrated Assessment Model. Part I: Projection of lake temperature, *Hydrol. Res. Lett.*, 4, 45–49, doi:10.3178/HRL.4.45.