

Flag bundles on Fano manifolds

Gianluca Occhetta^a, Luis E. Solá Conde^{a,*}, Jarosław A. Wiśniewski^b

^a*Dipartimento di Matematica, Università di Trento, via Sommarive 14 I-38123 Trento (TN), Italy*

^b*Instytut Matematyki UW, Banacha 2, 02-097 Warszawa, Poland*

Abstract

As an application of a recent characterization of complete flag manifolds as Fano manifolds having only \mathbb{P}^1 -bundles as elementary contractions, we consider here the case of a Fano manifold X of Picard number one supporting an unsplit family of rational curves whose subfamilies parametrizing curves through a fixed point are rational homogeneous, and we prove that X is homogeneous. In order to do this, we first study minimal sections on flag bundles over the projective line, and discuss how Grothendieck's theorem on principal bundles allows us to describe a flag bundle upon some special sections.

Résumé

Un résultat récent identifie les variétés de drapeaux complets comme les variétés de Fano n'ayant que des fibrations en \mathbb{P}^1 comme contractions élémentaires. On va se servir de cette caractérisation pour étudier une variété de Fano X de nombre de Picard un admettant une famille non scindée de courbes rationnelles dont les sous-familles qui paramétrisent les courbes par un point choisi sont rationnellement homogènes. On montre qu'un tel X est homogène. Dans ce but, on étudie d'abord les sections minimales de fibrés en drapeaux sur la droite projective et on discute comment le théorème de Grothendieck sur des fibrés principaux permet de décrire un fibré en drapeaux à partir de certaines sections spéciales.

Keywords: Fano manifolds, flag bundles, Grothendieck's theorem

2010 MSC: Primary 14J45, Secondary 14E30, 14M15, 14M17

1. Introduction

Although vector bundles and their projectivizations play an important role in every branch of Algebraic Geometry, a framework in which these objects are especially manageable is the one of algebraic varieties containing rational curves. The main reason for this is the fact, attributed to C. Segre for rank two (see [20, p. 44] for historical remarks on the general statement), that every vector bundle over the projective line \mathbb{P}^1 is isomorphic to a direct sum of line bundles. However, it was not until Alexandre Grothendieck's celebrated paper [6] that this theorem achieved an optimal form, in the framework of principal bundles. In fact, Grothendieck shows (by reducing the general case, via the adjoint representation of G , to the study of orthogonal bundles over \mathbb{P}^1) that every principal G -bundle –for G reductive– is diagonalizable (see Theorem 3.3 below).

Our interest in this topic comes from its relation with certain homogeneity criteria that we have been recently considering ([17, 19]) within a project whose goal is the Campana–Peternell conjecture, which predicts that every Fano manifold with nef tangent bundle is rational homogeneous. In a nutshell, we showed that flag manifolds are characterized within the class of Fano manifolds by having only \mathbb{P}^1 -bundles as elementary contractions. In particular, one may then try to use this result to prove the homogeneity of a certain Fano manifold X by “untangling” its families

*Corresponding author.

Email addresses: gianluca.occhetta@unitn.it (Gianluca Occhetta), eduardo.solaconde@unitn.it (Luis E. Solá Conde), J.Wisniewski@uw.edu.pl (Jarosław A. Wiśniewski)

of extremal rational curves, constructing upon X another Fano manifold \widetilde{X} dominating it and satisfying the above property.

More concretely, this “bottom-up” strategy may be roughly described as follows: we start from a Fano manifold whose homogeneity we want to check; we consider a (not necessarily complete) proper dominating family of minimal rational curves in X , $\mathcal{M} \xleftarrow{p} \mathcal{U} \xrightarrow{q} X$, and ask ourselves whether $\widetilde{X} = \mathcal{U}$ is again a Fano manifold and, in this case, proceed by substituting X by \widetilde{X} . If this procedure can be carried out until we get to a Fano manifold in which all the families of minimal rational curves are \mathbb{P}^1 -bundles, then the original variety X will be homogeneous.

This process can be shortened in the particular case in which q is smooth and its fibers $\mathcal{M}_x := q^{-1}(x)$ are homogeneous, since rational homogeneous bundles are determined by principal bundles, leading us immediately to a complete flag bundle $\widetilde{\mathcal{U}}$ dominating \mathcal{U} (see Section 2 below). In order to check that $\widetilde{\mathcal{U}}$ is in fact a complete flag, we need to study sections of the bundle $\widetilde{\mathcal{U}}$ over minimal rational curves in X , which is our motivation to give a “rational curves oriented” interpretation of Grothendieck’s theorem.

Section 3 is devoted to this topic. More concretely, we will study G/B -bundles (with G semisimple and $B \subset G$ a Borel subgroup) over the Riemann sphere \mathbb{P}^1 . We will see that (up to a choice of a Cartan subgroup $H \subset B$) in each of these bundles we may define a set of sections, that we call *fundamental*, that are in one-to-one correspondence with the Weyl group of G and that, under this correspondence, reflections of the root system correspond to \mathbb{P}^1 -bundles containing pairs of these sections. Moreover, we will show that one of these sections is minimal –in a deformation theoretical sense, see Definition 3.1– and that the G/B -bundle is determined by this minimal section and by its self-intersection numbers within the $\text{rk}(G)$ \mathbb{P}^1 -bundles containing it. This information may be then represented by what we call a *tagged Dynkin diagram* (see Theorem 3.23 for a precise statement).

As an application, we prove, in Section 4, the following statement:

Theorem 1.1. *Let X be a Fano manifold of Picard number one, and $p : \mathcal{U} \rightarrow \mathcal{M}$ be an unsplit dominating complete family of rational curves satisfying that the evaluation morphism $q : \mathcal{U} \rightarrow X$ is smooth. Assume furthermore that the fiber $\mathcal{M}_x = q^{-1}(x)$ is a rational homogeneous space for every $x \in X$. Then X is rational homogeneous.*

The relation of this statement with the Campana–Peternell conjecture comes from the fact that the smoothness of q is satisfied by any unsplit family of rational curves in X if we assume that T_X is nef.

We note also that this result resembles the main theorem in [7] (proven first by Mok for Hermitian symmetric spaces and homogeneous contact manifolds, see [16]), but there are certain differences between the two statements: on one hand Hwang and Hong need to assume that the image of \mathcal{M}_x into $\mathbb{P}(\Omega_{X,x})$, the so called *VMRT of \mathcal{M} at x* , is projectively equivalent to the VMRT of a rational homogeneous manifold X' , while we do not need to consider any particular projective embedding of \mathcal{M}_x . On the other hand, they only need to check the above property on a general point $x \in X$, while we need to assume that q is smooth, and that every \mathcal{M}_x is rational homogeneous. Note that if the relative Picard number $\rho(\mathcal{U}/\mathcal{M})$ is one, it is enough to check the homogeneity condition for a general $x \in X$ (see [12, Section 3]).

Unfortunately, one cannot expect to characterize all the rational homogeneous manifolds of Picard number one in this way, since there are some examples for which \mathcal{M}_x is not homogeneous ([14], see also [18, Proposition 2.20]). We expect that a similar treatment of non complete families of rational curves may lead to a solution of this problem.

Finally we note that Theorem 1.1 depends on a characterization of complete flag manifolds which is a slightly stronger version of the main theorem in [19], see Theorem A.1. We show how to prove this statement in Appendix A.

Warning: In previous papers ([17, 18, 19]) we used the term “smooth \mathbb{P}^1 -fibration” for any smooth morphism whose fibers are isomorphic to \mathbb{P}^1 , reserving the name “ \mathbb{P}^1 -bundle” for the Grothendieck projectivization of a rank two vector bundle. In the present paper, which is about smooth rational homogeneous fiber bundles, the term “ \mathbb{P}^1 -bundle” refers to a smooth morphism whose fibers are isomorphic to \mathbb{P}^1 .

2. Rational homogeneous bundles

Throughout this paper all the varieties will be smooth, projective and defined over the field of complex numbers. A (*smooth, projective*) *fiber bundle over X* is a (smooth, projective) morphism $q : E \rightarrow X$ between smooth varieties. We will be interested in the case in which q is smooth, projective, and also *isotrivial*, which means that all the fibers

of q over closed points are isomorphic; if F is a variety isomorphic to the fibers of q , we will say that the fiber bundle is an F -bundle. According to a classical theorem of Fischer and Grauert (cf. [2, page 29]), it follows that a smooth projective F -bundle q is *locally trivial* in the analytic topology: there exists then an open covering $\{U_i, i \in I\}$ of X with the analytic topology and isomorphisms $\phi_i : U_i \times F \rightarrow q^{-1}(U_i)$ commuting with the corresponding projections onto U_i , named *trivializations of q* . We may consider $\{\phi_i\}$ to be the whole atlas of trivializations of q , and then q may be reconstructed from the complex varieties $U_i \times F$ by means of the transitions

$$\phi_{ij} = \phi_j^{-1} \circ \phi_i : U_{ij} \times F \longrightarrow U_{ij} \times F, \quad (U_{ij} := U_i \cap U_j),$$

or, equivalently by the corresponding maps $\theta_{ij} : U_{ij} \rightarrow \text{Aut}(F)$ satisfying $\phi_{ij}(x, y) = (x, \theta_{ij}(x)(y))$. It turns out that the ϕ_{ij} define a cocycle $\theta \in H^1(X, \text{Aut}(F))$, which completely determines the F -bundle $q : E \rightarrow X$. Note that this cocycle is defined over the analytic space associated to X , that we denote also by X , by abuse of notation.

We will assume also that every fiber F of q is *rational homogeneous*, i.e., that it is isomorphic to the quotient of a semisimple group by a parabolic subgroup.

Remark 2.1. *Note that the representation of F as a quotient of a semisimple group is not unique. Besides eventual choice of two semisimple groups with the same universal cover, in some occasions a rational homogeneous variety F may be written as the quotient of two semisimple groups with different associated Lie algebras. This is the case of \mathbb{P}^{2n-1} (that can be written as a quotient of $\text{Sl}(2n)$, $\text{PGL}(2n-1)$ —these two with the same Lie algebra— and $\text{Sp}(2n)$), of the quadric Q^5 (quotient of SO_7 and G_2), and the spinor variety parametrizing $r-1$ dimensional linear subspaces in a quadric of dimension $2r-1$ (quotient of SO_{2r+1} and SO_{2r+2}). For irreducible rational homogeneous manifolds (i.e. rational homogeneous manifolds that are not products), this is the complete list (see [1, Ch. 3, p. 75]).*

However, it is known that the identity component of the automorphism group $\text{Aut}(F)$ is semisimple [8, Thm. 3.11], and it acts transitively on F . We will denote it by G_F , or G if there is no possible confusion, and then we may write F as a quotient G/P for some parabolic subgroup $P \subset G$. If we further assume that the base X is simply connected (this is the case, for instance, if X is Fano or, more generally, rationally connected), then it follows that the cocycle $\theta \in H^1(X, \text{Aut}(F))$ lies in $H^1(X, G)$; otherwise, since G is a normal subgroup of $\text{Aut}(F)$, θ would define a nonzero cocycle in $H^1(X, \text{Aut}(F)/G)$, which in turn defines an étale covering of X , a contradiction.

Assume now that $q : E \rightarrow X$ is an F -bundle defined by a cocycle $\theta \in H^1(X, G)$, for a rational homogeneous manifold F whose automorphism group has identity component G . Then θ defines a G -bundle $q_G : E_G \rightarrow X$, given by the glueing of $U_i \times G$ by means of the transition functions:

$$\theta_{ij} : U_{ij} \rightarrow G \hookrightarrow \text{Aut}(G),$$

where G embeds in $\text{Aut}(G)$ as the set of automorphism given by left multiplication with elements of G . There is then a well defined action of G on E_G , given by right multiplications by elements of G on every chart. We may now say that the G -bundle E_G is a *principal G -bundle* over X . If we now take any parabolic subgroup $P' \subset G$, we may consider the G/P' -bundle:


$$q_{P'} : E_{P'} := E_G \times_G G/P' := (E_G \times G/P') / \sim_G \rightarrow X,$$

where \sim_G is the equivalence relation defined by $(x, gP') \sim_G (xh, h^{-1}gP')$ for every $h \in G$. One may easily check this G/P' -bundle is defined also by the same cocycle $\theta \in H^1(X, G)$. Note that $H^1(X, G)$ is the cohomology of X , regarded as an analytic space, with values in G , so $E_{P'}$ is, a priori, only defined as a compact analytic space. However, since we can embed $E_{P'}$ into the projectivization of the holomorphic vector bundle $q_{P'*}(\omega_{E_{P'}/X}^{-r}) \cong E_G \times_{P'} H^0(G/P', \omega_{G/P'}^{-r})$ (for $r \gg 0$), which is an algebraic vector bundle by the GAGA principle ([21]), it follows that $E_{P'}$ is indeed a projective variety.

Furthermore, given two parabolic subgroups $P_1 \subset P_2 \subset G$, one has a natural map $q_{P_1, P_2} : E_{P_1} \rightarrow E_{P_2}$ satisfying $q_{P_2} \circ q_{P_1, P_2} = q_{P_1}$. In particular we will consider a Borel subgroup B of G contained in P , and define $\bar{E} := E_B$, with projection map $\bar{q} = q_B : \bar{E} \rightarrow X$, and, for every parabolic $P' \supset B$, a map $\bar{q}_{P'} := q_{B, P'} : \bar{E} \rightarrow E_{P'}$. In particular we have a projection $\bar{q}_P : \bar{E} \rightarrow E_P = E$, that we will denote by $\pi : \bar{E} \rightarrow E$. The bundle $\bar{q} : \bar{E} \rightarrow X$ will be called the *flag bundle* associated to $q : E \rightarrow X$. Note that π is also a flag bundle over E , whose fiber we write as G'/B' , where G' is the the semisimple group obtained by subsequently quotienting P by its unipotent subgroup and then by the center, and B' is the image of B in this quotient.

We will denote by k the *rank of the group* G , which can be defined as the dimension of a maximal torus contained in B , or as the number of simple roots of the root system of the Lie algebra \mathfrak{g} of G , or as the number of minimal parabolic subgroups of G containing B . Note that, denoting by P_1, \dots, P_k these minimal parabolic subgroups, the morphisms $\pi_i := \tilde{q}_{P_i} : \tilde{E} \rightarrow E_i := E_{P_i}$ are \mathbb{P}^1 -bundles.

For the reader's convenience, we present the following diagram that illustrates the notation used (corresponding to a minimal parabolic subgroup $P_i \subset P$):



Finally, we may describe the relative canonical class of a G/B -bundle as above. It is well known that $K_{G/B}$ may be described as follows. Consider $\Delta := \{\alpha_i, i = 1, \dots, k\}$ a base of simple roots of the root system Φ of the Lie algebra \mathfrak{g} of G , so that every positive root $\beta \in \Phi^+$ is a nonnegative integer combination of the α_i 's, and define the integers b_1, \dots, b_k by the formula:

$$\sum_{\beta \in \Phi^+} \beta = \sum_{i=1}^k b_i \alpha_i. \quad (1)$$

Consider also the relative canonical divisors K_t of the projections $G/B \rightarrow G/P_t$. Then:

$$K_{G/B} = \sum_{i=1}^k b_i K_i.$$

Table 1 below contains the values of the integers b_1, \dots, b_k , for G simple (the numbering of the nodes corresponds to the one in [10, p. 58]).

\mathcal{D}	b_1, \dots, b_k
A_k	$k, (k-1)2, \dots, 2(k-1), k$
B_k	$2k-1, 2(2k-2), \dots, (k-1)(k+1), k^2$
C_k	$2k, 2(2k-1), \dots, (k-1)(k+2), k(k+1)/2$
D_k	$2k-2, 2(2k-3), \dots, (k-2)(k+1), (k-1)k/2, (k-1)k/2$
E_6	16, 22, 30, 42, 30, 16
E_7	34, 49, 66, 96, 75, 52, 27
E_8	92, 136, 182, 270, 220, 168, 114, 58
F_4	16, 30, 42, 22
G_2	10, 6

Table 1: Coefficients of $K_{G/B}$ in terms of the relative canonical divisors K_t .

On the other hand, one may show that $T_{G/B}$ may be constructed upon the relative tangent bundles of the projections $G/B \rightarrow G/P_t$, via successive use of Lie brackets. We refer the reader to [17, Construction 1] for details on this construction. In our situation, we may show that, identifying $\pi_i|_{\tilde{q}^{-1}(x)} : \tilde{q}^{-1}(x) \rightarrow \tilde{q}_i^{-1}(x)$ with the projection $G/B \rightarrow G/P_t$, for any $x \in X$ then

$$\mathcal{O}_{\tilde{E}}(K_{\pi_i})|_{\tilde{q}^{-1}(x)} = \mathcal{O}_{G/B}(K_t), \quad (2)$$

and the corresponding successive use of Lie brackets produces the relative tangent bundle of \widetilde{E} . In particular, one obtains:

Lemma 2.2. *The relative canonical class of \widetilde{q} satisfies:*

$$K_{\widetilde{q}} = \sum_{i=1}^k b_i K_{\pi_i}.$$

The same argument allows us to compute (the pull-back to \widetilde{E} of) the relative canonical class of $q : E \rightarrow X$. Set $D = \{1, \dots, k\}$; let $I \subset D$ be the subset defining the parabolic subgroup P , i.e. the set of indices such that the Lie subalgebra $\mathfrak{p} \subset \mathfrak{g}$ of P satisfies:

$$\mathfrak{p} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Phi^+} \mathfrak{g}_{-\alpha} \oplus \bigoplus_{\alpha \in \Phi^+(I)} \mathfrak{g}_{\alpha}, \quad (3)$$

where $\Phi^+(I)$ denotes the subset of Φ^+ generated by the simple roots $\{\alpha_i, i \notin I\}$. Then, arguing as before, we have:

Lemma 2.3. *With the same notation as above, let us consider the integers c_1, \dots, c_k determined by:*

$$\sum_{\beta \in \Phi^+(I)} \beta = \sum_{i=1}^k c_i \alpha_i.$$

Then:

$$\pi^* K_q = \sum_{i=1}^k (b_i - c_i) K_{\pi_i}. \quad (4)$$

PROOF. It is enough to note $\pi^* K_q$ equals the difference $K_{\widetilde{q}} - K_{\pi}$, where the first summand has been computed in Lemma 2.2, and the second can be computed by applying the same lemma to the flag bundle $\pi : \widetilde{E} \rightarrow E$.

3. Flag bundles over \mathbb{P}^1 : remarks on Grothendieck's theorem

In this section we will study rational homogeneous bundles over \mathbb{P}^1 . More concretely, since all of them are contractions of flag bundles, we will concentrate on the case of a G/B -bundle $\widetilde{q} : \widetilde{E} \rightarrow \mathbb{P}^1$. We will use the notation introduced in Section 2.

Inspired by the case of projective bundles over \mathbb{P}^1 , that are determined by the restriction of their relative tangent bundles to a minimal section, we will study sections of $\widetilde{\pi}$ that satisfy the following minimality condition:

Definition 3.1. *A section $s : \mathbb{P}^1 \rightarrow \widetilde{E}$ of \widetilde{q} is called minimal if, for every $x \in \mathbb{P}^1$, the irreducible component H of the scheme $\text{Hom}_{\mathbb{P}^1}(\mathbb{P}^1, \widetilde{E}; x, s(x))$ parametrizing sections of \widetilde{q} sending x to $s(x)$ is zero-dimensional.*

The existence of minimal sections on the G/B -bundle \widetilde{E} is an immediate consequence of the Bend and Break Lemma (see for instance [4, Proposition 3.11]), once we prove the existence of a section of any kind; this, in turn, may be obtained, either from a much more general result of Graber, Harris and Starr, [5], or by means of ad hoc arguments for each of the possible types of semisimple group G , or (as we will do later) by means of Grothendieck's theorem on principal G -bundles.

Definition 3.2. *Given $\sigma : \mathbb{P}^1 \rightarrow \widetilde{E}$ a section of $\widetilde{q} : \widetilde{E} \rightarrow \mathbb{P}^1$, we may consider the projections $\pi_t : \widetilde{E} \rightarrow E_t$, where $P_t, t = 1, \dots, k$. As noted before, the π_t 's are \mathbb{P}^1 -bundles, and we may consider the surfaces \widetilde{E}_t appearing as the fiber products of \widetilde{E} and the section σ , for every t :*

$$\begin{array}{ccc} \widetilde{E}_t & \xrightarrow{\quad} & \widetilde{E} \\ \sigma_t \uparrow \pi_t & \nearrow \sigma & \downarrow \pi_t \\ \mathbb{P}^1 & \xrightarrow{\pi_t \circ \sigma} & \mathbb{P}^1 \end{array}$$

Then, for every $t = 1, \dots, k$, \widetilde{E}_t is a \mathbb{P}^1 -bundle, that we call the pivotal \mathbb{P}^1 -bundles of \widetilde{q} with respect to σ .

Note that the section σ defines a section $\sigma_t : \mathbb{P}^1 \rightarrow \widetilde{E}_t$. In the case in which σ is minimal in \widetilde{E} , then it is also minimal in \widetilde{E}_t . In particular the integers $d_t = K_{\pi_t} \cdot \sigma(\mathbb{P}^1)$ are non negative. The main goal of this section is to show that a minimal section σ , together with the non negative integers d_t , determine completely the G/B -bundle \widetilde{E} , and the main tool we will use is Grothendieck's theorem on principal G -bundles over \mathbb{P}^1 .

3.1. Grothendieck's theorem

Let G be a semisimple Lie group, and $H \subset G$ be a Cartan subgroup, which is a maximal abelian subgroup of G . Let N denote the normalizer of H in G , whose quotient N/H is called the *Weyl group of G with respect to H* . Given any smooth complex variety X we may consider the cohomology of X with values in the sheafified groups H and G (that, by abuse of notation, we denote with the same symbols), and the corresponding natural map $H^1(X, H) \rightarrow H^1(X, G)$. This map is equivariant with respect to the conjugation action of the groups N and H . Since the latter acts trivially on both sets, and $N \subset G$ acts trivially on the second, we finally have a map:

$$H^1(X, H)/W \rightarrow H^1(X, G) \quad (5)$$

Grothendieck's theorem says the following:

Theorem 3.3 (Grothendieck's theorem (cf. [6])). *With the same notation as above, for $X = \mathbb{P}^1$, the map (5) is a bijection.*

This theorem was originally stated for the case of a reductive group G , but we will stick to the semisimple case, which is the one we are interested in.

Remark 3.4. *The standard geometric interpretation of this theorem concerns the case $G = \mathrm{PGL}(r+1)$, where the theorem tells us that any \mathbb{P}^r -bundle over \mathbb{P}^1 is the projectivization of a direct sum of line bundles. For instance, for $r = 1$ and a cocycle $\theta \in H^1(\mathbb{P}^1, G)$, any choice of a Cartan subgroup $H \subset G$ corresponds to a choice of two sections C_0 and C_∞ of the corresponding \mathbb{P}^1 -bundle $\widetilde{E} : \widetilde{E} \rightarrow \mathbb{P}^1$, satisfying that $C_0 \cdot C_\infty = 0$. These two sections correspond to two quotients $\mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}^1}(a_i)$, $i = 1, 2$, where \widetilde{E} is isomorphic to the projectivization of a bundle $\mathcal{E} \cong \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \mathcal{O}_{\mathbb{P}^1}(a_2)$. Typically C_0 is associated with the quotient to $\mathcal{O}_{\mathbb{P}^1}(a_1)$, when we choose $a_1 \leq a_2$. Note that C_0 is a minimal section of the \mathbb{P}^1 -bundle, in the sense of Definition 3.1.*

Setup 3.5. *Throughout the rest of Section 3, G will denote a semisimple Lie group, $H \subset B \subset G$ a pair of Cartan and Borel subgroups, and $\theta \in H^1(\mathbb{P}^1, H)$ a cocycle, whose image into $H^1(\mathbb{P}^1, G)$ defines a G/B -bundle $\widetilde{q} : \widetilde{E} \rightarrow X$.*

Remark 3.6. *Note that all the possible Borel subgroups of G are conjugate, and the choice of different Borel subgroups produces isomorphic G/B -bundles. On the other hand, the choice of a preimage of θ into $H^1(\mathbb{P}^1, B)$ via the natural map $H^1(\mathbb{P}^1, B) \rightarrow H^1(\mathbb{P}^1, G)$ (in the standard language, the choice of a reduction of E_G via the inclusion $B \hookrightarrow G$), is equivalent to the choice of a holomorphic section of $\widetilde{q} : \widetilde{E} \rightarrow X$.*

Remark 3.7. *Following [6], the kernel L of the exponential map sending every $v \in \mathfrak{h}$ to $\exp(2\pi i v) \in H$ may be identified with the dual lattice of the lattice in \mathfrak{h}^\vee generated by the root system Φ . Moreover we have natural homomorphisms of groups:*

$$H^1(\mathbb{P}^1, H) \cong H^2(\mathbb{P}^1, L) \cong L \subset \mathfrak{h}, \quad (6)$$

satisfying that the conjugation action of W in $H^1(\mathbb{P}^1, H)$ corresponds to the restriction of the dual action of W on \mathfrak{h} , as the Weyl group of Φ .

3.2. Fundamental sections of a flag bundle

Let us start by noting some general facts on parabolic subgroups, and their Levi decompositions:

Remark 3.8. *Given a semisimple Lie group G and a Cartan subgroup H , let us denote with gothic fonts $\mathfrak{h}, \mathfrak{g}$, as usual, their associated Lie algebras. Then $\mathfrak{h} \subset \mathfrak{g}$ defines a root system $\Phi \subset \mathfrak{h}^\vee$. Any choice of a base of positive simple roots $\Delta = \{\alpha_t, t = 1, \dots, k\}$ corresponds to a choice of a fundamental chamber of the action of the Weyl group W on \mathfrak{h} , $\mathcal{P} = \{v \in \mathfrak{h} \mid \alpha_t \cdot v \geq 0, \forall t\} \subset \mathfrak{h}$, to a choice of a decomposition $\Phi = \Phi^+ \cup \Phi^-$ of the root system in two sets of positive*

and negative roots, and to a choice of a Borel subalgebra $\mathfrak{b} := \mathfrak{h} \oplus \bigoplus_{\beta \in \Phi^-} \mathfrak{g}_\beta$ and a corresponding Borel subgroup $B \subset G$ containing H ([9, 27.3]). Given such a Borel subgroup B , for every $t = 1, \dots, k$ we denote by $P_t \supset B$ the parabolic subgroups of G whose Lie algebras are $\mathfrak{b} \oplus \mathfrak{g}_{\alpha_t}$. Dividing P_t by its unipotent radical first, and then by its center, we obtain a semisimple group G_t , isomorphic to $\mathrm{PGL}(2)$, satisfying that the images H_t of H , and B_t of B , into G_t , are respectively a Cartan and a Borel subgroup of G_t . Note that we have an isomorphism $G_t/B_t \cong P_t/B$, and the natural map $P_t/B \subset G/B$ allows us to identify G_t/B_t with the fiber of the canonical projection $G/B \rightarrow G/P_t$ passing by the class of the identity modulo B . Moreover, the induced homomorphism of Lie groups

$$\phi : H \rightarrow \prod_t H_t \quad (7)$$

is finite and surjective, since one may easily check that its differential at the identity gives an isomorphism of Lie algebras $d\phi_e : \mathfrak{h} \rightarrow \bigoplus_t \mathfrak{h}_t$.

Remark 3.9. Conversely, if we start from a semisimple group G and a Borel subgroup B , we may consider the minimal parabolic subgroups of G containing B , P_1, \dots, P_k . Considering, as above, the semisimple part G_t of P_t , we have a natural surjective map $B \rightarrow \prod_t B_t$. Given, for every t , a Cartan subgroup $H_t \subset B_t$, the inverse image of $\prod_t H_t$ is a Cartan subgroup of G contained in B .

Example 3.10. In the case of $G = \mathrm{PGL}(2)$, each of the two possible choices of a basis of positive simple roots $\Delta = \{\alpha\}$ provides an isomorphism $\alpha : \mathfrak{h} \rightarrow \mathbb{C}$, under which the lattice L defined in Remark 3.7 may be identified with $\mathbb{Z} \subset \mathbb{C}$. Now, if we start from a cocycle $\theta \in H^1(\mathbb{P}^1, H)$, we may choose $\Delta = \{\alpha\}$ so that the image of θ in L via (6) has nonnegative α -coordinate. This integer is classically known as the invariant of the \mathbb{P}^1 -bundle defined by θ , which is the intersection of its minimal section C_0 with its relative canonical line bundle.

In our general setting, we may choose the fundamental chamber $\mathcal{P} \subset \mathfrak{h}$ so that the image of $\theta \in H^1(\mathbb{P}^1, H)$ into \mathfrak{h} belongs to \mathcal{P} and, by Remark 3.6, we may assume, without loss of generality, that B is the Borel subgroup of G containing H corresponding to this choice of \mathcal{P} . The corresponding base of simple roots is denoted by $\Delta = \{\alpha_1, \dots, \alpha_k\}$. Any other Borel subgroup of G containing H is the conjugation of B by an element $w \in W$, hence it will be denoted by B_w .

Definition 3.11. In the setup of 3.5, with the same notation as above the image of $\theta \in H^1(\mathbb{P}^1, H)$ into each of the sets $H^1(\mathbb{P}^1, B_w)$, $w \in W$, defines a section of the G/B -bundle $\bar{q} : \bar{E} \rightarrow \mathbb{P}^1$, that we denote by $\sigma_w : \mathbb{P}^1 \rightarrow C_w \subset \bar{E}$. These sections are called the fundamental sections of \bar{E} with respect to θ . In particular, the section corresponding to B will be called the minimal fundamental section of \bar{E} with respect to θ , and denoted by $\sigma : \mathbb{P}^1 \rightarrow C \subset \bar{E}$.

Remark 3.12. Note that the set of fundamental sections of \bar{E} is in one to one correspondence with the set of possible Weyl chambers of W in \mathfrak{h} , which is known to be bijective to W .

For instance, in the case of a \mathbb{P}^1 -bundle treated in Remark 3.4, given $H \subset G$ we have precisely two Borel subgroups containing it, each of them giving rise to one the (fundamental) sections C_0 or C_∞ . In our general setup, we may consider, for every Borel subgroup B_w containing H , and each positive simple root $\alpha_t \in \Delta$, the pivotal \mathbb{P}^1 -bundle $\bar{E}_{w,t} := (\pi_t \circ \sigma_w)^* \bar{E}$, fitting in the cartesian square:

$$\begin{array}{ccc} \bar{E}_{w,t} & \xrightarrow{\quad} & \bar{E} \\ \sigma_{w,t} \uparrow & \nearrow \sigma_w & \downarrow \pi_t \\ \mathbb{P}^1 & \xrightarrow{\pi_t \circ \sigma_w} & E_t \end{array}$$

Let us denote by C_w the image of the section σ_w . On the other hand, we may consider the reflection $r_t \in W$ corresponding to a positive simple root α_t , and the section $\sigma_{r_t w} : \mathbb{P}^1 \rightarrow \bar{E}_{r_t w, t}$, with image $C_{r_t w}$. Then we may claim the following, that follows from [9, 29.3 Lemma B]:

Proposition 3.13. *With the same notation as above, for every $w \in W$ and every $\alpha_t \in \Delta$,*

$$\widetilde{E}_{w,t} = \widetilde{E}_{r_t w, t},$$

and the sections C_w and $C_{r_t w}$ are the C_0 and C_∞ sections of the \mathbb{P}^1 -bundle $\widetilde{E}_{w,t}$.

Remark 3.14. *The correspondence between $\{C_w, C_{r_t w}\}$ and $\{C_0, C_\infty\}$ depends on the particular choices of t and w , and may be expressed as follows: C_w is the C_0 section of $\widetilde{E}_{w,t}$ if and only if $w(\alpha_t)$ is a positive root. In the particular case of $B_w = B$, it follows that the corresponding minimal fundamental section C is a minimal (fundamental) section for every pivotal \mathbb{P}^1 -bundle \widetilde{E}_t containing it.*

Remark 3.15. *Note also that each pivotal \mathbb{P}^1 -bundle \widetilde{E}_t is defined by the image θ_t of the cocycle θ via the natural map $H^1(\mathbb{P}^1, H) \rightarrow H^1(\mathbb{P}^1, H_t)$, induced by the group homomorphism $H \rightarrow H_t$ defined in Remark 3.8.*

3.3. The tag of a flag bundle

Let us consider now the differential at the identity of the homomorphism ϕ of complex tori (7), together with the corresponding exponential maps defined in Remark 3.7. We have a commutative diagram of group homomorphisms, with exact rows:

$$\begin{array}{ccccc} L & \longrightarrow & \mathfrak{h} & \xrightarrow{\exp(2\pi i _)} & H \\ \downarrow & & \downarrow \phi_{*e} & & \downarrow \phi \\ \bigoplus_t L_t & \longrightarrow & \bigoplus_t \mathfrak{h}_t & \xrightarrow{\exp(2\pi i _)} & \prod_t H_t \end{array} \quad (8)$$

where L_t denotes the kernel lattice for each H_t .

Definition 3.16. *With the same notation as above, denoting and by $\alpha'_t \in \mathfrak{h}_t^*$ the positive root associated to the Borel subgroup B_t , for each t , the homomorphism of lattices:*

$$\begin{array}{ccc} & \delta & \\ & \curvearrowright & \\ L & \xrightarrow{\phi_{*e}} \bigoplus_t L_t & \xrightarrow{\oplus_t \alpha'_t} \mathbb{Z}^k \end{array}$$

is called the tagging map of G , and the image of $\theta \in H^1(\mathbb{P}^1, H) \cong L$ via δ , $\delta(\theta)$ is called the tag of the G/B -bundle $\widetilde{q} : \widetilde{E} \rightarrow \mathbb{P}^1$ defined by θ .

Proposition 3.17. *With the same notation as above, the G/B -bundle $\widetilde{q} : \widetilde{E} \rightarrow \mathbb{P}^1$ defined by θ is completely determined by its tag $\delta(\theta) = (d_1, \dots, d_k)$. Moreover, for every t , $d_t = K_{\pi_t} \cdot C$, being C the minimal fundamental section of \widetilde{E} introduced in Definition 3.11.*

PROOF. For the first part it is enough to check that the map $L \rightarrow \bigoplus_t L_t$ is injective, and this follows from the fact that $\phi_{*e} : \mathfrak{h} \rightarrow \bigoplus_t \mathfrak{h}_t$ is an isomorphism. For the second, note that d_t equals the invariant of the corresponding pivotal \mathbb{P}^1 -bundle \widetilde{E}_t , hence it may be interpreted as the intersection number of the relative canonical divisor K_{π_t} with the minimal fundamental section C .

Remark 3.18. *As a consequence of Proposition 3.17, the tag of any G/B -bundle over \mathbb{P}^1 with respect to a cocycle θ lies in $\mathbb{Z}_{\geq 0}^k$. Then $\delta(\theta)$ shall be thought of as the choice of a non negative integer d_t for every positive simple root of G (with respect to the choice of a base of simple roots, as in Remark 3.8). In particular it is natural to represent it as a tagged Dynkin diagram, i.e. by ‘‘tagging’’ the t -th node of the Dynkin diagram, corresponding to the root α_t , with the non negative integer d_t . A tagged Dynkin diagram obtained in this way is called admissible with respect to G . Note that a flag manifold may be written as the quotient of different semisimple groups with the same Dynkin diagrams, for which the concept of admissibility is not the same, in general. Hence it is not true that, for a G/B -bundle over \mathbb{P}^1 , any k -tuple of non negative integers is admissible with respect to G .*

Example 3.19. For instance, a \mathbb{P}^n -bundle $\mathbb{P}(\bigoplus_i \mathcal{O}_{\mathbb{P}^1}(a_i))$, with $a_0 \leq \dots \leq a_n$, is determined by the tag $(a_1 - a_0, \dots, a_n - a_{n-1})$ on the Dynkin diagram A_n (with the standard numbering of its nodes). Hence, every tag is admissible for the group $\mathrm{PGL}(n+1)$, whereas for the group $G = \mathrm{Sl}(n+1)$, the lattice $\delta(L)$ of admissible tags has index n in \mathbb{Z}^n .

3.4. Minimal fundamental sections vs Minimal sections

Finally we will show the relation between the minimal fundamental sections of a G/B -bundle and its minimal sections.

Lemma 3.20. *With the same notation as above, the minimal fundamental section C of a G/B -bundle defined by a cocycle $\theta \in H^1(\mathbb{P}^1, H)$ is a minimal section in the sense of Definition 3.1.*

PROOF. Since, by construction, C has degree greater than or equal to zero with respect to the relative canonical divisors K_{π_t} , we may claim that the relative tangent bundle of \tilde{E} over C , that can be obtained (cf. [17, Construction 1]) from the relative canonical bundles $\mathcal{O}_{\tilde{E}}(K_{\pi_t})$ by means of successive Lie brackets, has no positive summands, and hence C has no deformations with a point fixed.

Conversely, we will show now that a minimal section of a G/B -bundle is a minimal fundamental section, for a certain choice of a Cartan subgroup. In other words, minimal sections are not necessarily unique, but they are determined by the different choices of Cartan subgroups of G .

Proposition 3.21. *Let $\tilde{q} : \tilde{E} \rightarrow \mathbb{P}^1$ be a G/B -bundle over \mathbb{P}^1 , and let $\sigma' : \mathbb{P}^1 \rightarrow C' \subset \tilde{E}$ be a minimal section in the sense of Definition 3.1. Then there exists a Cartan subgroup $H' \subset G$ such that C' is a minimal fundamental section with respect to H' .*

PROOF. The section σ' provides an inverse image θ' in $H^1(X, B)$ of the cocycle in $H^1(X, G)$ defining \tilde{E} . With the same notation as in Remark 3.8, we consider the images θ'_t of θ' into $H^1(\mathbb{P}^1, B_t)$, $t = 1, \dots, k$, defined by the minimal parabolic subgroups P_t containing B ; they define k pivotal \mathbb{P}^1 -bundles \tilde{E}'_t containing C' as a section. Since C' is minimal in \tilde{E} in the sense of Definition 3.1, it is also minimal in each \tilde{E}'_t . But on a \mathbb{P}^1 -bundle it is clear that a minimal section is a minimal fundamental section, from what it follows that $\theta'_t \in H^1(\mathbb{P}^1, B_t)$ has a (unique) inverse image in $H^1(\mathbb{P}^1, H'_t)$, for certain Cartan subgroups H'_t , $t = 1, \dots, k$. By Remark 3.9, the inverse image of $\prod_t H'_t$ into B is a Cartan subgroup H' , and the k -tuple $(\theta'_1, \dots, \theta'_k)$ has a unique inverse image $\theta'' \in H^1(\mathbb{P}^1, H')$, which maps into $\theta' \in H^1(\mathbb{P}^1, B)$. We conclude by noting that, by construction, C' is a minimal fundamental section of \tilde{E} with respect to H' .

Remark 3.22. *Given a G/B -bundle \tilde{q} , we have defined its tag with respect to a choice of Cartan subgroup $H \subset G$ and a cocycle $H^1(\mathbb{P}^1, H)$. But two of these possible choices, say $\theta \in H^1(\mathbb{P}^1, H)$ and $\theta' \in H^1(\mathbb{P}^1, H')$, are related by conjugation with an element $g \in G$, $H' = gHg^{-1}$. We may then consider a holomorphic map $f : \mathbb{C} \rightarrow G$ satisfying $f(0) = e$ and $f(1) = g$. We choose B and the minimal section C with respect to H as above, and consider, for every $z \in \mathbb{C}$, the corresponding cocycle $f(z)\theta f(z)^{-1} \in H^1(\mathbb{P}^1, f(z)Hf(z)^{-1})$ and its image into $H^1(\mathbb{P}^1, f(z)Bf(z)^{-1})$, that defines the minimal fundamental section of \tilde{q} with respect to $f(z)Hf(z)^{-1}$. We conclude that any two minimal fundamental sections of \tilde{q} are numerically equivalent, hence $\delta(\theta)$ depends only on the isomorphism class of the G/B -bundle.*

Summing up, we may write the following statement:

Theorem 3.23. *Let G be a semisimple Lie group, and $B \subset G$ be a Borel subgroup of G . For any admissible tagged Dynkin diagram \mathcal{D} there exist a G/B -bundle $\tilde{q} : \tilde{E} \rightarrow \mathbb{P}^1$, unique up to isomorphism, satisfying that the self-intersection of a minimal section C of \tilde{q} in the t -th pivotal \mathbb{P}^1 -bundle associated to C is equal to the tag d_t of the corresponding t -th node of \mathcal{D} .*

4. An application to the Campana–Pernell Conjecture

Along this section, which is devoted to the proof of Theorem 1.1, X will be a *CP-manifold*, that is a Fano manifold with nef tangent bundle. We will consider an unsplit complete family $\mathcal{M} \xleftarrow{p} \mathcal{U} \xrightarrow{q} X$ of rational curves in X . By definition, a *complete family of rational curves* is the universal family of curves parametrized by an irreducible component \mathcal{M} of $\text{RatCurves}^n(X)$. Then, the condition on the nefness of T_X tells us that the evaluation morphism q is dominant and smooth (see, for instance, [18, Prop. 2.10]). For any $x \in X$, the smooth subvariety $q^{-1}(x)$, which is assumed to be a rational homogeneous manifold, will be denoted by \mathcal{M}_x . Let us start by fixing some extra notation.

Notation 4.1. We will denote by G the identity component of $\text{Aut}(\mathcal{M}_x)$, so that every \mathcal{M}_x is isomorphic to the quotient of G by a parabolic subgroup P . Then a Borel subgroup B of G contained in P defines a G/B -bundle over X , that we denote by $\tilde{q} : \tilde{\mathcal{U}} \rightarrow X$, that admits a contraction $\pi : \tilde{\mathcal{U}} \rightarrow \mathcal{U}$. Note that π is also a flag bundle over \mathcal{U} , whose fiber we write as G'/B' (as in Section 2, G' is the semisimple group obtained by subsequently quotienting P by its unipotent subgroup and then by the center, and B' is the image of B in this quotient).

We will denote by k the rank of G , and set $D = \{1, \dots, k\}$. Every index $i \in D$ corresponds to a minimal parabolic subgroup $P_i \subset G$ containing B , that provides an elementary contraction $\pi_i : \tilde{\mathcal{U}} \rightarrow \mathcal{U}_i$, which is a \mathbb{P}^1 -bundle whose relative canonical classes we denote by K_i . We set $I := \{i \in D \mid P_i \subset P\}$, so that the contraction $\pi : \tilde{\mathcal{U}} \rightarrow \mathcal{U}$ factors via π_i , for every $i \in I$.

Let $f : \mathbb{P}^1 \rightarrow X$ be the normalization of any curve Γ of the family \mathcal{M} and consider the pull-back $f^*\mathcal{U}$; since Γ is an element of the family there is a natural section $s : \mathbb{P}^1 \rightarrow s(\mathbb{P}^1) \subset f^*\mathcal{U}$. Denote by $\tilde{s} : \mathbb{P}^1 \rightarrow s^*f^*\tilde{\mathcal{U}}$ a minimal section of the G'/B' -bundle $s^*f^*\tilde{\mathcal{U}}$ over \mathbb{P}^1 and by $\tilde{\Gamma}$ its image. We have then the following commutative diagram:

$$\begin{array}{ccccc}
 & & \tilde{s} & & \\
 & & \swarrow & & \\
 s^*f^*\tilde{\mathcal{U}} & \xrightarrow{\pi} & \mathbb{P}^1 & & \\
 \downarrow \lrcorner & & \downarrow s & \searrow & \\
 f^*\tilde{\mathcal{U}} & \xrightarrow{\pi} & f^*\mathcal{U} & \xrightarrow{q} & \mathbb{P}^1 \\
 \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow f \\
 \mathcal{M} & \xleftarrow{p \circ \pi} & \tilde{\mathcal{U}} & \xrightarrow{\pi} & \mathcal{U} & \xrightarrow{q} & X \\
 & & \swarrow p & & & & \\
 & & & & & &
 \end{array}$$

The curve $s(\mathbb{P}^1)$ is contracted by the composition map $f^*\mathcal{U} \rightarrow \mathcal{U} \rightarrow \mathcal{M}$ and, moreover, it cannot deform in $f^*\mathcal{U}$, otherwise its deformations would give a positive dimensional subvariety of \mathcal{M} determining the same curve in X . Hence, identifying $s^*f^*\tilde{\mathcal{U}}$ with its image in $f^*\tilde{\mathcal{U}}$, we may say that $\tilde{\Gamma}$ is a minimal sections of both the G/B -bundle $f^*\tilde{\mathcal{U}}$ and the G'/B' -bundle $s^*f^*\tilde{\mathcal{U}}$. By Proposition 3.21 this minimal section can be regarded as a minimal fundamental section for each of the bundles, and Theorem 3.23 tells us that the two bundles are determined by two tagged Dynkin diagrams supported on the Dynkin diagrams \mathcal{D} and \mathcal{D}' of G and G' , respectively.

Moreover, \mathcal{D}' corresponds to the Dynkin subdiagram of \mathcal{D} obtained by eliminating from \mathcal{D} the nodes indexed by $I \subset D$ and, by the interpretation of the tags given in Proposition 3.17, the tags agree with the inclusion $\mathcal{D}' \subset \mathcal{D}$.

Lemma 4.2. *With the same notation as above, the G'/B' -bundle $s^*f^*\tilde{\mathcal{U}}$ is trivial over \mathbb{P}^1 .*

PROOF. Recall that the t -th tag of G/B is given by $d_t := \tilde{\Gamma} \cdot K_t$. In view of Theorem 3.23 it is enough to show that $d_t = 0$ for every $t \notin I$. By Lemma 2.3 applied to the G/B -bundle $f^*\tilde{\mathcal{U}} \rightarrow \mathbb{P}^1$:

$$\pi^*K_q = \sum_{t=1}^k (b_t - c_t)K_{\pi_t}. \tag{9}$$

where the integers b_t, c_t are given by

$$\sum_{\beta \in \Phi^+} \beta = \sum_{i \in D} b_i \alpha_i, \quad \sum_{\beta \in \Phi^+(I)} \beta = \sum_{j \in D} c_j \alpha_j = \sum_{j \in D \setminus I} c_j \alpha_j. \quad (10)$$

Note that, by definition, we have $b_t \geq c_t$ for every $t \in D$. By Equation (9), we may then write, for any $x \in X$

$$\dim(\mathcal{M}_x) = \pi^* K_q \cdot \bar{\Gamma} = \sum_{j \notin I} (b_j - c_j) d_j + \sum_{i \in I} b_i d_i, \quad (11)$$

where the first equality follows from the fact that $\dim(\mathcal{M}_x) = -K_X \cdot \Gamma - 2 = K_q \cdot f(s(\mathbb{P}^1))$ and the projection formula.

Since the minimality of $\bar{\Gamma}$ implies that $d_t \geq 0$ for all $t \in D$, we may conclude by showing the following:

$$(4.2.1) \quad b_j - c_j > 0 \text{ for all } j \notin I.$$

$$(4.2.2) \quad \sum_{i \in I} b_i d_i \geq \dim(\mathcal{M}_x).$$

In order to prove (4.2.1), we notice that, having $b_j = c_j$ for some $j \notin I$, is equivalent to say that the only positive roots having nonzero coefficient with respect to α_j are the elements of $\Phi^+(I)$.

Since G is the identity component of $\text{Aut}(G/P)$, it follows that every connected component of the Dynkin diagram \mathcal{D} of G must contain a node corresponding to an index $i \in I$; in particular, the connected component \mathcal{D}' of \mathcal{D} containing the node corresponding to α_j contains a node corresponding to an index $i \in I$. But then the longest root β of the root subsystem determined by \mathcal{D}' is a root of \mathcal{D} , whose coordinates with respect to α_j and α_i are different from zero, hence $\beta \in \Phi^+ \setminus \Phi^+(I)$, which contradicts that $b_j = c_j$.

In order to show (4.2.2), we claim first that $d_i > 0$ for every $i \in I$. In fact, if $d_i = 0$ for some $i \in I$, the i -th pivotal \mathbb{P}^1 -bundle $\tilde{\mathcal{U}}_i$ which maps isomorphically into $f^* \mathcal{U}$ via π , would be isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$. But a section of $\tilde{\mathcal{U}}_i$ maps to $s(\mathbb{P}^1)$, which is contracted by $p \circ f$; hence the image of $\pi(\tilde{\mathcal{U}}_i)$ into \mathcal{M} would be a curve, whose points would parametrize the same curve in X , a contradiction.

At this point, in order to conclude it is enough to check that

$$\sum_{i \in I} b_i \geq \#(\Phi^+ \setminus \Phi^+(I)) = \dim(\mathcal{M}_x).$$

The last equality follows from the hypothesis $\mathcal{M}_x = G/P$. Since every root in $\Phi^+ \setminus \Phi^+(I)$ increases at least in a unity at least an integer b_i , the first inequality follows, and the lemma is proved.

Corollary 4.3. *Let X be as in Theorem 1.1. Then, with the same notation as above, the morphism $p \circ \pi : \tilde{\mathcal{U}} \rightarrow \mathcal{M}$ factors via a smooth \mathbb{P}^1 -bundle $\tilde{p} : \tilde{\mathcal{U}} \rightarrow \tilde{\mathcal{M}}$, for a smooth projective variety $\tilde{\mathcal{M}}$.*

PROOF. Write, as in Notation 4.1, any fiber F' of $\pi : \tilde{\mathcal{U}} \rightarrow \mathcal{U}$ as a quotient G'/B' . By Lemma 4.2, the fibers of the smooth morphism $p \circ \pi$ are isomorphic to the rational homogeneous manifold $G'/B' \times \mathbb{P}^1$. Moreover, since \mathcal{M} is rationally connected (because \mathcal{U} is rationally connected), hence simply connected, $p \circ \pi$ is defined by a cocycle $\theta' \in H^1(\mathcal{M}, G' \times \text{Aut}(\mathbb{P}^1))$. The image of this cocycle via the natural map to $H^1(\mathcal{M}, G')$ provides a G'/B' -bundle $\tilde{\pi} : \tilde{\mathcal{M}} \rightarrow \mathcal{M}$, whose pull-back to \mathcal{U} via p is precisely π .

We may now finish the proof of the main result of this Section.

PROOF OF THEOREM 1.1. Note that, by construction, $\rho_{\tilde{\mathcal{U}}} = \rho_{G/B} + \rho_X = \rho_{G/B} + 1$. On one hand, every minimal parabolic subgroup $P_i \supset B$, $i = 1, \dots, k$ provides a smooth \mathbb{P}^1 -bundle $\tilde{\mathcal{U}} \rightarrow \mathcal{U}_i$. Moreover, the classes Γ_i of the curves contracted by them are independent in $N_1(\tilde{\mathcal{U}})$. On the other hand, the class Γ of a fiber of the morphism \tilde{p} provided by Corollary 4.3 does not lie in the hyperplane of $N_1(\tilde{\mathcal{U}})$ generated by the Γ_i 's, hence $\{\Gamma, \Gamma_1, \dots, \Gamma_k\}$ is a basis of $N_1(\tilde{\mathcal{U}})$. We may now apply Theorem A.1, which is a refined version of the Main Theorem in [19], and claim that $\tilde{\mathcal{U}}$ is rational homogeneous. Then X , which is the image of a contraction of $\tilde{\mathcal{U}}$, is rational homogeneous, too.

Remark 4.4. Note that, along the proof of Theorem 1.1 and its preliminary lemmata, we have not really used the assumption on \mathcal{M} being a component of $\text{RatCurves}^n(X)$. In fact we are only using that $p : \mathcal{U} \rightarrow \mathcal{M}$ is a \mathbb{P}^1 -bundle with an evaluation morphism q onto X , for which the natural map $\mathcal{M} \rightarrow \text{RatCurves}^n(X)$ is surjective onto an irreducible component \mathcal{M}' and finite. In fact, it suffices to assume that $\mathcal{M} \rightarrow \mathcal{M}'$ is surjective and generically finite, since the existence of a curve contracted by $\mathcal{M} \rightarrow \mathcal{M}'$ easily implies the existence of a curve in a fiber of the G/P -bundle $q : \mathcal{U} \rightarrow X$ contracted also by the natural morphism from \mathcal{U} to the universal family \mathcal{U}' over \mathcal{M}' . Since this curve is free in \mathcal{U} , it follows that $\dim(\mathcal{U}) > \dim(\mathcal{U}')$ and hence $\dim(\mathcal{M}) > \dim(\mathcal{M}')$, a contradiction. Hence, this observation allows us to state the following, slightly more general, result.

Corollary 4.5. Let \mathcal{U} be a smooth complex projective algebraic variety, and $p : \mathcal{U} \rightarrow \mathcal{M}$, $q : \mathcal{U} \rightarrow X$ be two smooth fiber bundles, with fibers isomorphic to \mathbb{P}^1 and to a rational homogeneous manifold G/P , respectively. If moreover $b_2(X) = 1$, and the natural map $\mathcal{M} \rightarrow \text{RatCurves}^n(X)$ is a local isomorphism at a general point, then X , \mathcal{M} and \mathcal{U} are rational homogeneous.

A. Appendix: Characterization of flag manifolds revisited

The main result of [19] states that a Fano manifold whose elementary contractions are \mathbb{P}^1 -bundles is necessarily rational homogeneous. In this Appendix we will prove a slightly more general form of that statement, namely:

Theorem A.1. Let X be a smooth projective variety of Picard number n , satisfying that there exist $\Gamma_i \in N_1(X)$, $i = 1, \dots, n$, independent K_X -negative classes generating n extremal rays, whose associated elementary contractions $\pi_i : X \rightarrow X_i$ are smooth \mathbb{P}^1 -bundles. Then X is isomorphic to a flag manifold G/B , for some semisimple group G .

First of all we will show that, as in the proof of [19, Theorem 1.2], we may use the \mathbb{P}^1 -bundles to construct a set of reflections in $N^1(X)$, generating a finite group that turns out to be the Weyl group of a semisimple Lie algebra \mathfrak{g} . We then call *homogeneous model of X* the flag manifold G/B associated to the corresponding Lie group G . One may then identify appropriately the relative anticanonical divisors $-K_i$ of the \mathbb{P}^1 -bundles of X with the relative anticanonical divisors $-\bar{K}_i$ of the homogeneous model G/B , via a linear isomorphism $\psi : N^1(X) \rightarrow N^1(G/B)$ preserving the cohomology of line bundles.

We will then use this result to prove that the cone of curves of X is generated by the classes of the Γ_i 's. This implies that X is a Fano manifold, whose elementary contractions are \mathbb{P}^1 -bundles, and the result follows then from [19, Theorem 1.2].

A.1. The homogeneous model of X

Following [19], we consider W to be the subgroup of $\text{Gl}(N^1(X))$ generated by the involutions of $N^1(X)$ defined by $r_i(L) := L + (L \cdot \Gamma_i)K_i$, where, K_i denotes the relative canonical bundle of π_i . Moreover, for every i we consider the affine involution of $N^1(X)$ defined as the translation of r_i by $K_X/2$, that is $r'_i(L) := r_i(L - K_X/2) + K_X/2$ (note that $K_i \cdot \Gamma_i = K_X \cdot \Gamma_i = -2$, for all i , so we may also write $r'_i(L) := L + (L \cdot \Gamma_i + 1)K_i$). Let us denote by W' the group generated by them, which is the translation by $K_X/2$ of the group W . Then the arguments provided in [19, Section 2] work in our setting, and we may state the following:

Lemma A.2. Under the assumptions of Theorem A.1,

- (a) $H^j(X, L) = H^{j+\text{sgn}(L \cdot \Gamma_i+1)}(X, r'_i(L))$ for every $i = 1, \dots, n$, $j \in \mathbb{Z}$.
- (b) $\chi(X, L) = \pm \chi(X, w'(L))$, for every $L \in \text{Pic}(X)$ and for every $w' \in W'$.
- (c) The group W is finite.

PROOF. Follow verbatim Lemma 2.3, Lemma 2.7, Corollary 2.8 and Proposition 2.9 in [19].

Corollary A.3. With the same notation as above

- (a) W is the Weyl group of a reduced root system $\Phi := \{w(-K_i) \mid w \in W, i = 1, \dots, n\} \subset N^1(X)$, defining a semisimple Lie algebra \mathfrak{g} .

(b) The set $\Delta := \{-K_i, i = 1, \dots, n\}$ is a base of positive simple roots for Φ , and the Cartan matrix of Φ with respect to Δ is the intersection matrix $(-K_i \cdot \Gamma_j)_{ij}$; in particular, this matrix is nonsingular.

(c) Given the decomposition $\Phi = \Phi^+ \cup \Phi^-$ determined by the base Δ , the class $-K_X$ equals $\sum_{\beta \in \Phi^+} \beta$.

PROOF. Items (a) and (b) follow by Corollary 2.10 and Proposition 2.13 in [19]. For the third part, note that $B = \sum_{\beta \in \Phi^+} \beta$ satisfies $B \cdot \Gamma_i = 2$, for all i (see [10, 10.2, Cor. of Lemma B]). Since $-K_X$ satisfies the same property, the equality follows by (b).

We may then consider a semisimple Lie group G with Lie algebra \mathfrak{g} , and the corresponding flag manifold $\bar{X} = G/B$, that we call the *homogeneous model of X* . We denote its corresponding \mathbb{P}^1 -bundles, fibers, relative canonicals, reflections, etc, by $\bar{\pi}_i : \bar{X} \rightarrow \bar{X}_i, \bar{\Gamma}_i, \bar{K}_i, \bar{r}_i$, etc. By Corollary A.3, we may now define linear isomorphisms:

$$\psi : N^1(X) \rightarrow N^1(\bar{X}), \quad \psi' : N_1(X) \rightarrow N_1(\bar{X}),$$

sending K_i to \bar{K}_i and Γ_i to $\bar{\Gamma}_i$, and, therefore, satisfying that $\psi(L) \cdot \psi'(\Gamma) = L \cdot \Gamma$, for all $L \in N^1(X), \Gamma \in N_1(X)$, and sending Φ to $\bar{\Phi}$. Then:

Lemma A.4. *With the same notation as above, the isomorphism ψ sends K_X to \bar{K}_X . Moreover, we have isomorphisms of groups $W \cong \bar{W}, W' \cong \bar{W}'$, compatible via ψ with their actions on $N^1(X)$ and $N^1(\bar{X})$.*

PROOF. The first part follows from Corollary A.3 (c). For the second part, define the isomorphisms of groups by sending r_i to \bar{r}_i and r'_i to \bar{r}'_i .

One of our first goals will be to prove that the map ψ preserves the cohomology of line bundles. In order to do that, we need first to prove that the dimension of X equals the dimension of its homogeneous model.

Remark A.5. *Note that the hypotheses in Theorem A.1 suffice to claim that X is rationally chain connected with respect to the curves $\Gamma_i, i = 1, \dots, n$. In fact, we may consider the rationally connected fibration with respect to the families of deformations of the Γ_i 's (see [13, IV.4.16]), $X \rightarrow Z$, which is a proper map from an open subset of X onto its image. If $\dim(Z) > 0$, then we may choose an effective divisor on Z and consider its strict transform H in X . By construction, H has intersection zero with every Γ_i , a contradiction. In particular we may claim that:*

- $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$ (cf [13, IV.3.8]), and that, by Serre duality,
- $m := \dim(X)$ is the only integer satisfying that $H^m(X, K_X) \neq 0$.

Corollary A.6. *With the same notation as above, $m = \dim(\bar{X})$, which is equal to the length of the longest element of W .*

PROOF. It is enough to find an element of the affine Weyl group $w' \in W'$ satisfying that $w'(\mathcal{O}_X) = K_X$. In fact, if this is the case, the only index m in which K_X has nonzero cohomology would be determined by w' ; since on the other hand we would also have $w'(\mathcal{O}_{\bar{X}}) = K_{\bar{X}}$ (by Lemma A.4), the integer m would be the same for X and \bar{X} .

The condition required is equivalent to find an element $w \in W$ such that $w(-K_X) = K_X$. Now, it is well known (see [11, Section 1.8]) that the longest element of W satisfies this property.

Corollary A.7. *With the same notation as above, $\chi(X, L) = \chi(\bar{X}, \psi(L))$, for all $L \in \text{Pic}(X)$ satisfying that $\psi(L) \in \text{Pic}(\bar{X})$.*

PROOF. As in [19, Def. 2.6] we consider the polynomial function, of degree smaller than or equal to $m, \chi_X : N^1(X) \rightarrow \mathbb{R}$, satisfying that $\chi_X(L) = \chi(X, L)$ for every $L \in \text{Pic}(X)$. On one hand, one may prove that (cf. [19, Prop. 2.9]) χ_X vanishes on the hyperplanes $w'(K_X/2 + (\Gamma_i)^\perp)$ for every i and every $w' \in W'$, and there is one of these hyperplanes for every positive root $\beta \in \Phi^+$. Then χ_X vanishes in at least $m = \dim(\bar{X}) = \#\Phi^+$ hyperplanes of $N^1(X)$. Let $1 + F_\beta = 0$ denote the equation of the hyperplane determined by $\beta \in \Phi^+$; since the degree of χ_X is not greater than m , it follows that

$$\chi_X = a \prod_{\beta \in \Phi^+} (1 + F_\beta),$$

where $a = \chi_X(\mathcal{O}_X)$, which is equal to 1 by Remark A.5. In particular, χ_X is completely determined by the root system Φ , and our statement follows.

Proposition A.8. *With the same notation as above, $h^i(X, L)$ is equal to $h^i(\bar{X}, \psi(L))$, for all $L \in \text{Pic}(X) \cap \psi^{-1}(\text{Pic}(\bar{X}))$, and every integer i .*

PROOF. Let us also denote by $N \subset N^1(X)$ the dual cone of the cone generated by $\Gamma_1, \dots, \Gamma_n$. Note that it is not true a priori –as it is in the case of a flag manifold G/B , and of any X in the setup of [19]– that $N = \text{Nef}(X)$. On the other hand, arguing as in [19, Cor. 2.19], we may state that the interior of N is a fundamental chamber for the action of W on $N^1(X)$, and that the interior of $N' := N + K_X/2$ is a fundamental chamber for the action of W' on $N^1(X)$.

Let us consider a reduced sequence of reflections in W' , $(r'_{i_1}, \dots, r'_{i_m})$, associated with simple roots $(-K_{i_1}, \dots, -K_{i_m})$, whose composition is the longest element w'_0 of the group W' . Given a divisor $L \in N' \cap \text{Pic}(X)$ satisfying that $\psi(L) \in \text{Pic}(\bar{X})$, we may construct recursively the divisors $L_1 := r_{i_1}(L)$, $L_2 := r'_{i_2} \circ r'_{i_1}(L)$, \dots , $L_m := w'_0(L)$. We consider also the corresponding sequence $(\bar{L}_1, \dots, \bar{L}_m)$ of divisors in \bar{X} constructed upon $\bar{L} := \psi(L)$ via the same sequence reflections; note that $\bar{L}_j = \psi(L_j)$ for all j , by Lemma A.4. At every step, the cohomology of L_{j+1} is obtained from the cohomology of L_j by shifting its degree by $+1$ or -1 , as described in Lemma A.2 (a), depending only on the intersection number $L_j \cdot \Gamma_{i_{j+1}}$. In the case of $\bar{X} = G/B$ the degree of the shifting is $+1$ at every step, by the Borel–Weil–Bott theorem. Since $L_j \cdot \Gamma_{i_{j+1}} = \psi(L_j) \cdot \psi'(\Gamma_{i_{j+1}}) = \bar{L}_j \cdot \bar{\Gamma}_{i_{j+1}}$, it follows that at every step the degree of the shifting is $+1$ for X , as well. In particular, for any positive integer i , we have:

$$H^i(X, L) = H^{m+i}(X, w'_0(L)) = 0,$$

and we may conclude that:

$$H^0(X, L) = \chi(X, L) = \chi(\bar{X}, \psi(L)) = H^0(\bar{X}, \psi(L)).$$

Finally, using that the interior of N' is a fundamental chamber for the action of W' on $N^1(X)$, we may use Lemma A.2 (a) to conclude that the statement holds for every divisor $L \in \text{Pic}(X) \cap \psi^{-1}(\text{Pic}(\bar{X}))$.

A.2. The Mori cone of X

We are now ready to finish the proof of Theorem A.1, by reducing it to [19, Theorem 1.2] via the following description of the cone of effective curves of X :

Proposition A.9. *With the same notation as above, $\overline{\text{NE}}(X)$ is the simplicial cone generated by $\Gamma_1, \dots, \Gamma_n$. In particular X is a Fano manifold.*

To this end, we will use the Bott–Samelson varieties of X , whose construction we recall here (see [19, Section 3]):

Construction 1. *Given a sequence $\ell = (l_1, \dots, l_r)$, $l_i \in \{1, 2, \dots, n\}$, set $\ell[s] := (l_1, \dots, l_{r-s})$ for every $s \leq r$; in particular $\ell[0] = \ell$. Fix a point $x \in X$. Given a sequence $\ell = (l_1, \dots, l_r)$ of indices in $\{1, 2, \dots, n\}$, for every $s = 0, \dots, r$ we construct a manifold $Z_{\ell[s]}$, $s = 0, \dots, r$, called the Bott–Samelson variety associated to $\ell[s]$, together with morphisms*

$$f_{\ell[s]} : Z_{\ell[s]} \rightarrow X, \quad p_{\ell[s+1]} : Z_{\ell[s]} \rightarrow Z_{\ell[s+1]},$$

which are defined recursively as follows: for $s = r$ we set $Z_{\ell[r]} := \{x\}$ and let $f_{\ell[r]} : \{x\} \rightarrow X$ be the inclusion. Then for $s < r$ we consider the elementary contraction $\pi_{l_{r-s}} : X \rightarrow X_{l_{r-s}}$ determined by the extremal ray generated by $\Gamma_{l_{r-s}}$, the composition $g_{\ell[s+1]} := \pi_{l_{r-s}} \circ f_{\ell[s+1]} : Z_{\ell[s+1]} \rightarrow X_{l_{r-s}}$, and define $Z_{\ell[s]}$ upon $Z_{\ell[s+1]}$ as the fiber product with $\pi_{l_{r-s}}$:

$$\begin{array}{ccc} Z_{\ell[s]} & \xrightarrow{f_{\ell[s]}} & X \\ p_{\ell[s+1]} \downarrow & \nearrow f_{\ell[s+1]} & \downarrow \pi_{l_{r-s}} \\ Z_{\ell[s+1]} & \xrightarrow{g_{\ell[s+1]}} & X_{l_{r-s}} \end{array}$$

A key result for our proof of Proposition A.9 will be the following:

Proposition A.10. *[Cf. [19, Corollary 3.18]] Let $\ell = (l_1, \dots, l_r)$ be a sequence, with $l_j \in \{1, \dots, n\}$, for all j . Then $\dim f_{\ell}(Z_{\ell}) = \dim Z_{\ell}$ if and only if $w(\ell)$ is reduced.*

Since, with our current assumptions, we do not have the ampleness of $-K_X$, the proof given in [19] needs to be slightly modified, by using weaker versions of [19, Lemma 3.14 and Proposition 3.17]:

Lemma A.11. *Let D be a nef divisor on Z_ℓ . Then $H^i(Z_\ell, f_\ell^*(K_X/2) + D) = 0$ for every $i > 0$.*

PROOF. See [19, Proof of Lemma 3.14].

Proposition A.12. *Let $L \in \text{Pic}(X)$. Then $\chi(Z_\ell, f_\ell^*L) = \chi(\overline{Z}_\ell, \overline{f}_\ell^*\psi(L))$ and, if $L - K_X/2$ is nef, then $h^0(Z_\ell, f_\ell^*L) = h^0(\overline{Z}_\ell, \overline{f}_\ell^*\psi(L))$. In particular, if L is nef as well, then f_ℓ^*L is big if and only if $\overline{f}_\ell^*\psi(L)$ is big.*

PROOF. The first part follows verbatim from the proof of [19, Proposition 3.17]. For the second part we notice that $\psi(\text{Nef}(X)) \subset \text{Nef}(\overline{X})$, since this last cone is the dual of the cone generated by the $\overline{\Gamma}_i$'s, and that $\psi(K_X) = K_{\overline{X}}$. Therefore, if $L - K_X/2$ is nef, then the higher cohomologies of f_ℓ^*L and $\overline{f}_\ell^*\psi(L)$ vanish by Lemma A.11. The last statement follows from [15, Lemma 2.2.3], by applying the argument above to $rL - K_X/2 = (r-1)L + (L - K_X/2)$, for $r \gg 0$.

PROOF OF PROPOSITION A.10. Let us consider the homogeneous model \overline{X} of X and the corresponding Bott-Samelson variety \overline{Z}_ℓ , with evaluation $\overline{f}_\ell : \overline{Z}_\ell \rightarrow \overline{X}$. It is known that the property holds for \overline{f}_ℓ , since $\overline{f}_\ell(\overline{Z}_\ell)$ is the Schubert variety $\overline{Bw(\ell)B}/B$ of $\overline{X} = G/B$.

Take an ample line bundle $L \in \text{Pic}(X)$ satisfying that $L - K_X/2$ is nef. Since $\psi(\text{Nef}(X)) \subset \text{Nef}(\overline{X})$ (see the proof of Proposition A.12), then $\psi(L)$ is ample on \overline{X} . Hence $\overline{f}_\ell^*(\psi(L))$ is big if and only if $w(\ell)$ is reduced, by our first observation above. By Proposition A.12, we then have that f_ℓ^*L is big if and only if $w(\ell)$ is reduced. Clearly f_ℓ^*L is big if and only if $\dim f_\ell(Z_\ell) = \dim Z_\ell$, and the statement is proved.

PROOF OF PROPOSITION A.9. In order to prove that $\overline{\text{NE}}(X)$ is a simplicial cone generated by $\Gamma_1, \dots, \Gamma_n$, it is enough to show that for every proper subset $I \subset \{1, \dots, n\}$ there exists a contraction $\pi_I : X \rightarrow X_I$ satisfying that $\pi_{I*}(\Gamma_i) = 0$ if and only if $i \in I$.

Fix a proper subset $I \subset \{1, \dots, n\}$. Following [13, IV. Theorem 4.16], there exists a proper fibration $\pi_I : X^0 \rightarrow X_I^0$, defined on an open subset $X^0 \subset X$, whose fibers are Chains(I)-equivalence classes: by definition, two points in X are in the same Chains(I)-equivalence class if and only if there exists a connected chain of rational curves, whose irreducible components are curves in the classes Γ_i , $i \in I$, connecting them. In order to conclude the proof, it suffices to show that, π_I is defined on X , that is $X^0 = X$. But, by [3, Proposition 1], the indeterminacy locus $X \setminus X^0$ of π_I is contained in the union of Chains(I)-equivalence classes of dimension greater than the dimension of the general one, hence it is enough to prove that all the Chains(I)-equivalence classes have the same dimension.

Let $\ell = (l_1, \dots, l_r)$ be a reduced list such that $l_i \in I$, for all i , and $w(\ell)$ is the longest word in the subgroup $W_I \subset W$ generated by the reflections r_i , $i \in I$. Let $x \in X$ be any point, and Z_ℓ be the Bott-Samelson variety associated to ℓ such that $Z_{\ell[r]} = \{x\}$.

Clearly $f_\ell(Z_\ell)$ is contained in the Chains(I)-equivalence class containing x ; let us show that the opposite inclusion holds. Assume, by contradiction, that this is not the case. Then there exists $l_{r+1} \in I$ such that, being $\ell' = (l_1, \dots, l_r, l_{r+1})$, we have $f_\ell(Z_\ell) \subsetneq f_{\ell'}(Z_{\ell'})$. Since these two varieties are irreducible, $\dim Z_\ell = \dim(f_\ell(Z_\ell)) < \dim(f_{\ell'}(Z_{\ell'}))$, hence $\dim(f_{\ell'}(Z_{\ell'})) = \dim Z_{\ell'}$. This, by Proposition A.10, contradicts the fact that ℓ' is not reduced.

Then $f_\ell(Z_\ell)$ is the Chains(I)-equivalence class containing x . In particular, since ℓ is reduced, we have $\dim f_\ell(Z_\ell) = \dim Z_\ell = r$, by Proposition A.10 again, and all Chains(I)-equivalence class have the same dimension.

Acknowledgements

First author supported by PRIN project ‘‘Geometria delle variet  algebriche’’. First and second author supported by the Department of Mathematics of the University of Trento. Second and third author supported by the Polish National Science Center project 2013/08/A/ST1/00804.

References

- [1] Akhiezer, D. N., 1995. Lie group actions in complex analysis. Aspects of Mathematics, E27. Friedr. Vieweg & Sohn, Braunschweig.
URL <http://dx.doi.org/10.1007/978-3-322-80267-5>
- [2] Barth, W., Peters, C., Van de Ven, A., 1984. Compact complex surfaces. Vol. 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin.
URL <http://dx.doi.org/10.1007/978-3-642-96754-2>
- [3] Bonavero, L., Casagrande, C., Druel, S., 2007. On covering and quasi-unsplit families of curves. *J. Eur. Math. Soc. (JEMS)* 9 (1), 45–57.
URL <http://dx.doi.org/10.4171/JEMS/71>
- [4] Debarre, O., 2001. Higher-dimensional algebraic geometry. Universitext. Springer-Verlag, New York.
URL <http://dx.doi.org/10.1007/978-1-4757-5406-3>
- [5] Graber, T., Harris, J., Starr, J., 2003. Families of rationally connected varieties. *J. Amer. Math. Soc.* 16 (1), 57–67 (electronic).
URL <http://dx.doi.org/10.1090/S0894-0347-02-00402-2>
- [6] Grothendieck, A., 1957. Sur la classification des fibrés holomorphes sur la sphère de Riemann. *Amer. J. Math.* 79, 121–138.
- [7] Hong, J., Hwang, J.-M., 2008. Characterization of the rational homogeneous space associated to a long simple root by its variety of minimal rational tangents. In: *Algebraic geometry in East Asia—Hanoi 2005*. Vol. 50 of *Adv. Stud. Pure Math.* Math. Soc. Japan, Tokyo, pp. 217–236.
- [8] Huckleberry, A., 2008. Actions of complex Lie groups and the Borel-Weil correspondence. In: *Symmetries in complex analysis*. Vol. 468 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, pp. 99–123.
URL <http://dx.doi.org/10.1090/conm/468/09145>
- [9] Humphreys, J. E., 1975. Linear algebraic groups. Springer-Verlag, New York-Heidelberg, graduate Texts in Mathematics, No. 21.
- [10] Humphreys, J. E., 1978. Introduction to Lie algebras and representation theory. Vol. 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second printing, revised.
- [11] Humphreys, J. E., 1990. Reflection groups and Coxeter groups. Vol. 29 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge.
URL <http://dx.doi.org/10.1017/CB09780511623646>
- [12] Hwang, J.-M., 2006. Rigidity of rational homogeneous spaces. In: *International Congress of Mathematicians*. Vol. II. Eur. Math. Soc., Zürich, pp. 613–626.
- [13] Kollár, J., 1996. Rational curves on algebraic varieties. Vol. 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin.
URL <http://dx.doi.org/10.1007/978-3-662-03276-3>
- [14] Landsberg, J. M., Manivel, L., 2003. On the projective geometry of rational homogeneous varieties. *Comment. Math. Helv.* 78 (1), 65–100.
URL <http://dx.doi.org/10.1007/s000140300003>
- [15] Lazarsfeld, R., 2004. Positivity in algebraic geometry. I. Vol. 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, classical setting: line bundles and linear series.
URL <http://dx.doi.org/10.1007/978-3-642-18808-4>
- [16] Mok, N., 2008. Recognizing certain rational homogeneous manifolds of Picard number 1 from their varieties of minimal rational tangents. In: *Third International Congress of Chinese Mathematicians. Part 1, 2*. Vol. 2 of *AMS/IP Stud. Adv. Math.*, 42, pt. 1. Amer. Math. Soc., Providence, RI, pp. 41–61.
- [17] Muñoz, R., Occhetta, G., Solá Conde, L. E., Watanabe, K., 2015. Rational curves, Dynkin diagrams and Fano manifolds with nef tangent bundle. *Math. Ann.* 361 (3), 583–609.
- [18] Muñoz, R., Occhetta, G., Solá Conde, L. E., Watanabe, K., Wiśniewski, J. A., 2015. A survey on the Campana-Peternell conjecture. In: *Rendiconti dell’Istituto di Matematica dell’Università di Trieste. An International Journal of Mathematics*. Vol. 47. EUT Edizioni Università di Trieste.
- [19] Occhetta, G., Solá Conde, L. E., Watanabe, K., Wiśniewski, J. A., 2014. Fano manifolds whose elementary contractions are smooth \mathbb{P}^1 -fibrations, preprint [arXiv:1407.3658](https://arxiv.org/abs/1407.3658).
- [20] Okonek, C., Schneider, M., Spindler, H., 1980. Vector bundles on complex projective spaces. *Progress in Mathematics*, 3. Birkhäuser, Boston, Mass.
- [21] Serre, J.-P., 1955–1956. Géométrie algébrique et géométrie analytique. *Ann. Inst. Fourier, Grenoble* 6, 1–42.