

Draft Genome Sequences of Novel *Pseudomonas*, *Flavobacterium*, and *Sediminibacterium* Species Strains from a Freshwater Ecosystem

gen@meAnnouncements™

Federica Pinto,^a Adrian Tett,^a Federica Armanini,^a Francesco Asnicar,^a Adriano Boscaini,^b Edoardo Pasolli,^a Moreno Zolfo,^a Claudio Donati,^b Nico Salmaso,^b ^(b) Nicola Segata^a

^aCentre for Integrative Biology, University of Trento, Trento, Italy ^bEdmund Mach Foundation, San Michele all'Adige, Italy

AMERICAN SOCIETY FOR MICROBIOLOGY

ABSTRACT Freshwater ecosystems represent 0.01% of the water on Earth, but they support 6% of global biodiversity that is still mostly uncharacterized. Here, we describe the genome sequences of three strains belonging to novel species in the *Pseudomonas, Flavobacterium*, and *Sediminibacterium* genera recovered from a water sample of Lake Garda, Italy.

The uncharacterized microbial genetic diversity in natural environments is immense. Although high-throughput sequencing methods (1) and metagenomics (2) are recovering part of this diversity, ecosystems, such as freshwater habitats, which are estimated to support 6% of global biodiversity, remain poorly characterized (3). It is thus important to continue uncovering and analyzing microbial sequences from these environments.

In this project, we recovered and described three new microbial genomes of organisms inhabiting a pre-Alpine freshwater lake (Lake Garda, Italy). We specifically focused on potential symbionts of cyanobacterial organisms, and the genomic DNA was obtained from a nonaxenic culture of *Tychonema bourrellyi* (4). Paired-end libraries (Illumina) were prepared and run on the Illumina HiSeq 2500 platform (100-nucleotide [nt]-long paired-end reads). Raw reads were assembled using metaSPAdes version 3.10.1 (5) with default parameters, which generated 7,029 contigs larger than 1,000 bp with a total size of 74 Mbp. From this metagenomic assembly, three genomes belonging to the *Pseudomonas, Flavobacterium*, and *Sediminibacterium* genera were binned using the manually supervised anvi'o protocol based on the abundance and tetranucle-otide frequency distributions (6) and quality controlled by CheckM (7) (100%, 99%, and 99% predicted completeness, respectively, with an indication of contamination only for *Pseudomonas* at 0.27% and no strain heterogeneity).

The *Pseudomonas* sp. strain FEMGT703P genome has 4.41 Mb assembled into 14 contigs, with an N_{50} of 398,350 bp and a GC content of 59.93%. The assembly of the *Flavobacterium* sp. strain FEMGT703F genome resulted in 2.98 Mb, with an N_{50} of 372,530 bp and a GC content of 38.68%. The *Sediminibacterium* sp. strain FEMGT703S genome has 3.22 Mb, with an N_{50} of 530,195 bases and a GC content of 35.78%. The genomes were annotated using Prokka version 1.11 (8), which identified 4,064, 2,649, and 2,877 coding sequences for the *Pseudomonas*, *Flavobacterium*, and *Sediminibacterium* genomes, respectively. High-resolution phylogenetic profiling using PhyloPhIAn (9) and sequence similarity analysis using pyani (version 0.2.6; option "-m ANIb") both confirmed that the three genomes belong to new species when using a sequence identity cutoff of 95%.

The discovery of these new three microbial genomes in a nonaxenic *T. bourrellyi* culture might confirm the establishment of an ecological association between cyano-

Received 4 January 2018 Accepted 5 January 2018 Published 1 February 2018

Citation Pinto F, Tett A, Armanini F, Asnicar F, Boscaini A, Pasolli E, Zolfo M, Donati C, Salmaso N, Segata N. 2018. Draft genome sequences of novel *Pseudomonas, Flavobacterium*, and *Sediminibacterium* species strains from a freshwater ecosystem. Genome Announc 6: e00009-18. https://doi.org/10.1128/genomeA .00009-18.

Copyright © 2018 Pinto et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Nico Salmaso, nico.salmaso@fmach.it, or Nicola Segata, nicola.segata@unitn.it. bacteria and heterotrophic microbes (10, 11). However, more investigations are still needed to further characterize the microbial community diversity and interactions in freshwater systems.

Accession number(s). The sequences for this genome project have been deposited in GenBank under the accession no. PGCM00000000 (*Pseudomonas* sp. FEMGT703P), PGCN00000000 (*Flavobacterium* sp. FEMGT703F), and PGC00000000 (*Sediminibacterium* sp. FEMGT703S). The versions described in this paper are the first versions, PGCM01000000, PGCN01000000, and PGCO01000000, respectively.

ACKNOWLEDGMENTS

This work was financially supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement no. 704603 to Federica Pinto and Marie Skłodowska-Curie grant agreement no. 707345 to Edoardo Pasolli and by the European Research Council (ERC-STG project MetaPG) to Nicola Segata.

REFERENCES

- Hahn AS, Konwar KM, Louca S, Hanson NW, Hallam SJ. 2016. The information science of microbial ecology. Curr Opin Microbiol 31: 209–216. https://doi.org/10.1016/j.mib.2016.04.014.
- Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. 2017. Shotgun metagenomics, from sampling to analysis. Nat Biotechnol 35:833–844. https://doi.org/10.1038/nbt.3935.
- Dudgeon D, Arthington AH, Gessner MO, Kawabata ZI, Knowler DJ, Lévêque C, Naiman RJ, Prieur-Richard AH, Soto D, Stiassny MLJ, Sullivan CA. 2006. Freshwater biodiversity: importance, threats, status and conservation challenges. Biol Rev 81:163–182. https://doi.org/10 .1017/S1464793105006950.
- Pinto F, Tett A, Armanini F, Asnicar F, Boscaini A, Pasolli E, Zolfo M, Donati C, Salmaso N, Segata N. 2017. Draft genome sequence of the planktic cyanobacterium *Tychonema bourrellyi*, isolated from alpine lentic freshwater. Genome Announc 5:e01294-17. https://doi.org/10.1128/ genomeA.01294-17.
- Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile metagenomic assembler. Genome Res 27:824–834. https://doi .org/10.1101/gr.213959.116.

- Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319. https://doi.org/10.7717/peerj.1319.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res 25:1043–1055. https:// doi.org/10.1101/gr.186072.114.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069. https://doi.org/10.1093/bioinformatics/btu153.
- Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method for improved phylogenetic and taxonomic placement of microbes. Nat Commun 4:2304. https://doi.org/10.1038/ncomms3304.
- Alvarenga DO, Fiore MF, Varani AM. 2017. A metagenomic approach to cyanobacterial genomics. Front Microbiol 8:809. https://doi.org/10.3389/ fmicb.2017.00809.
- Seymour JR, Amin SA, Raina JB, Stocker R. 2017. Zooming in on the phycosphere: the ecological interface for phytoplankton-bacteria relationships. Nat Microbiol 2:17065. https://doi.org/10.1038/nmicrobiol .2017.65.





Correction for Pinto et al., "Draft Genome Sequences of Novel *Pseudomonas, Flavobacterium*, and *Sediminibacterium* Strains from a Freshwater Ecosystem"

Federica Pinto,^a Adrian Tett,^a Federica Armanini,^a Francesco Asnicar,^a Adriano Boscaini,^b Edoardo Pasolli,^a Moreno Zolfo,^a Claudio Donati,^b Nico Salmaso,^b ^(b) Nicola Segata^a

^aCentre for Integrative Biology, University of Trento, Trento, Italy ^bEdmund Mach Foundation, San Michele all'Adige, Italy

Volume 6, no. 5, e00009-18, 2018, https://doi.org/10.1128/genomeA.00009-18. Page 1: The article title should read as given above.

Published 8 March 2018

Citation Pinto F, Tett A, Armanini F, Asnicar F, Boscaini A, Pasolli E, Zolfo M, Donati C, Salmaso N, Segata N. 2018. Correction for Pinto et al., "Draft genome sequences of novel *Pseudomonas, Flavobacterium*, and *Sediminibacterium* strains from a freshwater ecosystem." Genome Announc 6:e00169-18. https://doi.org/10.1128/genomeA.00169-18.

Copyright © 2018 Pinto et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.