


Top-down Attention in the Face-Processing Network: an MRI-guided MEG Study using Multiple Simultaneous Frequency Tags

Daniel Baldauf & Eelke de Vries

Neuroimaging and electrophysiology have revealed multiple cortical face-selective regions that are spatially and functionally separable and form a distributed cortical network specialized for face perception. The aim of the present study was to investigate this functional specialization of the areas OFA, FFA, and STS by demonstrating the differential entrainment with the respective tagging-frequencies embedded in a face compound stimulus, as well as to investigate the attentional modulation of the related neural activations at those specific frequencies. For this, we used a frequency-tagging paradigm that allowed for independently frequency-tagging various parts of a face (eyes, mouth) as well as rhythmic changes in facial identity.



Top-down attention to any one of the three face components (face identity, face parts, and eye gaze) was found to facilitate both the neural activity in the respective sub-network related to processing of the attended stimulus and the behavioral performance in a target detection task.

We first identified the respective network components (ROIs) functionally with independent localizers. These functional localizers were also based on the MEG recordings, but instead of analyzing the rhythmic modulation of the signal during the complete, highly repetitive stimulus period, the functional localizers were solely based on the evoked magnetic fields (ERFs) in response to very first appearance of a face stimulus at the beginning of each trial. All regions of interest were defined in each subject's individual MRI space, which was co-registered with the MEG Squid array to guide the reconstruction of neural sources within their individual anatomical frame of reference.




Figure 2. (A) Behavioral responses were significantly faster and more accurate for validly cued targets compared to invalidly cued targets. (B) Mean coordinates of the FFA (red), OFA (green), and STS (blue), as defined in each experimental participant of the present study. (C-D) Entrainments of the three ROIs with the respective frequency-tags of the respectively preferred face-aspect. (F-H) Mean MEG power spectra of the Minimum-norm-estimates (MNE) for each ROI and attention condition.

The following spectroanalyses of the time-series in all three regions of interest of the face-processing network showed that it is possible to entrain several functionally specialized neural populations with the presentation rhythms of respectively relevant information.

By investigating – for the first time in MEG - the temporal dynamics of the complete occipital face processing network (OFA, FFA, and STS), we were able to dissect the functional compartmentalization of the system and the functional specialization of its components by demonstrating selective attentional modulation in each of the three regions.

In general, our MEG results are consistent with the view of an hierarchical organization of the three sub-networks : The network populations at an earlier level, such as OFA, and in parts also STS, are preferentially concerned with the analysis of crucial features or facial sub-components and seem to feed those representations of object parts forward to higher level face processing networks in inferior-temporal cortex, e.g. FFA, where the information about facial features and sub-components is integrated into a representation of facial identity. This relative arrangement in a hierarchical face-processing network is also supported by the telling differences in response latencies.

References:

- Bagherzadeh, Y., Baldauf, D., Lu, B., Pantazis, D., & Desimone, R. (2017). Alpha and gamma neurofeedback reinforce control of spatial attention. *Journal of Vision* 17 (10), 385-385
- Baldauf, D. Wolf, M., & Deubel, H. (2006), Deployment of visual attention before sequences of goal-directed hand movements. *Vision Research*, 46(26), 4355-4374.
- Baldauf, D., & Deubel, H. (2008a). Properties of attentional selection during the preparation of sequential saccades. *Experimental Brain Research*, 184(3), 411-425.
- Baldauf, D., & Deubel, H. (2008b). Visual attention during the preparation of bimanual movements. *Vision Research*, 48, 549-563.
- Baldauf, D., Cui, H., & Andersen, R.A. (2008a). The posterior parietal cortex encodes the first and second goal in sequential reaching arm movements. Annual meeting of the Society for Neuroscience, 2007. Program No. 397.3. San Diego, CA: Online.
- Baldauf, D., Cui, H., & Andersen, R.A. (2008b). The posterior parietal cortex encodes in parallel both goals for double-reach sequences. *Journal of Neuroscience*, 28(40):10081-10089.
- Baldauf, D., & Deubel, H. (2009). Attentional selection of multiple movement goal positions before rapid hand movement sequences: an ERP study. *Journal of Cognitive Neuroscience*, 21(1), 18-29.
- Baldauf, D., & Deubel, H. (2010). Attentional landscapes in reaching and grasping. *Vision Research*, 50, 999-1013.
- Baldauf, D. (2011). Chunking movements into sequence: The visual pre-selection of subsequent goals. *Neuropsychologia*, 49(5), 1383-1387.
- Baldauf, D. & Desimone, R. (2014). Neural mechanisms of object-based attention. *Science*, 344, 424-427.
- Baldauf, D. (2015). Top-down biasing signals of non-spatial, object-based attention. *Journal of Vision*, 15(12), 1395-1395.
- Baldauf, D., & Desimone, R. (2016). Mechanisms of spatial versus non-spatial, modality-based attention. Annual meeting of the Society for Neuroscience, 2016. Program no. 531.27. San Diego.
- Baldauf, D., Grossman, N., Hu, A.M., Boyden, E. & Desimone, R. (2016). Transcranial alternating current stimulation(tACS) reveals causal role of brain oscillations in visual attention. *Journal of Vision* 16 (12), 937-937