

A TEST FOR THE DIVERGENCE OF CERTAIN SERIES WITH POSITIVE TERMS

LUCA GOLDONI

ABSTRACT. In this short note we prove a simple divergent test for certain series with positive terms which are related in some way with the Theory of Dirichlet's series.

1. INTRODUCTION

While I was preparing some examples for ordinary Dirichlet series I encountered the following problem: I need to show that the series

$$\sum_{n=1}^{\infty} \frac{|\sin n|}{n}$$

is divergent. Since the method I used can be somewhat generalized it seems to me worthwhile to write down this short note.

2. THE THEOREM

Theorem 1. *Let be*

$$(1) \quad \sum_{n=1}^{\infty} \frac{|f(n)|}{n}.$$

if

- (1) $f : \mathbb{R} \rightarrow \mathbb{R}$ is a bounded function.
- (2) There exists a real number c and two bounded functions $g : \mathbb{R} \rightarrow \mathbb{R}$, $h : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$0 < f(c) \leq f(n+1)g(n) + h(n)f(n) \quad \forall n \in \mathbb{N}.$$

then the series (1) diverges.

Proof. Since the functions f , g , h are bounded on \mathbb{R} we can choose a positive constant M so that, for each $n \in \mathbb{N}$ it is $|f(n)| < M$, $|g(n)| < M$, $|h(n)| < M$. From the triangular inequality we have

$$0 < |f(c)| \leq |f(n+1)| |g(n)| + |f(n)| |h(n)| \leq M (|f(n+1)| + |f(n)|).$$

Date: May 13, 2018.

2000 Mathematics Subject Classification. 40A05, 30B50.

Key words and phrases. Infinite series, Divergence tests, Elementary proof.

Dipartimento di Matematica. Università di Trento.

so that

$$(|f(n+1)| + |f(n)|) \geq \frac{M}{f(c)}.$$

Now, since

$$\frac{|f(n)|}{n} = \left(\frac{|f(n)| - |f(n+1)|}{2n} \right) + \left(\frac{|f(n)| + |f(n+1)|}{2n} \right).$$

it follows that

$$\frac{|f(n)|}{n} \geq \left(\frac{|f(n)| - |f(n+1)|}{2n} \right) + \frac{M}{2f(c)} \frac{1}{n}.$$

Since, for each $k \in \mathbb{N}$ it is

$$S_k = \sum_{n=1}^k (|f(n)| - |f(n+1)|) = |f(1)| - |f(k+1)|$$

we have that

$$|S_k| \leq |f(1)| + M < +\infty$$

and so the series

$$\sum_{n=1}^{+\infty} \frac{|f(n)| - |f(n+1)|}{2n}$$

converges by Dirichlet's convergence test. Thus

$$\lim_{m \rightarrow \infty} \sum_{n=1}^m \frac{|f(n)|}{n} \geq \lim_{m \rightarrow \infty} \sum_{n=1}^m \frac{|f(n)| - |f(n+1)|}{2n} + \lim_{m \rightarrow \infty} \frac{M}{2f(c)} \sum_{n=1}^m \frac{1}{n} = +\infty$$

and the given series is divergent. \square

Corollary 1. *The series*

$$\sum_{n=1}^{\infty} \frac{|\sin(n)|}{n}, \quad \sum_{n=1}^{\infty} \frac{|\cos(n)|}{n}$$

are divergent.

Proof. It is enough to observe that for the first series we have

$$0 < \sin(1) = \sin(n+1-n) = \sin(n+1)\cos n - \cos(n+1)\sin n$$

so that

- $c = 1$.
- $g(n) = \cos n$ and $h(n) = -\cos(n+1)$.

while, for the second, it is

$$0 < \cos\left(\frac{\pi}{2} - 1\right) = \sin(1) = \sin(n+1)\cos n - \cos(n+1)\sin n$$

so that

- $c = \frac{\pi}{2} - 1$.
- $g(n) = \sin(n+1)$, $h(n) = -\sin(n)$.

\square

Corollary 2. *The Dirichlet's series*

$$\sum_{n=1}^{\infty} \frac{|\sin n|}{n^s}, \quad \sum_{n=1}^{\infty} \frac{|\cos n|}{n^s}$$

have abscissa of convergence $\sigma_c = 1$.

REFERENCES

- [1] G.H. Hardy "A course in Pure Mathematics" Cambridge University Press, 2004.
- [2] C.J. Tranter "Techniques of Mathematical analysis" Hodder and Stoughton, 1976.

UNIVERSITÀ DI TRENTO, DIPARTIMENTO DI MATEMATICA, V. SOMMARIVE
14, 56100 TRENTO, ITALY

E-mail address: goldoni@science.unitn.it