

STABLE REGULAR CRITICAL POINTS OF THE MUMFORD-SHAH FUNCTIONAL ARE LOCAL MINIMIZERS

M. BONACINI AND M. MORINI

ABSTRACT. In this paper it is shown that any regular critical point of the Mumford-Shah functional, with positive definite second variation, is an isolated local minimizer with respect to competitors which are sufficiently close in the L^1 -topology. A global minimality result in small tubular neighborhoods of the discontinuity set is also established.

1. INTRODUCTION

The *Mumford-Shah functional* is the most typical example of a class of variational problems called by E. De Giorgi *free discontinuity problems*, characterized by the competition between volume and surface terms. The minimization of such an energy was proposed in the seminal papers [22, 23] in the context of image segmentation, and plays an important role also in variational models for fracture mechanics. Its homogeneous version in a bounded open set $\Omega \subset \mathbb{R}^2$ is defined over pairs (Γ, u) , with Γ closed subset of $\bar{\Omega}$ and $u \in H^1(\Omega \setminus \Gamma)$, as

$$F(\Gamma, u) := \int_{\Omega \setminus \Gamma} |\nabla u|^2 dx + \mathcal{H}^1(\Gamma \cap \Omega). \quad (1.1)$$

Since its introduction, several results concerning the existence and regularity of minimizers, as well as the structure of the optimal set, have been obtained (see, *e.g.*, [2] for a detailed account on this topic).

In this paper we continue the study of second order optimality conditions for the functional in (1.1) initiated by F. Cagnetti, M.G. Mora and the second author in [6], where a suitable notion of second variation was introduced by considering one-parameter families of perturbations of the regular part of the discontinuity set. In [6] it was also shown that a critical point (Γ, u) with positive definite second variation minimizes the functional with respect to pairs of the form $(\Phi(\Gamma), v)$, where Φ is any diffeomorphism sufficiently close to the identity in the C^2 -norm, with $\Phi - Id$ compactly supported in Ω , and $v \in H^1(\Omega \setminus \Phi(\Gamma))$ satisfies $v = u$ on $\partial\Omega$.

In the main theorem of this paper we strongly improve the aforementioned result, by showing that in fact the positive definiteness of the second variation implies strict local minimality with respect to the weakest topology which is natural for this problem, namely the L^1 -topology. To be more precise, we prove that if (Γ, u) is a critical point with positive second variation, then there exists $\delta > 0$ such that

$$F(\Gamma, u) < F(K, v)$$

for all admissible pairs (K, v) , provided that v attains the same boundary conditions as u and $0 < \|u - v\|_{L^1(\Omega)} < \delta$. We mention that for technical reasons the boundary conditions imposed here are slightly different from those considered in [6], as we prescribe the Dirichlet condition only on a portion $\partial_D \Omega \subset \partial\Omega$ away from the intersection of the discontinuity set Γ with $\partial\Omega$.

The general strategy of the proof is close in spirit to the one devised in [12] for a different free-discontinuity problem. It consists in two fundamental steps: first, one shows that strict stability is sufficient to guarantee local minimality with respect to perturbations of the discontinuity set which

2010 *Mathematics Subject Classification.* 49K10 (49Q20).

Key words and phrases. Mumford-Shah functional, free discontinuity problems, second variation.

This is a post-peer-review, pre-copyedit version of an article published in *Annales de l'Institut Henry Poincaré C, Analyse non linéaire*. The final authenticated version is available online at: <http://dx.doi.org/10.1016/j.anihpc.2014.01.006>.

are close to the identity in the $W^{2,\infty}$ -norm (see Theorem 5.2). This amounts to adapting to our slightly different context the techniques developed in [6], with the main new technical difficulties stemming from allowing also boundary variations of the discontinuity set.

The second step of the outline consists in showing that the above local $W^{2,\infty}$ -minimality in fact implies the claimed local L^1 -minimality. This is done through a penalization/regularization approach, with an appeal to the regularity theory of quasi-minimizers of the area functional and of the Mumford-Shah functional (see [2]). More precisely, we start by showing that the local $W^{2,\infty}$ -minimality implies minimality with respect to small $C^{1,\alpha}$ -perturbations of the discontinuity set. This is perhaps the most technical part of the proof. The main idea is to restrict F to the class of pairs (Γ, v) such that $\|v - u\|_{W^{1,\infty}(\Omega \setminus \Gamma)} \leq 1$, so that the Dirichlet energy behaves like a volume term, and F can be regarded as a volume perturbation of the area functional. This allows to use the regularity theory for quasi-minimizers of the area functional to deduce the local $C^{1,\alpha}$ -minimality through a suitable contradiction argument.

A contradiction argument is also finally used to establish the sought L^1 -minimality. To give a flavor of this type of reasoning, we sketch here the main steps of this last part of the proof. One assumes by contradiction the existence of admissible pairs (Γ_n, u_n) with u_n converging to u in $L^1(\Omega)$, such that the minimality inequality fails along the sequence:

$$F(\Gamma_n, u_n) \leq F(\Gamma, u) \quad (1.2)$$

for every n . By an easy truncation argument, we may also assume that $\|u_n\|_\infty \leq \|u\|_\infty$, so that $u_n \rightarrow u$ in $L^p(\Omega)$ for every $p \geq 1$. Then we replace each (Γ_n, u_n) by a new pair (K_n, v_n) chosen as solution to a suitable penalization problem, namely

$$\min \left\{ F(K, w) + \beta \left(\sqrt{(\|w - u\|_{L^2(\Omega)}^2 - \varepsilon_n)^2 + \varepsilon_n^2} - \varepsilon_n \right) : (K, w) \text{ admissible, } w = u \text{ on } \partial_D \Omega \right\},$$

with $\varepsilon_n := \|u_n - u\|_{L^2(\Omega)}^2 \rightarrow 0$, and $\beta > 0$ large enough. Note that, by (1.2) and by minimality, we have

$$F(K_n, v_n) \leq F(\Gamma_n, u_n) \leq F(\Gamma, u). \quad (1.3)$$

The advantage is now that the pairs (K_n, v_n) satisfy a uniform *quasi-minimality* property (see Theorem 2.2). It is easy to show that, up to subsequences, the sequence (K_n, v_n) converges to a minimizer of the limiting problem

$$\min \left\{ F(K, w) + \beta \|w - u\|_{L^2(\Omega)}^2 : (K, w) \text{ admissible, } w = u \text{ on } \partial_D \Omega \right\}. \quad (1.4)$$

Now a calibration argument developed in [21] implies that we may choose β so large that (Γ, u) is the unique global minimizer of (1.4). With this choice of β we have in particular that $v_n \rightarrow u$ in L^1 , and in turn, by exploiting the regularity properties of quasi-minimizers of the Mumford-Shah functional, we infer that the corresponding discontinuity sets K_n are locally $C^{1,\alpha}$ -graphs and converge in the $C^{1,\alpha}$ -sense to Γ . Recalling (1.3), we have reached a contradiction to the $C^{1,\alpha}$ -minimality.

We remark that a similar two-steps strategy has been used also in [1] for a nonlocal isoperimetric problem related to the modeling of diblock copolymers, and in [8], where the appeal to the regularity of quasi-minimizers appears for the first time in the context of isoperimetric inequalities.

We regard our result as a first step of a more general study of second order minimality conditions for free-discontinuity problems. Besides considering more general functionals, it would be very interesting to extend our local minimality criterion to the case of discontinuity sets with singular points, like the so-called ‘‘triple junction’’, where three lines meet forming equal angles of $2\pi/3$, and the ‘‘crack-tip’’, where a line terminates at some point. This will be the subject of future investigation.

The paper is organized as follows. In Section 2 we fix the notation and we review some preliminary results concerning the regularity theory for quasi-minimizers of the Mumford-Shah functional. In Section 3 we collect the necessary definitions and state the main result. Section 4 is devoted to the computation of the second variation, when also boundary variations of the discontinuity set are allowed. The proof of the main theorem starts in Section 5 (where the local $W^{2,\infty}$ -minimality is addressed) and lasts for Sections 6 and 7 (where the $C^{1,\alpha}$ and the desired local L^1 -minimality,

respectively, are established). In Section 8 we describe some examples and applications of our minimality criterion. We highlight here Proposition 8.1 where we show that a regular critical point is always a global minimizer in sufficiently small tubular neighborhoods of the discontinuity set. We thus recover in our framework the result of [19], which was obtained by a calibration method.

In the final Appendix (Section 9) we prove some auxiliary technical lemmas needed in the paper.

2. NOTATION AND PRELIMINARIES

In this section we fix the notation and we recall some preliminary results.

2.1. Geometric preliminaries. Let Γ be a smooth embedded curve in \mathbb{R}^2 , let $\nu : \mathcal{U} \rightarrow \mathbb{S}^1$ be a smooth vector field defined in a tubular neighborhood \mathcal{U} of Γ and normal to Γ on Γ , and let $\tau := \nu^\perp$ be the unit tangent vector to Γ (where $^\perp$ stands for the clockwise rotation by $\frac{\pi}{2}$). If $g : \mathcal{U} \rightarrow \mathbb{R}^d$ is a smooth function, we denote by $D_\Gamma g(x)$ ($\nabla_\Gamma g(x)$ if $d = 1$) the *tangential differential* of g at $x \in \Gamma$, that is, the linear operator from \mathbb{R}^2 into \mathbb{R}^d given by $D_\Gamma g(x) := dg(x) \circ \pi_x$, where $dg(x)$ is the usual differential of g at x and π_x is the orthogonal projection on the tangent space to Γ at x . If g is a vector field from Γ to \mathbb{R}^2 we define also its *tangential divergence* as $\operatorname{div}_\Gamma g := \tau \cdot \partial_\tau g$.

The following *divergence formula* is a particular case of [24, 7.6]: for every smooth vector field $g : \mathcal{U} \rightarrow \mathbb{R}^2$ holds

$$\int_\Gamma \operatorname{div}_\Gamma g \, d\mathcal{H}^1 = \int_\Gamma H(g \cdot \nu) \, d\mathcal{H}^1 + \int_{\partial\Gamma} g \cdot \eta \, d\mathcal{H}^0. \quad (2.1)$$

Here $\partial\Gamma$ stands for the endpoints of Γ , η is a unit vector tangent to Γ and pointing out of Γ at each point of $\partial\Gamma$ (it coincides with τ , up to a sign) and the function H is defined in \mathcal{U} by $H := \operatorname{div} \nu$. Notice that H coincides, when restricted to Γ , with the *curvature* of Γ and, since $\partial_\nu \nu = 0$, we have $H = \operatorname{div}_\Gamma \nu = D\nu[\tau, \tau]$.

Let $\Phi : \bar{\mathcal{U}} \rightarrow \bar{\mathcal{U}}$ be a smooth orientation-preserving diffeomorphism and let $\Gamma_\Phi := \Phi(\Gamma)$. A possible choice for the unit normal to Γ_Φ is given by the vector field

$$\nu_\Phi = \frac{(D\Phi)^{-T}[\nu]}{|(D\Phi)^{-T}[\nu]|} \circ \Phi^{-1}, \quad (2.2)$$

while the vector η appearing in (2.1) becomes

$$\eta_\Phi = \frac{D\Phi[\eta]}{|D\Phi[\eta]|} \circ \Phi^{-1} \quad (2.3)$$

on $\partial\Gamma_\Phi$. We denote by H_Φ the curvature of Γ_Φ . We shall use the following identity, which is a particular case of the so-called generalized area formula (see, *e.g.*, [2, Theorem 2.91]): for every $\psi \in L^1(\Gamma_\Phi)$

$$\int_{\Gamma_\Phi} \psi \, d\mathcal{H}^1 = \int_\Gamma (\psi \circ \Phi) J_\Phi \, d\mathcal{H}^1, \quad (2.4)$$

where $J_\Phi := |(D\Phi)^{-T}[\nu]| \det D\Phi$ is the 1-dimensional Jacobian of Φ .

2.2. Partial regularity for quasi-minimizers of the Mumford-Shah functional. Given an open set $\Omega \subset \mathbb{R}^2$, we recall that the space $SBV(\Omega)$ of special functions of bounded variation is defined as the set of all functions $u \in L^1(\Omega)$ whose distributional derivative Du is a bounded Radon measure of the form

$$Du = \nabla u \mathcal{L}^2 + D^j u = \nabla u \mathcal{L}^2 + (u^+ - u^-) \nu_u \mathcal{H}^1 \llcorner S_u,$$

where $\nabla u \in L^1(\Omega; \mathbb{R}^2)$ is the approximate gradient of u , S_u is the jump set of u (which is countably $(\mathcal{H}^1, 1)$ -rectifiable), u^+ and u^- are the traces of u on S_u and ν_u is the approximate normal on S_u . We refer to [2] for a complete treatment of the space SBV and a precise definition of all the notions introduced above. In the sequel we will consider the following notion of convergence in the space SBV , motivated by the compactness theorem [2, Theorem 4.8].

Definition 2.1. We say that $u_n \rightarrow u$ in $SBV(\Omega)$ if $u_n \rightarrow u$ strongly in $L^1(\Omega)$, $\nabla u_n \rightharpoonup \nabla u$ weakly in $L^2(\Omega; \mathbb{R}^2)$, and $D^j u_n \rightharpoonup D^j u$ weakly* in the sense of measures in Ω .

Given $u \in SBV(\Omega)$, we introduce the quantities

$$D_u(x, r) := \int_{B_r(x) \cap \Omega} |\nabla u|^2 dy, \quad A_u(x, r) := \min_{T \in \mathcal{A}} \int_{\overline{S}_u \cap B_r(x)} \text{dist}^2(y, T) d\mathcal{H}^1(y),$$

where \mathcal{A} denotes the set of affine lines in \mathbb{R}^2 , and

$$E_u(x, r) := D_u(x, r) + r^{-2} A_u(x, r).$$

The result that we are going to recall expresses the fact that the rate of decay of E_u in small balls determines the $C^{1,\alpha}$ -regularity of the jump set of u , provided that u satisfies a quasi-minimality property. In order to state precisely the theorem, we introduce some more notation. We set $C_{\nu,r} := \{x \in \mathbb{R}^2 : |\pi_\nu(x)| < r, |x \cdot \nu| < r\}$ for $\nu \in \mathbb{S}^1$ and $r > 0$, where $\pi_\nu(x) = x - (x \cdot \nu)\nu$. If $g : (-r, r) \rightarrow \mathbb{R}$, we define the *graph* of g (with respect to the direction ν) to be the set

$$\text{gr}_\nu(g) := \{x = x' + g(x')\nu \in \mathbb{R}^2 : x' = \pi_\nu(x), |x'| < r\}.$$

Theorem 2.2. *Let $u \in SBV(\Omega)$ be a quasi-minimizer of the Mumford-Shah functional, that is, assume that there exists $\omega > 0$ such that for every ball $B_\rho(x)$*

$$\int_{\Omega \cap B_\rho(x)} |\nabla u|^2 dx + \mathcal{H}^1(S_u \cap B_\rho(x)) \leq \int_{\Omega \cap B_\rho(x)} |\nabla v|^2 dx + \mathcal{H}^1(S_v \cap B_\rho(x)) + \omega \rho^2 \quad (2.5)$$

for every $v \in SBV(\Omega)$ with $\{v \neq u\} \subset\subset B_\rho(x)$. There exist constants $R_0 > 0$, $\varepsilon_0 > 0$ (depending only on ω) such that if

$$E_u(x, r) < \varepsilon_0 r$$

for some $x \in \overline{S}_u \cap \Omega$ and $r < R := R_0 \wedge \text{dist}(x, \partial\Omega)$, then there exist a smaller radius $r' \in (0, r)$ (depending only on ω , R and r) and a function $f \in C^{1, \frac{1}{4}}(-r', r')$ with $f(0) = f'(0) = 0$ such that

$$(\overline{S}_u - x) \cap C_{\nu, r'} = \text{gr}_\nu(f),$$

where ν denotes the normal to S_u at x . Moreover, $\|f\|_{C^{1, \frac{1}{4}}} \leq C$ for some constant C depending only on ω .

The previous result (which holds also in dimension $N > 2$) is a consequence of [2, Theorem 8.2 and Theorem 8.3]: the only missing point is the uniform bound in $C^{1, \frac{1}{4}}$, which is not explicitly stated but can be deduced by checking that the constants appearing in the proof depend only on ω . Notice that the theorem provides the regularity of S_u in balls well contained in Ω ; concerning the regularity of the discontinuity set at the intersection with the boundary of Ω , under Neumann conditions, we have the following result, which is essentially contained in the book [10] (see, in particular, [10, Remark 79.42]; see also [18]).

Theorem 2.3. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, open set with boundary of class C^1 , and let $u \in SBV(\Omega)$ satisfy the same assumption of Theorem 2.2. Then there exist $b \in (0, 1)$ and $\tau > 0$ (depending only on ω and on Ω) such that, setting*

$$\Omega(\tau) := \{x \in \Omega : \text{dist}(x, \partial\Omega) < \tau\},$$

the intersection $\overline{S}_u \cap \Omega(\tau)$ is a finite disjoint union of curves of class $C^{1,b}$ intersecting $\partial\Omega$ orthogonally, with $C^{1,b}$ -norm uniformly bounded by a constant depending only on ω and Ω .

We conclude this preliminary section by recalling a well known property of quasi-minimizers of the Mumford-Shah functional, namely a lower bound on the \mathcal{H}^1 -dimensional density of the jump set in balls centered at any point of its closure. The estimate was proved in [11] in balls entirely contained in the domain Ω (see also [2, Theorem 7.21]); we refer also, when a Dirichlet condition is assumed at the boundary of the domain, to [7] for balls centered at $\partial\Omega$, and to [3] for balls possibly intersecting $\partial\Omega$ but not necessarily centered at $\partial\Omega$, and finally to [10, Section 77] in the case of balls intersecting $\partial\Omega$ when a Neumann condition is imposed.

In fact, for our purposes we will need to consider the mixed situation, where we impose a Dirichlet condition on a part $\partial_D\Omega$ of the boundary and a Neumann condition on the remaining part $\partial_N\Omega$. The result is still valid in this case, for balls centered at the intersection between the Dirichlet and the Neumann part of the boundary, under the additional assumption that $\partial_D\Omega$ and

$\partial_N \Omega$ meet orthogonally. We are not aware of any result of this kind in the existing literature, but the proof can be obtained by following closely the strategy of the original proof in [11], combined also with some new ideas contained in [3]. We will sketch the proof in Section 9.1, referring the reader to [4] for the details.

The precise statement is the following.

Theorem 2.4. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, open set, let $\partial_D \Omega \subset \partial \Omega$ be relatively open and of class C^1 , $\partial_N \Omega := \partial \Omega \setminus \overline{\partial_D \Omega}$ of class C^1 , and assume that $\partial_D \Omega$ meets $\partial_N \Omega$ orthogonally. Let $\Omega' \subset \mathbb{R}^2$ be a bounded, open set of class C^1 such that $\Omega \subset \Omega'$ and $\partial \Omega \cap \Omega' = \partial_D \Omega$. Let $u \in SBV(\Omega')$ be such that $\overline{S_u} \cap \overline{\partial_D \Omega} = \emptyset$ and $u \in W^{1,\infty}(\Omega' \setminus \overline{S_u})$.*

Let $w \in SBV(\Omega')$, with $w = u$ in $\Omega' \setminus \Omega$, satisfy for every $x \in \overline{\Omega}$ and for every $\rho > 0$

$$\int_{\Omega' \cap B_\rho(x)} |\nabla w|^2 dx + \mathcal{H}^1(S_w \cap B_\rho(x)) \leq \int_{\Omega' \cap B_\rho(x)} |\nabla v|^2 dx + \mathcal{H}^1(S_v \cap B_\rho(x)) + \omega \rho^2$$

for every $v \in SBV(\Omega')$ such that $v = u$ in $\Omega' \setminus \Omega$ and $\{v \neq w\} \subset \subset B_\rho(x)$. Then there exist $\rho_0 > 0$ and $\theta_0 > 0$ (depending only on ω , u and Ω) such that

$$\mathcal{H}^1(S_w \cap B_\rho(x)) \geq \theta_0 \rho$$

for every $\rho \leq \rho_0$ and $x \in \overline{S_w}$.

3. SETTING AND MAIN RESULT

Let $\Omega \subset \mathbb{R}^2$ be an open, bounded, connected set with boundary of class C^3 . We introduce the following space of *admissible pairs*

$$\mathcal{A}(\Omega) := \{(K, v) : K \subset \mathbb{R}^2 \text{ closed, } v \in H^1(\Omega \setminus K)\}$$

and we set

$$F(K, v) := \int_{\Omega \setminus K} |\nabla v|^2 dx + \mathcal{H}^1(K \cap \Omega) \quad \text{for } (K, v) \in \mathcal{A}(\Omega).$$

It will be useful to consider also a localized version of the functional: for $A \subset \Omega$ open we set

$$F((K, v); A) := \int_{A \setminus K} |\nabla v|^2 dx + \mathcal{H}^1(K \cap A).$$

Given an admissible pair $(K, v) \in \mathcal{A}(\Omega)$ and assuming that K is a regular curve connecting two points of $\partial \Omega$, we denote by ν a smooth vector field coinciding with the unit normal to K when restricted to the points of K , by H the curvature of K with respect to ν , and by η the unit co-normal of $K \cap \partial \Omega$ (see Section 2.1). For any function $z \in H^1(\Omega \setminus K)$ we denote the traces of z on the two sides of K by z^+ and z^- : precisely, for \mathcal{H}^1 -a.e. $x \in K$ we set

$$z^\pm(x) := \lim_{r \rightarrow 0^+} \frac{1}{|B_r(x) \cap V_x^\pm|} \int_{B_r(x) \cap V_x^\pm} z(y) dy,$$

where $V_x^\pm := \{y \in \mathbb{R}^2 : \pm(y-x) \cdot \nu(x) \geq 0\}$. With an abuse of notation, we denote by z^+ and z^- also the restrictions of z to Ω^+ and Ω^- respectively, where Ω^+ and Ω^- are the two connected components of $\Omega \setminus K$, with the normal vector field ν pointing into Ω^+ . Finally we denote by $\nu_{\partial \Omega}$ the exterior unit normal vector to $\partial \Omega$ and by $H_{\partial \Omega}$ the curvature of $\partial \Omega$ with respect to $\nu_{\partial \Omega}$.

Definition 3.1. We say that $(K, v) \in \mathcal{A}(\Omega)$ is a *regular pair* if K is a curve of class C^∞ connecting two points of $\partial \Omega$ and intersecting $\partial \Omega$ transversally, and there exists $\partial_D \Omega \subset \subset \partial \Omega \setminus K$ relatively open in $\partial \Omega$ such that v is of class C^1 on $\partial_D \Omega$ and

$$\int_{\Omega \setminus K} \nabla v \cdot \nabla z dx = 0 \quad \text{for every } z \in H^1(\Omega \setminus K) \text{ with } z = 0 \text{ on } \partial_D \Omega, \quad (3.1)$$

that is, v is a weak solution to

$$\begin{cases} \Delta v = 0 & \text{in } \Omega \setminus K, \\ \partial_\nu v^\pm = 0 & \text{on } K \cap \Omega, \\ \partial_{\nu_{\partial \Omega}} v = 0 & \text{on } \partial_N \Omega := \partial \Omega \setminus \partial_D \Omega. \end{cases}$$

We denote by $\mathcal{A}_{reg}(\Omega)$ the space of all such pairs.




FIGURE 1. An admissible subdomain U for a regular pair (K, v) (see Definition 3.2). Notice that U excludes the relative boundary of $\partial_D \Omega$.

Definition 3.2. Given a regular pair $(K, v) \in \mathcal{A}_{reg}(\Omega)$, we say that an open subset $U \subset \mathbb{R}^2$ with Lipschitz boundary is an *admissible subdomain* if $K \subset U$ and $\overline{U} \cap \mathcal{S} = \emptyset$, where \mathcal{S} denotes the relative boundary of $\partial_D \Omega$ in $\partial \Omega$. In this case we define the space $H_V^1(\Omega \setminus K)$ consisting of all functions $v \in H^1(\Omega \setminus K)$ such that $v = 0$ in $(\Omega \setminus U) \cup \partial_D \Omega$ (the condition on $\partial_D \Omega$ has to be intended in the sense of traces). Notice that equation (3.1) holds for every $z \in H_V^1(\Omega \setminus K)$.

This paper deals with regular critical pairs (Γ, u) , according to the following definition motivated by the formula for the first variation of the functional F (see (4.6) and Remark 4.8).

Definition 3.3. We say that a regular pair $(\Gamma, u) \in \mathcal{A}_{reg}(\Omega)$ is a *regular critical pair* for F if the following conditions are satisfied:

- (i) Γ meets $\partial \Omega$ orthogonally,
- (ii) *transmission condition*:

$$H = |\nabla_{\Gamma} u^+|^2 - |\nabla_{\Gamma} u^-|^2 \quad \text{on } \Gamma \cap \Omega, \quad (3.2)$$

- (iii) *non-vanishing jump condition*: $|u^+ - u^-| \geq c > 0$ on Γ .

Remark 3.4. The assumption of C^∞ -regularity of the curve Γ is not so restrictive as it may appear: indeed, as a consequence of the transmission condition (3.2) and of the fact that u satisfies (3.1), Γ is automatically analytical as soon as it is of class $C^{1,\alpha}$ (see [16]). Moreover, by (3.1) u is of class C^∞ up to $\Gamma \cap \Omega$ and the traces ∇u^+ , ∇u^- of ∇u are well defined on both sides of Γ . Notice that u may fail to be regular only at the relative boundary \mathcal{S} of $\partial_D \Omega$; since every admissible subdomain U excludes a neighborhood of \mathcal{S} , the values of u in such a neighborhood will not affect any of the arguments of the paper. Thus, we can always assume (by modifying u in a neighborhood of \mathcal{S} , if needed) that u^+ and u^- are of class C^1 in $\overline{\Omega^+}$ and $\overline{\Omega^-}$, respectively.

Besides the notion of critical pair, which amounts to the vanishing of the *first* variation of the functional, we also introduce the concept of stability, which is defined in terms of the positivity of the *second* variation. Its explicit expression at a regular critical pair (Γ, u) , which will be computed in Theorem 4.4, motivates the definition of the quadratic form $\partial^2 F((\Gamma, u); U) : H^1(\Gamma \cap \Omega) \rightarrow \mathbb{R}$ given by

$$\begin{aligned} \partial^2 F((\Gamma, u); U)[\varphi] := & -2 \int_{\Omega} |\nabla v_{\varphi}|^2 dx + \int_{\Gamma \cap \Omega} |\nabla_{\Gamma} \varphi|^2 d\mathcal{H}^1 + \int_{\Gamma \cap \Omega} H^2 \varphi^2 d\mathcal{H}^1 \\ & - \int_{\Gamma \cap \partial \Omega} H_{\partial \Omega} \varphi^2 d\mathcal{H}^0 \end{aligned} \quad (3.3)$$

where $v_\varphi \in H_U^1(\Omega \setminus \Gamma)$ solves

$$\int_{\Omega} \nabla v_\varphi \cdot \nabla z \, dx + \int_{\Gamma \cap \Omega} [z^+ \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^+) - z^- \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^-)] \, d\mathcal{H}^1 = 0 \quad (3.4)$$

for every $z \in H_U^1(\Omega \setminus \Gamma)$. Notice that the last integral in (3.3) in fact reduces to the sum $H_{\partial\Omega}(x_1)\varphi^2(x_1) + H_{\partial\Omega}(x_2)\varphi^2(x_2)$, where x_1 and x_2 are the intersections of Γ with $\partial\Omega$. The (nonlocal) dependence on U is realized through the function v_φ .

Remark 3.5. The second integral in equation (3.4) has to be intended in the duality sense between $H^{-\frac{1}{2}}(\Gamma \cap \Omega)$ and $H^{\frac{1}{2}}(\Gamma \cap \Omega)$. Indeed, by directly estimating the Gagliardo $H^{\frac{1}{2}}$ -seminorm one can check that the product $\varphi \nabla_\Gamma u^\pm$ belongs to $H^{\frac{1}{2}}(\Gamma \cap \Omega)$ as long as $\nabla_\Gamma u^\pm \in C^{0,\alpha}(\Gamma)$ for some $\alpha > \frac{1}{2}$. In turn, the latter regularity property is guaranteed by Lemma 9.2, recalling that u satisfies the condition (3.1).

Definition 3.6. We say that a regular critical pair (Γ, u) (see Definition 3.3) is *strictly stable* in an admissible subdomain U if

$$\partial^2 F((\Gamma, u); U)[\varphi] > 0 \quad \text{for every } \varphi \in H^1(\Gamma \cap \Omega) \setminus \{0\}. \quad (3.5)$$

The aim of this paper is to discuss the relation between the notion of strict stability of a regular critical pair and the one of local minimality. It is easily seen that the positive semidefiniteness of the quadratic form $\partial^2 F((\Gamma, u); U)$ is a necessary condition for local minimality in U (see [6, Theorem 3.15]). In the main result of the paper we prove that its *strict positivity* is in fact a *sufficient* condition for a regular critical pair to be a local minimizer in the L^1 -sense:

Theorem 3.7. *Let (Γ, u) be a strictly stable regular critical pair in an admissible subdomain U , according to Definition 3.6. Then u is an isolated local minimizer for F in U , in the sense that there exists $\delta > 0$ such that*

$$F(\Gamma, u) < F(K, v) \quad (3.6)$$

for every $(K, v) \in \mathcal{A}(\Omega)$ such that $v = u$ in $(\Omega \setminus U) \cup \partial_D \Omega$ and $0 < \|u - v\|_{L^1(\Omega)} < \delta$.

Remark 3.8. In order to simplify the proofs and the notations we decided to state and prove the previous result only in the simplified situation where Ω is connected and Γ is a regular curve joining two points of $\partial\Omega$. It is straightforward to check that Theorem 3.7 can be generalized to the case where Γ is a finite, disjoint union of curves of class C^∞ , each one connecting two points of $\partial\Omega$ and meeting $\partial\Omega$ orthogonally.

Remark 3.9. The non-vanishing jump condition (point (iii) of Definition 3.3) is not a technical assumption and cannot be dropped: indeed, it is possible to construct examples (see the Remark after Theorem 3.1 in [9]) satisfying all the assumptions of Theorem 3.7 except for this one, for which the conclusion of the theorem does not hold. In our strategy, this hypothesis is needed in order to deduce, in Proposition 7.2, by applying the calibration constructed in [21], that the unique solution of the penalization problem (7.4) is u itself, if β is sufficiently large.

Remark 3.10. A remark concerning the C^3 -regularity assumption on $\partial\Omega$ is also in order. In fact, a careful inspection of the proofs presented in the remaining part of the paper shows that C^3 -regularity is only needed in a neighborhood of $\Gamma \cap \partial\Omega$, whereas away from that intersection C^1 -regularity of $\partial\Omega$ would suffice.

Remark 3.11. We expect the main result of this paper to hold also in higher dimensions. The main technical obstacle to extending our approach to $N \geq 3$ is the lack of a higher dimensional version of Theorem 2.3. Although we didn't check the details, we believe that with such a regularity theorem at hands, the overall strategy presented here would go through.

We conclude with the following consequence of Theorem 3.7, which states that given any family of equicoercive functionals \mathcal{F}_ε which Γ -converge to the relaxed version of F with respect to the L^1 -topology, we can approximate each strictly stable regular critical pair for F by a sequence of local minimizers of the functionals \mathcal{F}_ε . This follows from the abstract result observed in [17, Theorem 4.1]. There is a vast literature concerning the approximation of the Mumford-Shah functional in the sense of Γ -convergence (see, for instance, [5]).

Theorem 3.12 (link with Γ -convergence). *Let (Γ, u) be a strictly stable regular critical pair in an admissible subdomain U . Let $\mathcal{F}_\varepsilon : L^1(\Omega) \rightarrow \mathbb{R} \cup \{+\infty\}$ be a family of equicoercive and lower semi-continuous functionals which Γ -converge as $\varepsilon \rightarrow 0$ to the relaxed functional (see the beginning of Section 7)*

$$\mathcal{F}(v) := \begin{cases} \int_\Omega |\nabla v|^2 dx + \mathcal{H}^1(S_v) & \text{if } v \in SBV(\Omega), v = u \text{ on } (\Omega \setminus U) \cup \partial_D \Omega, \\ +\infty & \text{otherwise in } L^1(\Omega) \end{cases}$$

with respect to the L^1 -topology. Then there exists $\varepsilon_0 > 0$ and a family $(u_\varepsilon)_{\varepsilon < \varepsilon_0}$ of local minimizers of \mathcal{F}_ε such that $u_\varepsilon \rightarrow u$ in $L^1(\Omega)$ as $\varepsilon \rightarrow 0$.

4. COMPUTATION OF THE SECOND VARIATION

In this section we compute the second variation of the functional F . To start with, we fix some notation: for any one-parameter family of functions $(g_s)_{s \in \mathbb{R}}$ we denote the partial derivative with respect to the variable s of the map $(s, x) \mapsto g_s(x)$, evaluated at (t, x) , by $\dot{g}_t(x)$. We usually omit the subscript when $t = 0$. In the following, we fix a regular pair $(K, v) \in \mathcal{A}_{reg}(\Omega)$ and an admissible subdomain U .

Definition 4.1. A flow $(\Phi_t)_t$ is said to be *admissible* for (K, v) in U if it is generated by a vector field $X \in C^2(\mathbb{R}^2; \mathbb{R}^2)$ such that $\text{supp } X \subset\subset U \setminus \partial_D \Omega$ and $X \cdot \nu_{\partial \Omega} = 0$ on $\partial \Omega$, that is, Φ_t solves the equation $\dot{\Phi}_t = X \circ \Phi_t$, $\Phi_0 = Id$.

Remark 4.2. The condition $X \cdot \nu_{\partial \Omega} = 0$ guarantees that the trajectories of points in $\partial \Omega$ remain on $\partial \Omega$: thus $\Phi_t(\bar{\Omega}) = \bar{\Omega}$ for every t . Observe also that, since $\text{supp } X \subset\subset U \setminus \partial_D \Omega$, we have that $K_{\Phi_t} \subset U \setminus \partial_D \Omega$ for every t , where we set $K_{\Phi_t} := \Phi_t(K)$.

Given an orientation preserving diffeomorphism $\Phi \in C^\infty(\bar{\Omega}; \bar{\Omega})$ such that $\text{supp}(\Phi - Id) \subset\subset U \setminus \partial_D \Omega$, we define v_Φ as the unique solution in $H^1(\Omega \setminus K_\Phi)$ (up to additive constants in the connected components of $\Omega \setminus K_\Phi$ whose boundary does not contain $\partial_D \Omega$) to

$$\begin{cases} \int_{\Omega \setminus K_\Phi} \nabla v_\Phi \cdot \nabla z = 0 & \text{for every } z \in H_U^1(\Omega \setminus K_\Phi), \\ v_\Phi = v & \text{in } (\Omega \setminus U) \cup \partial_D \Omega. \end{cases} \quad (4.1)$$

Definition 4.3. Let $(\Phi_t)_t$ be an admissible flow for (K, v) in U . We define the *first and second variations of F at (K, v) in U along $(\Phi_t)_t$* to be

$$\frac{d}{dt} F((K_{\Phi_t}, v_{\Phi_t}); U)|_{t=0}, \quad \frac{d^2}{dt^2} F((K_{\Phi_t}, v_{\Phi_t}); U)|_{t=0}$$

respectively, where v_{Φ_t} is defined as in (4.1) with Φ replaced by Φ_t .

Notice that this definition makes sense since the existence of the derivatives is guaranteed by the regularity result proved in [6, Proposition 8.1], which can be adapted to the present setting. In particular, this result implies that the map $(t, x) \mapsto v_{\Phi_t}(x)$ is differentiable with respect to the variable t and that $\dot{v}_{\Phi_t} \in H_U^1(\Omega \setminus K_{\Phi_t})$. We set $\dot{v} := \dot{v}_{\Phi_0}$.

In the following theorem we compute explicitly the second variation of the functional F . We stress that, comparing with the analogous result obtained in [6, Theorem 3.6], we allow here the admissible variations to affect also the intersection of the discontinuity set K with the boundary of Ω , while in the quoted paper only variations compactly supported in Ω were considered. As a consequence, in the present situation boundary terms arise when integration by parts are performed: in particular this happens for the derivatives of the surface term, while the first and second variations of the volume term remain unchanged. We refer also to [25], where a similar computation for the second variation of the surface area was carried out taking into account boundary effects, in the case of a critical set (the novelty here is that we will be able to get an expression of the second variation at a generic regular pair, not necessarily critical).

Theorem 4.4. *Let $(K, \nu) \in \mathcal{A}_{reg}(\Omega)$ be a regular pair for F , let U be an admissible subdomain, and let $(\Phi_t)_t$ be an admissible flow in U associated to a vector field X . Then the function \dot{v} belongs to $H^1_U(\Omega \setminus K)$ and satisfies the equation*

$$\int_{\Omega} \nabla \dot{v} \cdot \nabla z \, dx + \int_{K \cap \Omega} \left[\operatorname{div}_K((X \cdot \nu) \nabla_K v^+) z^+ - \operatorname{div}_K((X \cdot \nu) \nabla_K v^-) z^- \right] d\mathcal{H}^1 = 0 \quad (4.2)$$

for every $z \in H^1_U(\Omega \setminus K)$. Moreover, the second variation of F at (K, ν) in U along $(\Phi_t)_t$ is given by

$$\begin{aligned} \frac{d^2}{dt^2} F((K_{\Phi_t}, \nu_{\Phi_t}); U)|_{t=0} &= 2 \int_{K \cap \Omega} (\dot{v}^+ \partial_\nu \dot{v}^+ - \dot{v}^- \partial_\nu \dot{v}^-) d\mathcal{H}^1 + \int_{K \cap \Omega} |\nabla_K(X \cdot \nu)|^2 d\mathcal{H}^1 \\ &+ \int_{K \cap \Omega} H^2(X \cdot \nu)^2 d\mathcal{H}^1 + \int_{K \cap \Omega} f(Z \cdot \nu - 2X^\parallel \cdot \nabla_K(X \cdot \nu) + D\nu[X^\parallel, X^\parallel] - H(X \cdot \nu)^2) d\mathcal{H}^1 \\ &+ \int_{K \cap \partial\Omega} Z \cdot \eta d\mathcal{H}^0, \end{aligned} \quad (4.3)$$

where $f := |\nabla_K v^-|^2 - |\nabla_K v^+|^2 + H$, $Z := DX[X]$, and we split the field X in its tangential and normal components to K :

$$X = X^\parallel + (X \cdot \nu)\nu \quad \text{on } K. \quad (4.4)$$

Remark 4.5. As in (3.4), the second integral in equation (4.2) has to be intended in the duality sense between $H^{-\frac{1}{2}}(K \cap \Omega)$ and $H^{\frac{1}{2}}(K \cap \Omega)$ (see Remark 3.5). Integrations by parts yields

$$- \int_{\Omega} |\nabla \dot{v}|^2 \, dx = \int_{K \cap \Omega} [\dot{v}^+ \partial_\nu \dot{v}^+ - \dot{v}^- \partial_\nu \dot{v}^-] d\mathcal{H}^1.$$

Before proving Theorem 4.4, we collect in the following lemma some auxiliary identities which will be used in the computation of the second variation.

Lemma 4.6. *The following identities hold:*

- (a) $\dot{\nu} = -(D_K X)^T[\nu] - D_K \nu[X] = -\nabla_K(X \cdot \nu)$ on K ;
- (b) $\frac{\partial}{\partial t}(\eta_{\Phi_t} \circ \Phi_t)|_{t=0} = (D_K X)^T[\nu, \eta]\nu$ on $K \cap \partial\Omega$;
- (c) $(X \cdot \nu)\dot{\nu} \cdot \eta + X \cdot \frac{\partial}{\partial t}(\eta_{\Phi_t} \circ \Phi_t)|_{t=0} = -H(X \cdot \nu)(X \cdot \eta)$ on $K \cap \partial\Omega$;
- (d) $DX[X, \nu_{\partial\Omega}] + D\nu_{\partial\Omega}[X, X] = 0$ on $K \cap \partial\Omega$.

Proof. Equality (a) is proved in [6, Lemma 3.8, (f)]. To prove (b), we set $v_t := D\Phi_t[\eta]$ and recalling (2.3) we have

$$\begin{aligned} \frac{\partial}{\partial t}(\eta_{\Phi_t} \circ \Phi_t)|_{t=0} &= \frac{\partial}{\partial t} \left(\frac{v_t}{|v_t|} \right) \Big|_{t=0} = \dot{v} - (\dot{v} \cdot \eta)\eta \\ &= DX[\eta] - DX[\eta, \eta]\eta = DX[\eta, \nu]\nu, \end{aligned}$$

which is (b). We obtain (c) by combining (a) and (b):

$$(X \cdot \nu)\dot{\nu} \cdot \eta + X \cdot \frac{\partial}{\partial t}(\eta_{\Phi_t} \circ \Phi_t)|_{t=0} = -(X \cdot \nu)D_K \nu[X, \eta] = -H(X \cdot \nu)(X \cdot \eta),$$

where the last equality follows by writing $X = (X \cdot \nu)\nu + (X \cdot \eta)\eta$ and observing that $D_K \nu[\nu] = 0$. Equation (d) follows by differentiating with respect to t at $t = 0$ the identity

$$(X \circ \Phi_t) \cdot (\nu_{\partial\Omega} \circ \Phi_t) = 0,$$

which holds on $K \cap \partial\Omega$. □

Proof of Theorem 4.4. We split the proof of the theorem into three steps.

Step 1. Derivation of the equation solved by \dot{v} . As already observed, the result contained in [6, Proposition 8.1] guarantees that $\dot{v} \in H^1_U(\Omega \setminus K)$. Given any test function $z \in H^1_U(\Omega \setminus K)$ with

$\text{supp } z \cap K = \emptyset$, for t small enough we have $\text{supp } z \subset \Omega \setminus K_{\Phi_t}$, and in particular $z \in H^1_U(\Omega \setminus K_{\Phi_t})$. Hence by (4.1) we deduce

$$\int_{\Omega} \nabla v_{\Phi_t} \cdot \nabla z \, dx = 0,$$

so that differentiating with respect to t at $t = 0$ we obtain that \dot{v} is harmonic in $(\Omega \cap U) \setminus K$ and $\nabla \dot{v} \cdot \nu_{\partial\Omega} = 0$ on $(\partial\Omega \cap U) \setminus \partial_D\Omega$. In addition, it is shown in Step 1 of the proof of [6, Theorem 3.6] that

$$\partial_\nu v^\pm = \text{div}_K((X \cdot \nu)\nabla_K v^\pm) \quad \text{on } K \cap \Omega.$$

By this expression we have that $\partial_\nu v^\pm \in H^{-\frac{1}{2}}(K \cap \Omega)$ (see Remark 3.5), and hence the previous conditions are equivalent to (4.2) by integration by parts.

Step 2. Computation of the first variation. The same computation carried out in Step 2 of the proof of [6, Theorem 3.6] leads to

$$\frac{d}{dt} \int_{\Omega} |\nabla v_{\Phi_t}|^2 \, dx = \int_{\Omega} \text{div}(|\nabla v_{\Phi_t}|^2 X) \, dy.$$

Hence, applying the divergence theorem we obtain

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} |\nabla v_{\Phi_t}|^2 \, dx &= \int_{\partial\Omega} |\nabla v_{\Phi_t}|^2 (X \cdot \nu_{\partial\Omega}) \, d\mathcal{H}^1 + \int_{K_{\Phi_t} \cap \Omega} (|\nabla v_{\Phi_t}^-|^2 - |\nabla v_{\Phi_t}^+|^2) (X \cdot \nu_{\Phi_t}) \, d\mathcal{H}^1 \\ &= \int_{K_{\Phi_t} \cap \Omega} (|\nabla_{K_{\Phi_t}} v_{\Phi_t}^-|^2 - |\nabla_{K_{\Phi_t}} v_{\Phi_t}^+|^2) (X \cdot \nu_{\Phi_t}) \, d\mathcal{H}^1 \end{aligned}$$

where to deduce the last equality we used $X \cdot \nu_{\partial\Omega} = 0$ and the fact that $\partial_{\nu_{\Phi_t}} v_{\Phi_t}^\pm$ vanishes on K_{Φ_t} . Concerning the surface term, we start from the well known formula for the first variation of the area functional (see, for instance, [24, Chapter 2, Section 9]) and we use the divergence theorem on $K_{\Phi_t} \cap \Omega$, to obtain

$$\begin{aligned} \frac{d}{dt} \mathcal{H}^1(K_{\Phi_t} \cap \Omega) &= \int_{K_{\Phi_t} \cap \Omega} \text{div}_{K_{\Phi_t}} X \, d\mathcal{H}^1 \\ &= \int_{K_{\Phi_t} \cap \Omega} H_{\Phi_t} (X \cdot \nu_{\Phi_t}) \, d\mathcal{H}^1 + \int_{K_{\Phi_t} \cap \partial\Omega} X \cdot \eta_{\Phi_t} \, d\mathcal{H}^0, \end{aligned}$$

where we recall that H_{Φ_t} stands for the curvature of K_{Φ_t} . Thus we can conclude that

$$\frac{d}{dt} F((K_{\Phi_t}, v_{\Phi_t}); U) = \int_{K_{\Phi_t} \cap \Omega} f_t (X \cdot \nu_{\Phi_t}) \, d\mathcal{H}^1 + \int_{K_{\Phi_t} \cap \partial\Omega} X \cdot \eta_{\Phi_t} \, d\mathcal{H}^0, \quad (4.5)$$

where $f_t := |\nabla_{K_{\Phi_t}} v_{\Phi_t}^-|^2 - |\nabla_{K_{\Phi_t}} v_{\Phi_t}^+|^2 + H_{\Phi_t}$. In particular, evaluating (4.5) at $t = 0$ we obtain

$$\frac{d}{dt} F((K_{\Phi_t}, v_{\Phi_t}); U)|_{t=0} = \int_{K \cap \Omega} f(X \cdot \nu) \, d\mathcal{H}^1 + \int_{K \cap \partial\Omega} X \cdot \eta \, d\mathcal{H}^0. \quad (4.6)$$

Step 3. Computation of the second variation. We have to differentiate again (4.5) at $t = 0$. By a change of variables we have

$$\begin{aligned} \frac{d^2}{dt^2} F((K_{\Phi_t}, v_{\Phi_t}); U)|_{t=0} &= \int_{K \cap \Omega} \frac{\partial}{\partial t} (f_t \circ \Phi_t)|_{t=0} (X \cdot \nu) \, d\mathcal{H}^1 \\ &+ \int_{K \cap \Omega} f \frac{\partial}{\partial t} (\dot{\Phi}_t \cdot (\nu_{\Phi_t} \circ \Phi_t) J_{\Phi_t})|_{t=0} \, d\mathcal{H}^1 + \frac{d}{dt} \left(\int_{K_{\Phi_t} \cap \partial\Omega} X \cdot \eta_{\Phi_t} \, d\mathcal{H}^0 \right) \Big|_{t=0} \\ &=: I_1 + I_2 + I_3. \end{aligned} \quad (4.7)$$

The first integral I_1 is equal to

$$I_1 = \int_{K \cap \Omega} \dot{f}(X \cdot \nu) \, d\mathcal{H}^1 + \int_{K \cap \Omega} (\nabla f \cdot \nu)(X \cdot \nu)^2 \, d\mathcal{H}^1 + \int_{K \cap \Omega} (\nabla_K f \cdot X^\parallel)(X \cdot \nu) \, d\mathcal{H}^1, \quad (4.8)$$

while using [6, Lemma 3.8, (g)] we have

$$I_2 = \int_{K \cap \Omega} f \text{div}_K((X \cdot \nu)X) \, d\mathcal{H}^1 + \int_{K \cap \Omega} f(Z \cdot \nu - 2X^\parallel \cdot \nabla_K(X \cdot \nu) + D\nu[X^\parallel, X^\parallel]) \, d\mathcal{H}^1. \quad (4.9)$$

Applying the divergence formula on $K \cap \Omega$ we obtain

$$\begin{aligned} & \int_{K \cap \Omega} (\nabla_K f \cdot X^\parallel)(X \cdot \nu) d\mathcal{H}^1 + \int_{K \cap \Omega} f \operatorname{div}_K((X \cdot \nu)X) d\mathcal{H}^1 \\ &= \int_{K \cap \Omega} f H(X \cdot \nu)^2 d\mathcal{H}^1 + \int_{K \cap \partial\Omega} f(X \cdot \nu)(X \cdot \eta) d\mathcal{H}^0, \end{aligned} \quad (4.10)$$

while using [6, formula (3.17)] we get

$$\int_{K \cap \Omega} (\nabla f \cdot \nu)(X \cdot \nu)^2 d\mathcal{H}^1 = \int_{K \cap \Omega} (H^2 - 2fH)(X \cdot \nu)^2 d\mathcal{H}^1. \quad (4.11)$$

Differentiating f_t with respect to t we obtain

$$\int_{K \cap \Omega} \dot{f}(X \cdot \nu) d\mathcal{H}^1 = \int_{K \cap \Omega} (2\nabla_K v^- \cdot \nabla_K \dot{v}^- - 2\nabla_K v^+ \cdot \nabla_K \dot{v}^+ + \dot{H})(X \cdot \nu) d\mathcal{H}^1, \quad (4.12)$$

and an integration by parts yields

$$\begin{aligned} & 2 \int_{K \cap \Omega} (\nabla_K v^\pm \cdot \nabla_K \dot{v}^\pm)(X \cdot \nu) d\mathcal{H}^1 \\ &= -2 \int_{K \cap \Omega} \dot{v}^\pm \operatorname{div}_K((X \cdot \nu)\nabla_K v^\pm) d\mathcal{H}^1 + 2 \int_{K \cap \partial\Omega} \dot{v}^\pm(X \cdot \nu)(\nabla_K v^\pm \cdot \eta) d\mathcal{H}^0 \\ &= -2 \int_{K \cap \Omega} \dot{v}^\pm \partial_\nu \dot{v}^\pm d\mathcal{H}^1, \end{aligned} \quad (4.13)$$

where the last equality follows by (4.2) and by observing that ∇v^\pm vanishes on $K \cap \partial\Omega$, as v satisfies homogeneous Neumann boundary conditions on K and on $\partial\Omega$ (∇v is regular up to $K \cap \partial\Omega$ by Lemma 9.2). Since $\partial_\nu \dot{v} \cdot \nu = -\dot{v} \cdot \partial_\nu \nu = 0$, we have $\operatorname{div} \dot{v} = \operatorname{div}_K \dot{v}$ and in turn $\dot{H} = \operatorname{div}_K \dot{v}$. Hence, integrating by parts and using (a) of Lemma 4.6, we deduce

$$\begin{aligned} & \int_{K \cap \Omega} \dot{H}(X \cdot \nu) d\mathcal{H}^1 = \int_{K \cap \Omega} \operatorname{div}_K \dot{v}(X \cdot \nu) d\mathcal{H}^1 \\ &= - \int_{K \cap \Omega} \dot{v} \cdot \nabla_K(X \cdot \nu) d\mathcal{H}^1 + \int_{K \cap \partial\Omega} (X \cdot \nu)(\dot{v} \cdot \eta) d\mathcal{H}^0 \\ &= \int_{K \cap \Omega} |\nabla_K(X \cdot \nu)|^2 d\mathcal{H}^1 + \int_{K \cap \partial\Omega} (X \cdot \nu)(\dot{v} \cdot \eta) d\mathcal{H}^0. \end{aligned} \quad (4.14)$$

We finally compute I_3 :

$$\begin{aligned} I_3 &= \frac{d}{dt} \left(\int_{K_{\Phi_t} \cap \partial\Omega} X \cdot \eta_{\Phi_t} d\mathcal{H}^0 \right) \Big|_{t=0} = \sum_{x \in K \cap \partial\Omega} \frac{\partial}{\partial t} \left(X(\Phi_t(x)) \cdot \eta_{\Phi_t}(\Phi_t(x)) \right) \Big|_{t=0} \\ &= \sum_{x \in K \cap \partial\Omega} Z(x) \cdot \eta(x) + \sum_{x \in K \cap \partial\Omega} X(x) \cdot \frac{\partial}{\partial t} (\eta_{\Phi_t} \circ \Phi_t(x)) \Big|_{t=0}. \end{aligned} \quad (4.15)$$

We finally obtain the expression (4.3) by collecting (4.7)–(4.15), using equality (c) of Lemma 4.6, and observing that the quantity $f - H = |\nabla_K v^-|^2 - |\nabla_K v^+|^2$ vanishes on $K \cap \partial\Omega$ (since, as observed before, $\nabla v^\pm = 0$ on $K \cap \partial\Omega$). \square

Remark 4.7. We observe that we can easily obtain an expression for the second variation of the functional F at a generic t . Indeed, by exploiting the property $\Phi_{t+s} = \Phi_t \circ \Phi_s$ of the flow, we have

$$\frac{d^2}{dh^2} F((K_{\Phi_h}, v_{\Phi_h}); U) \Big|_{h=t} = \frac{d^2}{ds^2} F((\Phi_{t+s}(K), v_{\Phi_{t+s}}); U) \Big|_{s=0} = \frac{d^2}{ds^2} F(\Phi_s(K_{\Phi_t}), (v_{\Phi_t})_{\Phi_s}) \Big|_{s=0},$$

and we can directly apply Theorem 4.4 to the regular pair (K_{Φ_t}, v_{Φ_t}) .

Remark 4.8. The formula (4.6) for the first variation of F motivates the definition of critical pair (see Definition 3.3). Indeed, assuming that (4.6) vanishes for each vector field X which is tangent to $\partial\Omega$, we first obtain that $f = 0$ on $K \cap \Omega$ by considering arbitrary vector fields with $\operatorname{supp} X \subset\subset \Omega$. Then, using this information and dropping the requirement on the support of X , we deduce the orthogonality of K and $\partial\Omega$.

Corollary 4.9. *Assume that (Γ, u) is a regular critical pair. Then*

$$\begin{aligned} \frac{d^2}{dt^2} F((\Gamma_{\Phi_t}, u_{\Phi_t}); U)|_{t=0} &= -2 \int_{\Omega} |\nabla \dot{u}|^2 dx + \int_{\Gamma \cap \Omega} |\nabla_{\Gamma}(X \cdot \nu)|^2 d\mathcal{H}^1 \\ &\quad + \int_{\Gamma \cap \Omega} H^2(X \cdot \nu)^2 d\mathcal{H}^1 - \int_{\Gamma \cap \partial\Omega} H_{\partial\Omega}(X \cdot \nu)^2 d\mathcal{H}^0, \end{aligned} \quad (4.16)$$

where $H_{\partial\Omega} := \operatorname{div} \nu_{\partial\Omega}$ denotes the curvature of $\partial\Omega$.

Proof. The first integral in (4.3) can be rewritten as $-2 \int_{\Omega} |\nabla \dot{u}|^2 dx$ thanks to (4.2) (see Remark 4.5). To obtain the expression in (4.16) it is now sufficient to observe that at a critical pair we have $f = 0$ on $K \cap \Omega$, $X \cdot \eta = X \cdot \nu_{\partial\Omega} = 0$ on $K \cap \partial\Omega$, and

$$Z \cdot \eta = DX[X, \nu_{\partial\Omega}] = -D\nu_{\partial\Omega}[X, X] = -(X \cdot \nu)^2 D\nu_{\partial\Omega}[\nu, \nu] = -H_{\partial\Omega}(X \cdot \nu)^2$$

on $K \cap \partial\Omega$ by (d) of Lemma 4.6. \square

4.1. The second order condition. In the following we assume that (Γ, u) is a regular critical pair and U is an admissible subdomain. Notice that the expression of the second variation of F at (Γ, u) proved in Corollary 4.9 motivates the definition of the quadratic form (3.3) and the notion of strict stability that we introduced in Definition 3.6.

Following the approach of [6], we start paving the way for the main result by proving two equivalent formulations of condition (3.5), one in terms of the first eigenvalue of a suitable compact linear operator defined on $H^1(\Gamma \cap \Omega)$ and the other in terms of a dual minimum problem. Let us start by introducing the following bilinear form on $H^1(\Gamma \cap \Omega)$:

$$(\varphi, \psi)_{\sim} := \int_{\Gamma \cap \Omega} \nabla_{\Gamma} \varphi \cdot \nabla_{\Gamma} \psi d\mathcal{H}^1 + \int_{\Gamma \cap \Omega} H^2 \varphi \psi d\mathcal{H}^1 - \int_{\Gamma \cap \partial\Omega} H_{\partial\Omega} \varphi \psi d\mathcal{H}^0 \quad (4.17)$$

for every $\varphi, \psi \in H^1(\Gamma \cap \Omega)$. The proof of the following proposition can be obtained by simply adapting [6, Proposition 4.2] to our slightly different situation.

Proposition 4.10. *Assume that*

$$(\varphi, \varphi)_{\sim} > 0 \quad \text{for every } \varphi \in H^1(\Gamma \cap \Omega) \setminus \{0\}. \quad (4.18)$$

Then $(\cdot, \cdot)_{\sim}$ is a scalar product which defines an equivalent norm on $H^1(\Gamma \cap \Omega)$, denoted by $\|\cdot\|_{\sim}$.

The announced equivalent formulations of condition (3.5) are stated in the following proposition. Also for this proof we refer the reader to Proposition 4.3, Theorem 4.6 and Theorem 4.10 in [6], which can be directly adapted with the natural modifications, taking into account Remark 3.5.

Proposition 4.11. *The following statements are equivalent.*

- (i) *Condition (3.5) is satisfied.*
- (ii) *Condition (4.18) holds, and the monotone, compact, self-adjoint operator $T : H^1(\Gamma \cap \Omega) \rightarrow H^1(\Gamma \cap \Omega)$ defined by duality as*

$$(T\varphi, \psi)_{\sim} = -2 \int_{\Gamma \cap \Omega} [v_{\varphi}^+ \operatorname{div}_{\Gamma}(\psi \nabla_{\Gamma} u^+) - v_{\varphi}^- \operatorname{div}_{\Gamma}(\psi \nabla_{\Gamma} u^-)] d\mathcal{H}^1 \quad (4.19)$$

for every $\varphi, \psi \in H^1(\Gamma \cap \Omega)$ (where v_{φ} is defined in (3.4)), satisfies

$$\lambda_1(U) := \max_{\|\varphi\|_{\sim}=1} (T\varphi, \varphi)_{\sim} < 1 \quad (4.20)$$

(where the dependence on U is realized through the function v_{φ}).

- (iii) *Condition (4.18) holds, and defining, for $v \in H_U^1(\Omega \setminus \Gamma)$, Φ_v as the unique solution in $H^1(\Gamma \cap \Omega)$ to*

$$(\Phi_v, \psi)_{\sim} = -2 \int_{\Gamma \cap \Omega} [v^+ \operatorname{div}_{\Gamma}(\psi \nabla_{\Gamma} u^+) - v^- \operatorname{div}_{\Gamma}(\psi \nabla_{\Gamma} u^-)] d\mathcal{H}^1$$

for every $\psi \in H^1(\Gamma \cap \Omega)$, one has

$$\mu(U) := \min \left\{ 2 \int_{\Omega} |\nabla v|^2 dx : v \in H_U^1(\Omega \setminus \Gamma), \|\Phi_v\|_{\sim} = 1 \right\} > 1. \quad (4.21)$$

We will omit the dependence on U for λ_1 and μ where there is no risk of ambiguity.

Remark 4.12. Notice that if condition (4.18) is satisfied, then by Proposition 4.10 and by the Riesz Theorem the operator T is well defined. By (3.4) we immediately have

$$(T\varphi, \psi)_{\sim} = 2 \int_{\Omega} \nabla v_{\varphi} \cdot \nabla v_{\psi} \, dx.$$

Moreover comparing with (3.3) we see that

$$\partial^2 F((\Gamma, u); U)[\varphi] = -(T\varphi, \varphi)_{\sim} + \|\varphi\|_{\sim}^2.$$

Corollary 4.13. *Assume (3.5). Then there exists a constant $C > 0$ such that*

$$\partial^2 F((\Gamma, u); U)[\varphi] \geq C \|\varphi\|_{H^1(\Gamma \cap \Omega)}^2 \quad \text{for every } \varphi \in H^1(\Gamma \cap \Omega).$$

Proof. By Remark 4.12 and (4.20)

$$\partial^2 F((\Gamma, u); U)[\varphi] = \|\varphi\|_{\sim}^2 - (T\varphi, \varphi)_{\sim} \geq (1 - \lambda_1) \|\varphi\|_{\sim}^2,$$

hence the conclusion follows by Proposition 4.11 and Proposition 4.10. \square

From the definition in (4.21) it is clear that μ depends monotonically on the domain U . This is made explicit by the following corollary.

Corollary 4.14. *Let U_1, U_2 be admissible subdomains for (Γ, u) , with $U_1 \subset U_2$. Then $\mu(U_1) \geq \mu(U_2)$. In particular, if condition (3.5) is satisfied in U_2 , then it also holds in U_1 .*

Corollary 4.15. *Assume that condition (3.5) holds in U . Let U_n be a decreasing sequence of admissible subdomains for (Γ, u) such that U is the interior part of $\bigcap_n U_n$. Then (3.5) holds in U_n , if n is sufficiently large.*

Proof. In view of (4.21) it is sufficient to show that $\lim_n \mu(U_n) \geq \mu(U)$. Let $v_n \in H_{U_n}^1(\Omega \setminus \Gamma)$ be a solution to (4.21) with U replaced by U_n . Then $v_n \in H_{U_1}^1(\Omega \setminus \Gamma)$ and $2 \int_{\Omega} |\nabla v_n|^2 \, dx = \mu(U_n) \leq \mu(U)$, where the inequality follows from Corollary 4.14. Hence, up to subsequences, $v_n \rightharpoonup v \in H_{U_1}^1(\Omega \setminus \Gamma)$. Moreover, $v = 0$ a.e. in $U_1 \setminus U$, so that $v \in H_U^1(\Omega \setminus \Gamma)$ and v is admissible in problem (4.21) (by the compactness of the map $v \mapsto \Phi_v$): we conclude that

$$\lim_{n \rightarrow \infty} \mu(U_n) = \lim_{n \rightarrow \infty} 2 \int_{\Omega} |\nabla v_n|^2 \, dx \geq 2 \int_{\Omega} |\nabla v|^2 \, dx \geq \mu(U),$$

as claimed. \square

Remark 4.16. We observe that, in proving Theorem 3.7, we can assume without loss of generality that U is an open set of class C^∞ and that ∂U and $\partial \Omega$ are orthogonal where they intersect. Indeed, assume to have proved the theorem under these additional assumptions. If U is any admissible subdomain for (Γ, u) , we can find a decreasing sequence of admissible subdomains U_n of class C^∞ , with boundaries meeting $\partial \Omega$ orthogonally, such that U is the interior part of $\bigcap_n U_n$. It follows from Corollary 4.15 that the second variation is strictly positive in U_n for n large enough, and hence (Γ, u) is an isolated local minimizer in U_n . This immediately yields the conclusion also in the initial domain U .

5. LOCAL $W^{2,\infty}$ -MINIMALITY

In this section, as a first step toward the proof of Theorem 3.7, we show how the strategy developed in [6] can be adapted to the present setting in order to prove that the positiveness condition (3.5) is sufficient for a regular critical pair to be a local minimizer with respect to variations of class $W^{2,\infty}$ of the discontinuity set. For the rest of the section (Γ, u) will be a fixed strictly stable regular critical pair in an admissible subdomain U . For $\eta > 0$, we denote by

$$\mathcal{N}_\eta(\Gamma) := \{x \in \mathbb{R}^2 : \text{dist}(x, \Gamma) < \eta\}$$

the η -tubular neighborhood of Γ .

In order to give a proper notion of sets which are close to Γ in the $W^{2,\infty}$ -sense, we now introduce a suitable flow in U whose trajectories intersect Γ orthogonally. To this aim, we start

by fixing $\eta_0 > 0$ such that $\mathcal{N}_{\eta_0}(\Gamma) \subset\subset U \setminus \partial_D \Omega$, and a vector field $X \in C^2(\mathbb{R}^2; \mathbb{R}^2)$ such that $\text{supp } X \subset\subset U \setminus \partial_D \Omega$, $X = \nu$ on Γ , $X \cdot \nu_{\partial \Omega} = 0$ on $\partial \Omega$, and $|X| = 1$ in $\mathcal{N}_{\eta_0}(\Gamma)$. We denote by $\Psi : \mathbb{R} \times \bar{\Omega} \rightarrow \bar{\Omega}$ the flow generated by X :

$$\frac{\partial}{\partial t} \Psi(t, x) = X(\Psi(t, x)), \quad \Psi(0, x) = x.$$

Observe that (by taking a smaller η_0 if necessary) for every $y \in \mathcal{N}_{\eta_0}(\Gamma)$ are uniquely determined two points $\pi(y) \in \Gamma$ and $\tau(y) \in \mathbb{R}$ such that $y = \Psi(\tau(y), \pi(y))$. The existence of the maps π and τ , as well as the fact that they are of class C^2 , is guaranteed by the Implicit Function Theorem.

We define, for $\delta > 0$, the following class of functions:

$$\mathcal{D}_\delta := \{\psi \in C^2(\Gamma) : \|\psi\|_{C^2(\Gamma)} < \delta\}.$$

We can extend each function $\psi \in \mathcal{D}_\delta$ to $\mathcal{N}_{\eta_0}(\Gamma)$ by setting $\psi(y) := \psi(\pi(y))$, in such a way that ψ is constant along the trajectories of the flow Ψ . We associate with ψ the diffeomorphism $\Phi^\psi(x) := \Psi(\psi(x), x)$, and we remark that

$$\|\Phi^\psi - Id\|_{C^2(\Gamma)} \leq C \|\psi\|_{C^2(\Gamma)} \quad (5.1)$$

for some constant C independent of $\psi \in \mathcal{D}_\delta$. Finally, we define the set

$$\Gamma_\psi := \Phi^\psi(\Gamma) = \{\Psi(\psi(x), x) : x \in \Gamma\}, \quad (5.2)$$

and the function $u_\psi := u_{\Phi^\psi}$ as the unique solution in $H^1(\Omega \setminus \Gamma_\psi)$ to

$$\int_{\Omega \setminus \Gamma_\psi} \nabla u_\psi \cdot \nabla z = 0 \quad \text{for every } z \in H_U^1(\Omega \setminus \Gamma_\psi)$$

with $u_\psi = u$ in $(\Omega \setminus U) \cup \partial_D \Omega$. We will also denote by $\nu_\psi := \nu_{\Phi^\psi}$ and $\eta_\psi := \eta_{\Phi^\psi}$ the vectors defined on Γ_ψ and $\Gamma_\psi \cap \partial \Omega$ by (2.2) and (2.3) respectively, and by $H_\psi := \text{div}_{\Gamma_\psi} \nu_\psi$ the curvature of Γ_ψ .

Remark 5.1. For $\psi \in \mathcal{D}_\delta$, the function u_ψ is a weak solution to the Neumann problem

$$\begin{cases} \Delta u_\psi = 0 & \text{in } (\Omega \cap U) \setminus \Gamma_\psi, \\ \partial_{\nu_\psi} u_\psi = 0 & \text{on } \Gamma_\psi \cap \Omega, \\ \partial_{\nu_{\partial \Omega}} u_\psi = 0 & \text{on } (\partial \Omega \cap \bar{U}) \setminus \partial_D \Omega, \end{cases}$$

and the sets Γ_ψ are uniformly bounded in C^2 , by (5.1). Hence, by classical results and by using Lemma 9.2 to deal with the regularity in a neighborhood of the boundary $\Gamma_\psi \cap \partial \Omega$, we obtain that the functions u_ψ^\pm are of class $C^{1,\gamma}$ up to $\Gamma_\psi \cap \bar{\Omega}$, for some $\gamma \in (\frac{1}{2}, 1)$, with $C^{1,\gamma}$ -norm uniformly bounded with respect to $\psi \in \mathcal{D}_\delta$. More precisely,

$$\sup_{\psi \in \mathcal{D}_\delta} \|\nabla_\Gamma(u_\psi^\pm \circ \Phi^\psi)\|_{C^{0,\gamma}(\Gamma \cap \bar{\Omega}; \mathbb{R}^2)} < +\infty,$$

and, as an application of Ascoli–Arzelà Theorem, we also have

$$\sup_{\psi \in \mathcal{D}_\delta} \|\nabla_\Gamma(u_\psi^\pm \circ \Phi^\psi) - \nabla_\Gamma u^\pm\|_{C^{0,\alpha}(\Gamma \cap \bar{\Omega}; \mathbb{R}^2)} \rightarrow 0$$

for every $\alpha \in (0, \gamma)$, as $\delta \rightarrow 0$.

The main result of this section is the following.

Theorem 5.2. *Let (Γ, u) be a strictly stable regular critical pair in an admissible subdomain U , according to Definition 3.6. Then (Γ, u) is an isolated local $W^{2,\infty}$ -minimizer for F in U , in the sense that there exist $\delta > 0$ and $C > 0$ such that*

$$F(\Gamma_\psi, v) \geq F(\Gamma, u) + C \|\psi\|_{H^1(\Gamma \cap \Omega)}^2$$

for every $\psi \in W^{2,\infty}(\Gamma \cap \Omega)$ such that $\|\psi\|_{W^{2,\infty}(\Gamma \cap \Omega)} < \delta$, and for every $v \in H^1(\Omega \setminus \Gamma_\psi)$ with $v = u$ in $(\Omega \setminus U) \cup \partial_D \Omega$ (where the set Γ_ψ is defined in (5.2)).

The remaining part of this section is entirely devoted to the proof of Theorem 5.2. We start by fixing $\delta_0 > 0$ such that $\Gamma_\psi \subset \mathcal{N}_{\eta_0}(\Gamma)$ for every $\psi \in \mathcal{D}_{\delta_0}$, where $\mathcal{N}_{\eta_0}(\Gamma)$ is the tubular neighborhood of Γ fixed at the beginning of this section. Our first task is to associate, with every $\psi \in \mathcal{D}_{\delta_0}$, an admissible flow $(\Phi_t)_t$ connecting Γ to Γ_ψ : this can be easily done by setting

$$\Phi_t(x) := \Psi(t\psi(x), x). \quad (5.3)$$

The flow Φ_t is admissible in U (according to Definition 4.1), as it is generated by the vector field

$$X_\psi := \psi X, \quad (5.4)$$

where X is defined at the beginning of this section. Moreover it satisfies $\Phi_1(\Gamma) = \Gamma_\psi$, and

$$\|\Phi_t - Id\|_{C^2(\Gamma)} \leq C \|\psi\|_{C^2(\Gamma)} \quad (5.5)$$

for every $t \in [0, 1]$, where C is a positive constant independent of $\psi \in \mathcal{D}_{\delta_0}$. We also introduce the vector field

$$Z_\psi := DX_\psi[X_\psi] = \psi^2 DX[X] \quad (5.6)$$

(the last equality follows by a direct computation, by observing that $\nabla\psi \cdot X = 0$ since ψ is constant along the trajectories of the flow generated by X). Notice that by (5.4) and (5.6) we immediately have the estimates

$$|X_\psi| \leq |\psi|, \quad |Z_\psi| \leq C |\psi|^2 \quad \text{in } \mathcal{N}_{\eta_0}(\Gamma), \quad (5.7)$$

where C is a positive constant independent of ψ . In the following lemma we collect some technical estimates concerning the above construction that will be used in the proof of the main result of this section.

Lemma 5.3. *Given $\varepsilon > 0$, there exists $\delta(\varepsilon) > 0$ such that for every $\psi \in \mathcal{D}_{\delta(\varepsilon)}$ the following estimates hold:*

- (a) $\frac{1}{2} \|\psi\|_{H^1(\Gamma \cap \Omega)}^2 \leq \|X_\psi \cdot \nu_\psi\|_{H^1(\Gamma_\psi \cap \Omega)}^2 \leq 2 \|\psi\|_{H^1(\Gamma \cap \Omega)}^2$;
- (b) $|X_\psi \cdot \eta_\psi| \leq \varepsilon |\psi|$ on $\Gamma_\psi \cap \partial\Omega$.
- (c) $\frac{1}{2} \|\psi\|_{H^1(\Gamma \cap \Omega)}^2 \leq \|\psi\|_{H^1(\Gamma_\psi \cap \Omega)}^2 \leq 2 \|\psi\|_{H^1(\Gamma \cap \Omega)}^2$.

Proof. To prove (a), we first note that given $\sigma > 0$ we can find $\delta(\sigma) \in (0, \delta_0)$ such that for every $\psi \in \mathcal{D}_{\delta(\sigma)}$ we have on Γ_ψ

$$\nu_\psi = \nu \circ \Phi_1^{-1} + \tilde{\nu} \quad \text{with} \quad \|\tilde{\nu}\|_{C^1(\Gamma_\psi)} \leq \sigma \quad (5.8)$$

and

$$\|X - X \circ \Phi_1^{-1}\|_{C^1(\Gamma_\psi)} \leq \sigma \quad (5.9)$$

(where $\Phi_1 = \Phi^\psi$, by (5.3)). Hence on Γ_ψ

$$X_\psi \cdot \nu_\psi = \psi X \cdot \nu_\psi = \psi((X \cdot \nu) \circ \Phi_1^{-1} + (X - X \circ \Phi_1^{-1}) \cdot \nu \circ \Phi_1^{-1} + X \cdot \tilde{\nu}) =: \psi(1 + R_1)$$

(where we used the fact that $(X \cdot \nu) \circ \Phi_1^{-1} = 1$), and

$$\begin{aligned} \nabla_{\Gamma_\psi}(X_\psi \cdot \nu_\psi) &= (\nabla_{\Gamma_\psi} \psi) X \cdot \nu_\psi + \psi \nabla_{\Gamma_\psi}(X \cdot \nu_\psi) \\ &= (\nabla_{\Gamma_\psi} \psi)(1 + R_1) + \psi \nabla_{\Gamma_\psi}(1 + (X - X \circ \Phi_1^{-1}) \cdot \nu \circ \Phi_1^{-1} + X \cdot \tilde{\nu}) \\ &=: (\nabla_{\Gamma_\psi} \psi)(1 + R_1) + \psi R_2. \end{aligned}$$

Recalling (5.8) and (5.9), the L^∞ -norm of R_1 and R_2 can be made as small as we want by taking σ small enough, and in turn from the previous identities we obtain (a).

To prove (b), we first observe that, by reducing $\delta(\sigma)$ if necessary, we can guarantee that for every $\psi \in \mathcal{D}_{\delta(\sigma)}$

$$\eta_\psi = \eta \circ \Phi_1^{-1} + \tilde{\eta} \quad \text{with} \quad |\tilde{\eta}| \leq \sigma \quad \text{on } \Gamma_\psi \cap \partial\Omega. \quad (5.10)$$

We deduce that on $\Gamma_\psi \cap \partial\Omega$

$$|X_\psi \cdot \eta_\psi| = |\psi X \cdot \eta_\psi| = |\psi((X \cdot \eta) \circ \Phi_1^{-1} + (X - X \circ \Phi_1^{-1}) \cdot \eta \circ \Phi_1^{-1} + X \cdot \tilde{\eta})| \leq \varepsilon |\psi|$$

where the last inequality follows by observing that $(X \cdot \eta) \circ \Phi_1^{-1} = 0$, and by (5.9) and (5.10) (choosing σ small enough, depending on ε). This proves (b).

Finally, by a change of variables (using the area formula (2.4)) we have

$$\|\psi\|_{H^1(\Gamma_\psi \cap \Omega)}^2 = \int_{\Gamma \cap \Omega} \left(|\psi \circ \Phi^\psi|^2 + \frac{|\nabla_\Gamma(\psi \circ \Phi^\psi)|^2}{|D\Phi^\psi[\tau]|^2} \right) J_{\Phi^\psi} d\mathcal{H}^1,$$

and (c) follows by (5.1) and recalling that $\psi \circ \Phi^\psi = \psi$ on Γ . \square

Given $\psi \in \mathcal{D}_{\delta_0}$, we can define a bilinear form on $H^1(\Gamma_\psi \cap \Omega)$ as in (4.17), by setting

$$(\varphi, \vartheta)_{\sim, \psi} := \int_{\Omega \cap \Gamma_\psi} \nabla_{\Gamma_\psi} \varphi \cdot \nabla_{\Gamma_\psi} \vartheta d\mathcal{H}^1 + \int_{\Omega \cap \Gamma_\psi} H_\psi^2 \varphi \vartheta d\mathcal{H}^1 - \int_{\Gamma_\psi \cap \partial\Omega} D\nu_{\partial\Omega}[\nu_\psi, \nu_\psi] \varphi \vartheta d\mathcal{H}^0.$$

The positivity assumption (3.5) guarantees that, if δ is sufficiently small, it is possible to control the H^1 -norm on Γ_ψ in terms of the norm $\|\cdot\|_{\sim, \psi}$ associated with $(\cdot, \cdot)_{\sim, \psi}$, uniformly with respect to $\psi \in \mathcal{D}_\delta$. This is the content of the following proposition, whose proof is analogous to [6, Lemma 5.3] (the only difference lies in the presence of a boundary term, which can be treated similarly to the others).

Proposition 5.4. *In the hypotheses of Theorem 5.2, there exist $C_1 > 0$ and $\delta_1 \in (0, \delta_0)$ such that for every $\psi \in \mathcal{D}_{\delta_1}$*

$$\|\varphi\|_{H^1(\Gamma_\psi \cap \Omega)} \leq C_1 \|\varphi\|_{\sim, \psi} \quad \text{for every } \varphi \in H^1(\Gamma_\psi \cap \Omega).$$

The previous result allows us to introduce, for $\psi \in \mathcal{D}_{\delta_1}$, a compact operator $T_\psi : H^1(\Gamma_\psi \cap \Omega) \rightarrow H^1(\Gamma_\psi \cap \Omega)$ defined by

$$(T_\psi \varphi, \vartheta)_{\sim, \psi} = -2 \int_{\Gamma_\psi \cap \Omega} [v_{\varphi, \psi}^+ \operatorname{div}_{\Gamma_\psi}(\vartheta \nabla_{\Gamma_\psi} u_\psi^+) - v_{\varphi, \psi}^- \operatorname{div}_{\Gamma_\psi}(\vartheta \nabla_{\Gamma_\psi} u_\psi^-)] d\mathcal{H}^1 \quad (5.11)$$

for every $\varphi, \vartheta \in H^1(\Gamma_\psi \cap \Omega)$, where $v_{\varphi, \psi} \in H_U^1(\Omega \setminus \Gamma_\psi)$ is the solution to

$$\int_{\Omega} \nabla v_{\varphi, \psi} \cdot \nabla z dx + \int_{\Gamma_\psi \cap \Omega} [z^+ \operatorname{div}_{\Gamma_\psi}(\varphi \nabla_{\Gamma_\psi} u_\psi^+) - z^- \operatorname{div}_{\Gamma_\psi}(\varphi \nabla_{\Gamma_\psi} u_\psi^-)] d\mathcal{H}^1 = 0$$

for every $z \in H_U^1(\Omega \setminus \Gamma_\psi)$. We define also $\lambda_{1, \psi}$ similarly to (4.20). The following semicontinuity property of the eigenvalues $\lambda_{1, \psi}$ will be crucial in the proof of Theorem 5.2. We omit the proof of this result, since it is the same as in [6, Lemma 5.4]: we only observe that Remark 5.1 guarantees that we have the same convergence as in [6, formula (5.14)], so that we can reproduce word by word the same argument.

Proposition 5.5. *In the hypotheses of Theorem 5.2,*

$$\limsup_{\|\psi\|_{C^2(\Gamma)} \rightarrow 0} \lambda_{1, \psi} \leq \lambda_1.$$

We are finally ready to prove the local minimality result stated at the beginning of this section.

Proof of Theorem 5.2. We divide the proof into two steps.

Step 1. We first show that there exist $\delta \in (0, \delta_1)$ and $c > 0$ such that for every $\psi \in \mathcal{D}_\delta$

$$F(\Gamma_\psi, u_\psi) \geq F(\Gamma, u) + c \|\psi\|_{H^1(\Gamma \cap \Omega)}^2. \quad (5.12)$$

Given $\psi \in \mathcal{D}_\delta$, with $\delta \in (0, \delta_1)$ to be chosen, consider the admissible flow $(\Phi_t)_t$ associated with ψ , according to (5.3), and its tangent vector field X_ψ . Setting $g_\psi(t) := F(\Gamma_{\Phi_t}, u_{\Phi_t})$, we claim that there exist $c > 0$ and $\delta > 0$ such that

$$g_\psi''(t) \geq 2c \|\psi\|_{H^1(\Gamma \cap \Omega)}^2 \quad \text{for every } t \in [0, 1] \text{ and } \psi \in \mathcal{D}_\delta. \quad (5.13)$$

Once this is proved, claim (5.12) will follow immediately: indeed, as $g_\psi'(0) = 0$ since (Γ, u) is a critical pair, and recalling that $\Gamma_{\Phi_1} = \Gamma_\psi$, we deduce

$$\begin{aligned} F(\Gamma, u) &= g_\psi(0) = g_\psi(1) - \int_0^1 (1-t) g_\psi''(t) dt \\ &\leq F(\Gamma_\psi, u_\psi) - c \|\psi\|_{H^1(\Gamma \cap \Omega)}^2, \end{aligned}$$

which is (5.12).

We now come to the proof of (5.13). In order to simplify the notation, we set $\nu_t := \nu_{\Phi_t}$, $\eta_t := \eta_{\Phi_t}$, $\Gamma_t := \Gamma_{\Phi_t}$, and $H_t := H_{\Phi_t}$. By Remark 4.7, recalling the definition of $T_{t\psi}$ (see (5.11)), we deduce that

$$\begin{aligned} g''_{\psi}(t) &= -(T_{t\psi}(X_{\psi} \cdot \nu_t), X_{\psi} \cdot \nu_t)_{\sim, t\psi} + \int_{\Gamma_t \cap \Omega} \left(H_t^2(X_{\psi} \cdot \nu_t)^2 + |\nabla_{\Gamma_t}(X_{\psi} \cdot \nu_t)|^2 \right) d\mathcal{H}^1 \\ &\quad + \int_{\Gamma_t \cap \Omega} f_t(Z_{\psi} \cdot \nu_t - 2X_{\psi}^{\parallel} \cdot \nabla_{\Gamma_t}(X_{\psi} \cdot \nu_t) + D\nu_t[X_{\psi}^{\parallel}, X_{\psi}^{\parallel}] - H_t(X_{\psi} \cdot \nu_t)^2) d\mathcal{H}^1 \\ &\quad + \int_{\Gamma_t \cap \partial\Omega} Z_{\psi} \cdot \eta_t d\mathcal{H}^0, \end{aligned}$$

where $f_t = |\nabla_{\Gamma_t} u_{\Phi_t}^-|^2 - |\nabla_{\Gamma_t} u_{\Phi_t}^+|^2 + H_t$. Since

$$\begin{aligned} 0 &= Z_{\psi} \cdot \nu_{\partial\Omega} + D\nu_{\partial\Omega}[X_{\psi}, X_{\psi}] \\ &= Z_{\psi} \cdot \nu_{\partial\Omega} + D\nu_{\partial\Omega}[\nu_t, \nu_t](X_{\psi} \cdot \nu_t)^2 + ((X_{\psi} \cdot \eta_t)^2 \eta_t + 2(X_{\psi} \cdot \nu_t)(X_{\psi} \cdot \eta_t) \nu_t) \cdot D\nu_{\partial\Omega}[\eta_t] \end{aligned}$$

on $\Gamma_t \cap \partial\Omega$ by Lemma 4.6 (d), we can rewrite $g''_{\psi}(t)$ as

$$\begin{aligned} g''_{\psi}(t) &= -(T_{t\psi}(X_{\psi} \cdot \nu_t), X_{\psi} \cdot \nu_t)_{\sim, t\psi} + \|X_{\psi} \cdot \nu_t\|_{\sim, t\psi}^2 \\ &\quad + \int_{\Gamma_t \cap \Omega} f_t(Z_{\psi} \cdot \nu_t - 2X_{\psi}^{\parallel} \cdot \nabla_{\Gamma_t}(X_{\psi} \cdot \nu_t) + D\nu_t[X_{\psi}^{\parallel}, X_{\psi}^{\parallel}] - H_t(X_{\psi} \cdot \nu_t)^2) d\mathcal{H}^1 \\ &\quad + \int_{\Gamma_t \cap \partial\Omega} Z_{\psi} \cdot (\eta_t - \nu_{\partial\Omega}) d\mathcal{H}^0 \\ &\quad - \int_{\Gamma_t \cap \partial\Omega} ((X_{\psi} \cdot \eta_t)^2 \eta_t + 2(X_{\psi} \cdot \nu_t)(X_{\psi} \cdot \eta_t) \nu_t) \cdot D\nu_{\partial\Omega}[\eta_t] d\mathcal{H}^0. \end{aligned} \quad (5.14)$$

We now carefully estimate each term in the previous expression. In the following, C will denote a generic positive constant, independent of $\psi \in \mathcal{D}_{\delta_1}$, which may change from line to line.

As (Γ, u) satisfies condition (3.5), Proposition 4.11 implies that $\lambda_1 < 1$, so that by Proposition 5.5 we can find $\delta_2 \in (0, \delta_1)$ such that for every $\psi \in \mathcal{D}_{\delta_2}$

$$\lambda_{1, \psi} < \frac{1}{2}(\lambda_1 + 1) < 1. \quad (5.15)$$

Fix $\varepsilon > 0$ to be chosen later, and let $\delta(\varepsilon) > 0$ be given by Lemma 5.3 (assume without loss of generality that $\delta(\varepsilon) < \delta_2$). We remark that, if $\psi \in \mathcal{D}_{\delta(\varepsilon)}$, then $t\psi \in \mathcal{D}_{\delta(\varepsilon)}$ for every $t \in [0, 1]$, and $X_{t\psi} = tX_{\psi}$: hence we can apply (a) and (b) of Lemma 5.3 to $t\psi$, and we easily obtain that

$$\frac{1}{2} \|\psi\|_{H^1(\Gamma \cap \Omega)}^2 \leq \|X_{\psi} \cdot \nu_t\|_{H^1(\Gamma_t \cap \Omega)}^2 \leq 2 \|\psi\|_{H^1(\Gamma \cap \Omega)}^2, \quad (5.16)$$

$$|X_{\psi} \cdot \eta_t| \leq \varepsilon |\psi| \quad \text{on } \Gamma_t \cap \partial\Omega, \quad (5.17)$$

for every $\psi \in \mathcal{D}_{\delta(\varepsilon)}$ and for every $t \in [0, 1]$.

Fix now any $\psi \in \mathcal{D}_{\delta(\varepsilon)}$. From the definition of $\lambda_{1, \psi}$ and (5.15) we have

$$\begin{aligned} &-(T_{t\psi}(X_{\psi} \cdot \nu_t), X_{\psi} \cdot \nu_t)_{\sim, t\psi} + \|X_{\psi} \cdot \nu_t\|_{\sim, t\psi}^2 \geq (1 - \lambda_{1, t\psi}) \|X_{\psi} \cdot \nu_t\|_{\sim, t\psi}^2 \\ &\geq \frac{1 - \lambda_1}{2} \|X_{\psi} \cdot \nu_t\|_{\sim, t\psi}^2 \geq \frac{1 - \lambda_1}{2C_1^2} \|X_{\psi} \cdot \nu_t\|_{H^1(\Gamma_t \cap \Omega)}^2 \geq \frac{1 - \lambda_1}{4C_1^2} \|\psi\|_{H^1(\Gamma \cap \Omega)}^2, \end{aligned} \quad (5.18)$$

where the last two inequalities follow from Proposition 5.4 and from (5.16).

By Remark 5.1 the map

$$\psi \in \mathcal{D}_{\delta(\varepsilon)} \mapsto \| |\nabla_{\Gamma_{\psi}} u_{\psi}^-|^2 - |\nabla_{\Gamma_{\psi}} u_{\psi}^+|^2 + H_{\psi} \|_{L^{\infty}(\Gamma_{\psi} \cap \Omega)}$$

is continuous with respect to the C^2 -topology; hence, as it vanishes for $\psi = 0$ by (3.2), possibly reducing $\delta(\varepsilon)$ we have that for every $\psi \in \mathcal{D}_{\delta(\varepsilon)}$

$$\| |\nabla_{\Gamma_{\psi}} u_{\psi}^-|^2 - |\nabla_{\Gamma_{\psi}} u_{\psi}^+|^2 + H_{\psi} \|_{L^{\infty}(\Gamma_{\psi} \cap \Omega)} < \varepsilon.$$

We deduce that

$$\begin{aligned}
& \int_{\Gamma_t \cap \Omega} f_t(Z_\psi \cdot \nu_t - 2X_\psi^\parallel \cdot \nabla_{\Gamma_t}(X_\psi \cdot \nu_t) + D\nu_t[X_\psi^\parallel, X_\psi^\parallel] - H_t(X_\psi \cdot \nu_t)^2) d\mathcal{H}^1 \\
& \geq -\varepsilon \|Z_\psi \cdot \nu_t - 2X_\psi^\parallel \cdot \nabla_{\Gamma_t}(X_\psi \cdot \nu_t) + D\nu_t[X_\psi^\parallel, X_\psi^\parallel] + H_t(X_\psi \cdot \nu_t)^2\|_{L^1(\Gamma_t \cap \Omega)} \\
& \geq -\varepsilon C \left(\|\psi\|_{L^2(\Gamma_t \cap \Omega)}^2 + \|\nabla_{\Gamma_t}(X_\psi \cdot \nu_t)\|_{L^2(\Gamma_t \cap \Omega)} \|\psi\|_{L^2(\Gamma_t \cap \Omega)} \right) \\
& \geq -\varepsilon C \|\psi\|_{H^1(\Gamma \cap \Omega)}^2,
\end{aligned} \tag{5.19}$$

where we used also (5.7), (5.16), and (c) of Lemma 5.3.

By reducing $\delta(\varepsilon)$ if necessary we can assume

$$\max_{x \in \Gamma_\psi \cap \partial\Omega} |\eta_\psi(x) - \nu_{\partial\Omega}(x)| < \varepsilon \quad \text{for every } \psi \in \mathcal{D}_{\delta(\varepsilon)},$$

so that using again (5.7) and (c) of Lemma 5.3 we obtain

$$\int_{\Gamma_t \cap \partial\Omega} Z_\psi \cdot (\eta_t - \nu_{\partial\Omega}) d\mathcal{H}^0 \geq -\varepsilon C \int_{\Gamma_t \cap \partial\Omega} \psi^2 d\mathcal{H}^0 \geq -\varepsilon C \|\psi\|_{H^1(\Gamma \cap \Omega)}^2. \tag{5.20}$$

Finally, we proceed in a similar way to estimate the last integral in (5.14): by (5.7) and (5.17)

$$-\int_{\Gamma_t \cap \partial\Omega} ((X_\psi \cdot \eta_t)^2 \eta_t + 2(X_\psi \cdot \nu_t)(X_\psi \cdot \eta_t) \nu_t) \cdot D\nu_{\partial\Omega}[\eta_t] d\mathcal{H}^0 \geq -\varepsilon C \|\psi\|_{H^1(\Gamma \cap \Omega)}^2. \tag{5.21}$$

Collecting (5.18)–(5.21), by (5.14) we conclude that for every $\psi \in \mathcal{D}_{\delta(\varepsilon)}$ and for every $t \in [0, 1]$

$$g_\psi''(t) \geq \left(\frac{1 - \lambda_1}{4C_1^2} - \varepsilon C \right) \|\psi\|_{H^1(\Gamma \cap \Omega)}^2$$

for some positive constant C (independent of ψ), so that by choosing ε sufficiently small we obtain the claim (5.13) and, in turn, (5.12).

Step 2. The conclusion of the theorem follows now by approximation: given any $\psi \in W^{2,\infty}(\Gamma \cap \Omega)$ with $\|\psi\|_{W^{2,\infty}(\Gamma \cap \Omega)} < \delta$, we can find a sequence $\psi_n \in \mathcal{D}_\delta$ converging to ψ in $W^{1,\infty}(\Gamma \cap \Omega)$ for which the conclusion obtained in the previous step holds:

$$F(\Gamma_{\psi_n}, u_{\psi_n}) \geq F(\Gamma, u) + c \|\psi_n\|_{H^1(\Gamma \cap \Omega)}^2.$$

Noting that $F(\Gamma_{\psi_n}, u_{\psi_n}) \rightarrow F(\Gamma_\psi, u_\psi)$ as a consequence of the $W^{1,\infty}$ -convergence of ψ_n to ψ , by passing to the limit in the previous inequality we conclude that the same estimate holds for ψ . Hence the conclusion of the theorem follows by recalling that $F(\Gamma_\psi, v) \geq F(\Gamma_\psi, u_\psi)$ for every $v \in H^1(\Omega \setminus \Gamma_\psi)$ with $v = u$ in $(\Omega \setminus U) \cup \partial_D \Omega$, by definition of u_ψ . \square

6. LOCAL $C^{1,\alpha}$ -MINIMALITY

In this section we show that the $W^{2,\infty}$ -minimality property proved in the previous section implies that (Γ, u) is also a minimizer with respect to small $C^{1,\alpha}$ -perturbations of the discontinuity set. We start by a preliminary construction that will be needed in the proof.

Remark 6.1. Let X be the vector field defined at the beginning of Section 5, which, we recall, coincides with ν on Γ and is tangent to $\partial\Omega$, and let Ψ be the flow generated by X . We want to define a one-parameter family of smooth curves $(\Gamma_\delta)_\delta$, for $\delta \in (-\delta_0, \delta_0)$, with $\Gamma_0 = \Gamma$, such that X is normal to each curve of the family, and whose union is a tubular neighborhood of Γ . In order to do this, let $x_0 \in \Gamma \cap \partial\Omega$ and let $x_\delta := \Psi(\delta, x_0)$. We then define Γ_δ as the trajectory of the flow generated by X^\perp starting from x_δ , where the vector field X^\perp is obtained by a rotation of X by $\frac{\pi}{2}$. This construction provides a family of curves with the desired properties.

We can then define a family of tubular neighborhoods of Γ in Ω whose boundaries meet $\partial\Omega$ orthogonally, by setting for $\delta \in (-\delta_0, \delta_0)$

$$\mathcal{I}_\delta(\Gamma) := \bigcup_{|s| < \delta} \Gamma_s.$$

Proposition 6.2. *Let (Γ, u) be a strictly stable regular critical pair in an admissible subdomain U , and let $\alpha \in (0, 1)$. There exists $\delta > 0$ such that*

$$F(\Gamma, u) < F(\Phi(\Gamma), v)$$

for every diffeomorphism $\Phi \in C^{1,\alpha}(\overline{\Omega}; \overline{\Omega})$ with $0 < \|\Phi - Id\|_{C^{1,\alpha}(\overline{\Omega})} \leq \delta$ and $\text{supp}(\Phi - Id) \subset\subset U \setminus \partial_D \Omega$, and for every $v \in H^1(\Omega \setminus \Phi(\Gamma))$ with $v = u$ in $(\Omega \setminus U) \cup \partial_D \Omega$.

Proof. Assume by contradiction that there exist sequences $\sigma_n \rightarrow 0$ and $\Phi_n \in C^{1,\alpha}(\overline{\Omega}; \overline{\Omega})$, with

$$\text{supp}(\Phi_n - Id) \subset\subset U \setminus \partial_D \Omega, \quad 0 < \|\Phi_n - Id\|_{C^{1,\alpha}(\overline{\Omega})} \leq \sigma_n,$$

such that $F(\Phi_n(\Gamma), u_n) \leq F(\Gamma, u)$, where $u_n := u_{\Phi_n}$ is defined as in (4.1). Notice that, arguing as in Remark 5.1, we have that u_n^\pm are of class $C^{1,\alpha}$ up to $\Phi_n(\Gamma)$, and

$$\|\nabla(u_n^\pm \circ \Phi_n) - \nabla u^\pm\|_{L^\infty(\Omega^\pm)} \rightarrow 0.$$

We first extend u^+ and u^- to $C^{1,\alpha}$ -functions in Ω^- and Ω^+ , respectively, by using [13, Theorem 6.2.5]. We similarly extend $u_n^\pm \circ \Phi_n$ to $C^{1,\alpha}$ -functions \tilde{u}_n^\pm in Ω^\mp , and we set $v_n^\pm := \tilde{u}_n^\pm \circ \Phi_n^{-1}$: since the extension operator constructed in [13, Theorem 6.2.5] is continuous, we have that

$$\|\nabla v_n^\pm - \nabla u^\pm\|_{L^\infty(\Omega)} \leq \delta_n$$

for some $\delta_n \rightarrow 0$. Finally, as $\|\Phi_n - Id\|_{C^{1,\alpha}} \rightarrow 0$, we can also assume that $\Phi_n(\Gamma) \subset \mathcal{I}_{\delta_n}(\Gamma)$.

Consider the following obstacle problems

$$\min \left\{ J(E, v^+, v^-) : E \subset \Omega, \Omega^+ \setminus \mathcal{I}_{\delta_n}(\Gamma) \subset E \subset \Omega^+ \cup \mathcal{I}_{\delta_n}(\Gamma), v^\pm - u^\pm \in W^{1,\infty}(\Omega), \right. \\ \left. v^+ \chi_E + v^- \chi_{E^c} = u \text{ in } (\Omega \setminus U) \cup \partial_D \Omega, \|\nabla v^\pm - \nabla u^\pm\|_{L^\infty(\Omega)} \leq 1 \right\}, \quad (6.1)$$

where

$$J(E, v^+, v^-) := \int_E |\nabla v^+|^2 + \int_{\Omega \setminus E} |\nabla v^-|^2 + P(E, \Omega),$$

and let (F_n, w_n^+, w_n^-) be a solution to (6.1) (which exists by the direct method of the Calculus of Variations). Since $(\Phi_n(\Omega^+), v_n^+, v_n^-)$ is an admissible competitor, we deduce that

$$J(F_n, w_n^+, w_n^-) \leq J(\Phi_n(\Omega^+), v_n^+, v_n^-) = F(\Phi_n(\Gamma), u_n) \leq F(\Gamma, u). \quad (6.2)$$

We now divide the proof into three steps.

Step 1. We claim that, if $\gamma > 0$ is sufficiently large (independently of n), then (F_n, w_n^+, w_n^-) is also a solution to the penalized problem (without obstacle)

$$\min \left\{ \tilde{J}(E, v^+, v^-) : E \subset \Omega, v^\pm - u^\pm \in W^{1,\infty}(\Omega), \|\nabla v^\pm - \nabla u^\pm\|_{L^\infty(\Omega)} \leq 1, \right. \\ \left. v^\pm = u \text{ in } (\Omega^\pm \setminus U) \cup (\partial_D \Omega \cap \overline{\Omega}^\pm) \right\}, \quad (6.3)$$

where

$$\tilde{J}(E, v^+, v^-) := \int_E |\nabla v^+|^2 + \int_{\Omega \setminus E} |\nabla v^-|^2 + P(E, \Omega) + \gamma |E \Delta T_n(E)|$$

and $T_n(E) := E \cup (\Omega^+ \setminus \mathcal{I}_{\delta_n}(\Gamma)) \cap (\Omega^+ \cup \mathcal{I}_{\delta_n}(\Gamma))$.

In order to prove the claim, fix any competitor (F, w^+, w^-) for problem (6.3). We denote by ν_E the generalized inner normal to a finite perimeter set E . Since $\nu_{T_n(F)} = X$ almost everywhere on $\partial^* T_n(F) \cap \Gamma_{\delta_n}$, and $|X| \leq 1$, we can estimate the difference of the perimeters of F and $T_n(F)$ in Ω^+ as follows:

$$\begin{aligned} P(F, \Omega^+) - P(T_n(F), \Omega^+) &= \int_{(\partial^* F \setminus \partial^* T_n(F)) \cap \Omega^+} d\mathcal{H}^1 - \int_{(\partial^* T_n(F) \setminus \partial^* F) \cap \Omega^+} d\mathcal{H}^1 \\ &\geq \int_{(\partial^* F \setminus \partial^* T_n(F)) \cap \Omega^+} X \cdot \nu_F d\mathcal{H}^1 - \int_{(\partial^* T_n(F) \setminus \partial^* F) \cap \Omega^+} X \cdot \nu_{T_n(F)} d\mathcal{H}^1 \\ &= \int_{(F \Delta T_n(F)) \cap \Omega^+} \text{div} X \geq -\|\text{div} X\|_\infty |(F \Delta T_n(F)) \cap \Omega^+|, \end{aligned}$$

where we used the divergence theorem taking into account that $X \cdot \nu_{\partial\Omega} = 0$ on $\partial\Omega$. A similar estimate holds in Ω^- , and we conclude that

$$P(F, \Omega) - P(T_n(F), \Omega) \geq -\|\operatorname{div} X\|_\infty |F \Delta T_n(F)|.$$

Concerning the volume terms, since ∇w^\pm are uniformly bounded in L^∞ by a constant Λ depending only on $\|\nabla u\|_\infty$ we have

$$\left| \int_F |\nabla w^+|^2 - \int_{T_n(F)} |\nabla w^+|^2 \right| \leq \Lambda^2 |F \Delta T_n(F)|,$$

and a similar estimate holds for w^- in the complements of the sets F and $T_n(F)$. Hence we deduce by minimality of (F_n, w_n^+, w_n^-)

$$\begin{aligned} \tilde{J}(F, w^+, w^-) - \tilde{J}(F_n, w_n^+, w_n^-) &\geq J(F, w^+, w^-) - J(T_n(F), w^+, w^-) + \gamma |F \Delta T_n(F)| \\ &\geq (\gamma - 2\Lambda^2 - \|\operatorname{div} X\|_\infty) |F \Delta T_n(F)| \geq 0 \end{aligned}$$

if $\gamma > 2\Lambda^2 + \|\operatorname{div} X\|_\infty$. This shows that (F_n, w_n^+, w_n^-) is also a solution to (6.3).

Step 2. Each set F_n is a quasi-minimizer of the area functional in Ω , that is there exists a constant $\omega > 0$ (independent of n) such that for every set $F \subset \Omega$ of finite perimeter with $F \Delta F_n \subset\subset B_r(x)$ ($x \in \mathbb{R}^2$, $r > 0$) we have

$$P(F_n, \Omega) \leq P(F, \Omega) + \omega |F \Delta F_n|. \quad (6.4)$$

This can be directly deduced from the fact that (F_n, w_n^+, w_n^-) solves (6.3), using in particular the L^∞ bound on ∇w_n^\pm to estimate the Dirichlet integrals by $|F \Delta F_n|$.

Combining this information with the Hausdorff convergence of F_n to Ω^+ (whose boundary inside Ω is regular), we deduce that each F_n has $C^{1, \frac{1}{2}}$ boundary inside Ω (for n sufficiently large) which converges to Γ in the $C^{1, \alpha}$ -sense for all $\alpha \in (0, \frac{1}{2})$. This is a well-known consequence of the classical regularity theory of quasi-minimizers of the area functional, which is stated in our precise setting in [15, Theorem 3.5] (see also the references contained therein). In particular, the regularity up to the boundary $\partial\Omega$ follows from a work by Grüter [14], which guarantees in addition that the intersection of $\overline{\partial F_n} \cap \overline{\Omega}$ with $\partial\Omega$ is orthogonal.

Hence there exist diffeomorphisms $\Psi_n : \overline{\Omega} \rightarrow \overline{\Omega}$ of class $C^{1, \alpha}$ such that $F_n = \Psi_n(\Omega^+)$, $\overline{\partial F_n} \cap \overline{\Omega} = \Psi_n(\Gamma)$ and $\|\Psi_n - Id\|_{C^{1, \alpha}(\Gamma)} \rightarrow 0$. In turn, by Lemma 9.1 we conclude that $\overline{\partial F_n} \cap \overline{\Omega} = \Gamma_{\psi_n}$ for some functions $\psi_n \in C^{1, \alpha}(\Gamma)$ such that $\psi_n \rightarrow 0$ in $C^{1, \alpha}(\Gamma)$.

We also remark that ∇w_n^\pm are Hölder continuous up to Γ_{ψ_n} , and they converge uniformly to ∇u^\pm . Indeed, by considering the Dirichlet minimizer u_{Ψ_n} in $\Omega \setminus \Psi_n(\Gamma)$ under the usual boundary conditions, we have by elliptic regularity (as in Remark 5.1) that $\nabla_\Gamma(u_{\Psi_n}^\pm \circ \Psi_n)$ is Hölder continuous and converges uniformly to $\nabla_\Gamma u^\pm$. Hence, for n large enough, and also taking into account the continuity of the extension operator, u_{Ψ_n} satisfies the constraint $\|\nabla u_{\Psi_n}^\pm - \nabla u^\pm\|_\infty \leq 1$ so that we conclude that $w_n^\pm = u_{\Psi_n}^\pm$.

Step 3. By the quasi-minimality property (6.4) and the $C^{1, \alpha}$ -regularity of Γ_{ψ_n} , we deduce by a standard argument (see, e.g., [15, Proposition 3.2]) that the curvatures H_{ψ_n} of the sets Γ_{ψ_n} are uniformly bounded by the constant ω . In turn, this provides the $W^{2, \infty}$ -regularity of Γ_{ψ_n} .

If we now write the Euler-Lagrange equations for problem (6.1), we get

$$H_{\psi_n} = \begin{cases} |\nabla w_n^+|^2 - |\nabla w_n^-|^2 & \text{on } \Gamma_{\psi_n} \cap \mathcal{I}_{\delta_n}(\Gamma), \\ H_{\Gamma_{\pm\delta_n}} & \text{on } \Gamma_{\psi_n} \cap \Gamma_{\pm\delta_n}, \end{cases}$$

where $H_{\Gamma_{\pm\delta_n}}$ denotes the curvature of the curve $\Gamma_{\pm\delta_n}$. Moreover, as (Γ, u) is a critical pair, by (3.2) we have

$$H_\Gamma = |\nabla u^+|^2 - |\nabla u^-|^2 \quad \text{on } \Gamma.$$

Hence, by the uniform convergence of ∇w_n^\pm to ∇u^\pm and observing that the curvature $H_{\Gamma_{\pm\delta_n}}$ is uniformly close to H_Γ by the regularity of the flow generating the family of curves $(\Gamma_\delta)_\delta$, we deduce that

$$\|H_{\psi_n} \circ \Psi_n - H_\Gamma\|_{L^\infty(\Gamma)} \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

which implies, by Lemma 9.1, that $\|\psi_n\|_{W^{2, \infty}(\Gamma)} \rightarrow 0$.

We can conclude that, setting $w_n := w_n^+ \chi_{F_n} + w_n^- \chi_{F_n^c}$, by (6.2)

$$F(\Gamma_{\psi_n}, w_n) = J(F_n, w_n^+, w_n^-) \leq F(\Gamma, u),$$

which implies, by the isolated local $W^{2,\infty}$ -minimality of (Γ, u) proved in Theorem 5.2, that for all n large enough $\psi_n = 0$ and $w_n = u$. As a consequence, $(\Phi_n(\Omega^+), v_n^+, v_n^-)$ is itself a solution to (6.1): by repeating all the previous argument for this sequence instead of (F_n, w_n^+, w_n^-) , we conclude as before that $\Phi_n = Id$, which is the desired contradiction. \square

7. PROOF OF THEOREM 3.7

In this section we complete the proof of Theorem 3.7. It will be useful to introduce the relaxed functional

$$\mathcal{F}(u) := \int_{\Omega} |\nabla u|^2 dx + \mathcal{H}^1(S_u) \quad \text{for } u \in SBV(\Omega)$$

and, for $B \subset \Omega$ Borel set, its local version

$$\mathcal{F}(u; B) := \int_{\Omega \cap B} |\nabla u|^2 dx + \mathcal{H}^1(S_u \cap B).$$

Remark 7.1. We recall here that if $(K, v) \in \mathcal{A}(\Omega)$ is an admissible pair with $\mathcal{H}^{N-1}(K) < +\infty$ and $v \in L^\infty(\Omega)$, then the function v is in $SBV(\Omega)$ and satisfies $\mathcal{H}^1(S_v \setminus K) = 0$ (see [2, Proposition 4.4]); in particular, we have $\mathcal{F}(v) \leq F(K, v)$. On the other hand, if (Γ, u) is a regular critical pair, then $u \in SBV(\Omega)$, $S_u = \Gamma$ and $\mathcal{F}(u) = F(\Gamma, u)$.

We can now start the proof of Theorem 3.7. By Remark 4.16 we are allowed to perform the proof under the additional assumption that U has boundary of class C^∞ intersecting $\partial\Omega$ orthogonally. Moreover, from Remark 7.1 it follows that in order to obtain the result it is sufficient to show that there exists $\delta > 0$ such that $\mathcal{F}(u) < \mathcal{F}(v)$ for every $v \in SBV(\Omega)$ with $v = u$ in $(\Omega \setminus U) \cup \partial_D\Omega$ and $0 < \|v - u\|_{L^1(\Omega)} < \delta$.

Hence we assume by contradiction that there exists a sequence $v_n \in SBV(\Omega)$, with $v_n = u$ in $(\Omega \setminus U) \cup \partial_D\Omega$, such that $0 < \|v_n - u\|_{L^1(\Omega)} \rightarrow 0$ and

$$\mathcal{F}(v_n) \leq \mathcal{F}(u). \quad (7.1)$$

By a truncation argument, we can assume that $\|v_n\|_{L^\infty(\Omega)} \leq \|u\|_{L^\infty(\Omega)} =: M < +\infty$.

We introduce a bounded open set Ω' such that $\Omega \subset \Omega'$ and $\Omega' \cap \partial\Omega = \partial_D\Omega$, in order to enforce the boundary condition on $\partial_D\Omega$. We can extend u in $\Omega' \setminus \Omega$ to a function $u \in SBV(\Omega')$ such that $\mathcal{H}^1(S_u \cap \partial_D\Omega) = 0$ and $\|u\|_{L^\infty(\Omega')} \leq M$. Moreover, we can also assume that $v_n \in SBV(\Omega')$ and $v_n = u$ in $\Omega' \setminus (U \cap \Omega)$. In particular, $\mathcal{H}^1(S_{v_n} \cap \partial_D\Omega) = 0$ and hence $\mathcal{F}(v_n; \Omega') \leq \mathcal{F}(u; \Omega')$.

We set $\varepsilon_n := \|v_n - u\|_{L^2(\Omega)}^2 \rightarrow 0$,

$$h_n(t) := \begin{cases} \sqrt{(t - \varepsilon_n)^2 + \varepsilon_n^2} - \varepsilon_n, & \text{if } t > \varepsilon_n, \\ 0, & \text{if } 0 \leq t \leq \varepsilon_n, \end{cases}$$

and we consider, for $\beta > 0$ to be chosen later, a solution w_n to the following penalized minimum problem:

$$\min \left\{ \mathcal{F}(w; \Omega') + \beta h_n(\|w - u\|_{L^2(\Omega)}^2) : w \in SBV(\Omega'), w = u \text{ in } \Omega' \setminus (U \cap \Omega) \right\}. \quad (7.2)$$

The existence of a solution to (7.2) is guaranteed by the lower semi-continuity and compactness theorems in SBV (see [2, Theorem 4.7 and Theorem 4.8]), and we can also assume $\|w_n\|_{L^\infty(\Omega')} \leq M$. Observe in addition that, by (7.1) and by minimality of w_n , we have

$$\mathcal{F}(w_n; \Omega') \leq \mathcal{F}(v_n; \Omega') + \beta h_n(\|w_n - u\|_{L^2(\Omega)}^2) \leq \mathcal{F}(v_n; \Omega') \leq \mathcal{F}(u; \Omega'). \quad (7.3)$$

In particular the energies $\mathcal{F}(w_n; \Omega')$ are equibounded, and in turn, again by the compactness and lower semi-continuity theorems in SBV , up to subsequences w_n converges in $SBV(\Omega')$ (see Definition 2.1) and in $L^p(\Omega')$ for every $p \in [1, +\infty)$ to a function $z \in SBV(\Omega')$ which solves the minimum problem

$$\min \left\{ \mathcal{F}(w; \Omega') + \beta \int_{\Omega} |w - u|^2 dx : w \in SBV(\Omega'), w = u \text{ in } \Omega' \setminus (U \cap \Omega) \right\}. \quad (7.4)$$

Indeed, if $w \in SBV(\Omega')$ is an admissible function for problem (7.4), then by semi-continuity and by minimality of w_n we immediately deduce that

$$\begin{aligned} \mathcal{F}(z; \Omega') + \beta \int_{\Omega} |z - u|^2 dx &\leq \liminf_{n \rightarrow \infty} \left(\mathcal{F}(w_n; \Omega') + \beta h_n(\|w_n - u\|_{L^2(\Omega)}^2) \right) \\ &\leq \liminf_{n \rightarrow \infty} \left(\mathcal{F}(w; \Omega') + \beta h_n(\|w - u\|_{L^2(\Omega)}^2) \right). \end{aligned}$$

By the result in [21], based on the construction of a suitable calibration, we can identify the solution to the limit problem.

Proposition 7.2. *If β is sufficiently large, then the unique solution to (7.4) is u itself.*

Notice that in [21] only pure Neumann boundary conditions are considered (i.e., $\partial_D \Omega = \emptyset$). Nevertheless, exactly the same construction applies to our setting without any change (see also [20, Remark 4.3.5]).

Hence, by choosing $\beta > 0$ sufficiently large, we have that $w_n \rightarrow u$ in $SBV(\Omega')$. In addition, by lower semi-continuity of \mathcal{F} and by (7.3) we deduce that $\mathcal{F}(w_n; \Omega') \rightarrow \mathcal{F}(u; \Omega')$ as $n \rightarrow \infty$, which combined with the lower semi-continuity of the two terms in the functional (which holds separately, by [2, Theorem 4.7]) yields

$$\lim_{n \rightarrow +\infty} \int_{\Omega'} |\nabla w_n|^2 dx = \int_{\Omega'} |\nabla u|^2 dx, \quad \lim_{n \rightarrow +\infty} \mathcal{H}^1(S_{w_n}) = \mathcal{H}^1(S_u). \quad (7.5)$$

In the following lemma we localize the previous convergence in open sets and we prove a continuity property that will be used subsequently.

Lemma 7.3. *For every open set $A \subset \mathbb{R}^2$ such that $|\partial A| = 0$ and $\mathcal{H}^1(S_u \cap \partial A) = 0$ we have*

$$\int_{\Omega' \cap A} |\nabla w_n|^2 dx \rightarrow \int_{\Omega' \cap A} |\nabla u|^2 dx, \quad \mathcal{H}^1(S_{w_n} \cap A) \rightarrow \mathcal{H}^1(S_u \cap A)$$

as $n \rightarrow +\infty$. Moreover, for every bounded continuous function $f \in C^0(\Omega')$ we have

$$\int_{S_{w_n} \cap A} f d\mathcal{H}^1 \rightarrow \int_{S_u \cap A} f d\mathcal{H}^1.$$

Proof. The first part of the statement follows easily from the lower semi-continuity of both terms in the functional, which holds in every open set, combined with (7.5). To prove the second part, fix any continuous and bounded function $f : \Omega' \rightarrow \mathbb{R}$. Assuming without loss of generality that $f \geq 0$ (for the general case, one can split f into positive and negative parts), we have to show that

$$\int_0^{\max f} \mathcal{H}^1(S_{w_n} \cap A \cap \{f > t\}) dt \rightarrow \int_0^{\max f} \mathcal{H}^1(S_u \cap A \cap \{f > t\}) dt.$$

The sets $A_t = \{f > t\}$ are open and they satisfy $|\partial A_t| = 0$, $\mathcal{H}^1(S_u \cap \partial A_t) = 0$ for all except at most for countable many t . Then, by the assumptions on A , the same is true for the sets $A \cap A_t$, and hence by the first part of the lemma we have

$$\mathcal{H}^1(S_{w_n} \cap A \cap \{f > t\}) \rightarrow \mathcal{H}^1(S_u \cap A \cap \{f > t\}) \quad \text{for a.e. } t \in (0, \max f),$$

and by the Dominated Convergence Theorem we obtain the conclusion. \square

In the following proposition we show that w_n satisfies a quasi-minimality property. This is an essential step in our strategy to prove Theorem 3.7: indeed, as a consequence of the regularity theory for quasi-minimizers of the Mumford-Shah functional we obtain firstly that a uniform lower bound on the 1-dimensional density of S_{w_n} holds, and moreover we will be able to deduce the $C^{1,\alpha}$ -convergence of S_{w_n} to S_u (see Proposition 7.6).

Proposition 7.4. *Each function w_n is a quasi-minimizer of the Mumford-Shah functional, that is, there exists a positive constant ω (independent of n) such that if $x \in \overline{\Omega}'$ and $\rho > 0$ then*

$$\mathcal{F}(w_n; B_\rho(x) \cap \Omega') \leq \mathcal{F}(v; B_\rho(x) \cap \Omega') + \omega \rho^2 \quad (7.6)$$

for every $v \in SBV(\Omega')$ with $v = u$ in $\Omega' \setminus (U \cap \Omega)$ and $\{v \neq w_n\} \subset\subset B_\rho(x)$.

Proof. Let v be as in the statement, and set $v^M := (-M) \vee (v \wedge M)$ (where $M = \|u\|_\infty$). Then, since $v^M \in SBV(\Omega')$ is an admissible competitor in problem (7.2), $\{v^M \neq w_n\} \subset \{v \neq w_n\}$ (as $\|w_n\|_\infty \leq M$) and $\mathcal{F}(v^M) \leq \mathcal{F}(v)$, we have by minimality of w_n

$$\begin{aligned} \mathcal{F}(w_n; B_\rho(x) \cap \Omega') &\leq \mathcal{F}(v^M; B_\rho(x) \cap \Omega') + \beta h_n \left(\int_\Omega |v^M - u|^2 dy \right) - \beta h_n \left(\int_\Omega |w_n - u|^2 dy \right) \\ &\leq \mathcal{F}(v; B_\rho(x) \cap \Omega') + \beta \left| \int_{B_\rho(x) \cap \Omega} |v^M - u|^2 dy - \int_{B_\rho(x) \cap \Omega} |w_n - u|^2 dy \right| \\ &\leq \mathcal{F}(v; B_\rho(x) \cap \Omega') + 8M^2 \beta \pi \rho^2, \end{aligned}$$

where we used the fact that h_n is 1-Lipschitz in the second inequality. Hence (7.6) follows by choosing $\omega := 8M^2 \beta \pi$. \square

Corollary 7.5. *Each set S_{w_n} is essentially closed: $\mathcal{H}^1(\overline{S_{w_n}} \setminus S_{w_n}) = 0$. Moreover, the sets $\overline{S_{w_n}}$ converge to $\overline{S_u}$ in $\overline{\Omega'}$ in the sense of Kuratowski:*

- (i) for every $x_n \in \overline{S_{w_n}}$ such that $x_n \rightarrow x$, then $x \in \overline{S_u}$;
- (ii) for every $x \in \overline{S_u}$ there exist $x_n \in \overline{S_{w_n}}$ such that $x_n \rightarrow x$.

Proof. Thanks to the quasi-minimality property proved in the previous proposition and to the fact that ∂U and $\partial \Omega$ meet orthogonally, we can apply Theorem 2.4 to infer the existence of constants $\vartheta_0 > 0$, $\rho_0 > 0$ (independent of n) such that for every $x \in \overline{S_{w_n}} \cap \overline{\Omega'}$ and for every $\rho \leq \rho_0$

$$\mathcal{H}^1(S_{w_n} \cap B_\rho(x)) \geq \vartheta_0 \rho. \quad (7.7)$$

The properties in the statement are standard consequences of (7.7) (see [2]). \square

Corollary 7.5 provides the Hausdorff convergence of $\overline{S_{w_n}}$ to $\overline{S_u}$ in $\overline{\Omega'}$, which allows us to assume, from now on, that $\overline{S_{w_n}}$ is contained in a tubular neighborhood of $\overline{S_u}$ contained in U . We now come to the main consequence of the regularity theory for quasi-minimizers. We follow here the notation introduced in Section 2.2.

We first observe that, using the good description of $\overline{S_{w_n}}$ near $\partial \Omega$ given by Theorem 2.3, we can find $\tau > 0$ such that $\overline{S_{w_n}} \cap \Omega(\tau)$ is a $C^{1,\alpha}$ -curve for some $\alpha \in (0, 1)$, with $C^{1,\alpha}$ -norms uniformly bounded with respect to n and meeting $\partial \Omega$ orthogonally. Combining this information with the Hausdorff convergence to $\overline{S_u}$, we deduce that the sets $\overline{S_{w_n}}$ converge to $\overline{S_u}$ in $\Omega(\tau)$ in the $C^{1,\beta}$ -sense, for every $\beta < \alpha$. In the following proposition we obtain the same convergence in the interior of Ω .

Proposition 7.6. *There exists a finite covering of $\Gamma \cap (\Omega \setminus \Omega(\tau))$ of the form $\bigcup_{i=1}^{N_0} (x_i + C_{\nu_i, \rho_i})$ where $x_i \in \Gamma$, $\nu_i = \nu_\Gamma(x_i)$, and functions $f_i^{(n)} : (-\rho_i, \rho_i) \rightarrow (-\rho_i, \rho_i)$ of class $C^{1,\alpha}$ (for some $\alpha \in (0, 1)$) such that*

$$(\overline{S_{w_n}} - x_i) \cap C_{\nu_i, \rho_i} = \text{gr}_{\nu_i}(f_i^{(n)})$$

for n sufficiently large and $i = 1, \dots, N_0$. Moreover, the sequence $f_i^{(n)}$ converges to f_i in $C^{1,\beta}$ as $n \rightarrow +\infty$ for every $\beta < \alpha$, where $f_i : (-\rho_i, \rho_i) \rightarrow (-\rho_i, \rho_i)$ is such that

$$(\Gamma - x_i) \cap C_{\nu_i, \rho_i} = \text{gr}_{\nu_i}(f_i).$$

Proof. Fix any point $x_0 \in \Gamma \cap (\Omega \setminus \Omega(\tau))$. By the regularity of u and $\Gamma = \overline{S_u}$, we can find $r_0 > 0$ such that $B_{r_0}(x_0) \subset \Omega \cap U$, $\mathcal{H}^1(S_u \cap \partial B_{r_0}(x_0)) = 0$ and

$$E_u(x_0, r_0) < \varepsilon_0 \frac{r_0}{8},$$

where ε_0 is given by Theorem 2.2. Lemma 7.3 immediately implies that $D_{w_n}(x_0, r_0) \rightarrow D_u(x_0, r_0)$ and that for every affine plane T

$$\int_{\overline{S_{w_n}} \cap B_{r_0}(x_0)} \text{dist}^2(y, T) d\mathcal{H}^1(y) \rightarrow \int_{\overline{S_u} \cap B_{r_0}(x_0)} \text{dist}^2(y, T) d\mathcal{H}^1(y).$$

From the previous convergence it follows also that $\limsup_{n \rightarrow \infty} A_{w_n}(x_0, r_0) \leq A_u(x_0, r_0)$, since if the minimum value defining $A_u(x_0, r_0)$ is attained at an affine plane T_0 , then

$$A_{w_n}(x_0, r_0) \leq \int_{\bar{S}_{w_n} \cap B_{r_0}(x_0)} \text{dist}^2(y, T_0) d\mathcal{H}^1(y) \rightarrow \int_{\bar{S}_u \cap B_{r_0}(x_0)} \text{dist}^2(y, T_0) d\mathcal{H}^1(y) = A_u(x_0, r_0).$$

Hence $\limsup_{n \rightarrow \infty} E_{w_n}(x_0, r_0) \leq E_u(x_0, r_0)$, so that for n sufficiently large we have

$$E_{w_n}(x_0, r_0) < \varepsilon_0 \frac{r_0}{8}.$$

By Corollary 7.5 we can find a sequence $x_n \in \bar{S}_{w_n}$ converging to x_0 , so that $B_{r_0/2}(x_n) \subset B_{r_0}(x_0)$ for n large enough and thus

$$E_{w_n}(x_n, r_0/2) = D_{w_n}(x_n, r_0/2) + \frac{4}{r_0^2} A_{w_n}(x_n, r_0/2) \leq 4E_{w_n}(x_0, r_0) < \varepsilon_0 \frac{r_0}{2}.$$

We can then apply Theorem 2.2: we find a radius $r_1 \in (0, r_0)$ and functions $g_n : (-r_1, r_1) \rightarrow \mathbb{R}$ uniformly bounded in $C^{1, \frac{1}{4}}$, with $g_n(0) = g_n'(0) = 0$, such that $(\bar{S}_{w_n} - x_n) \cap C_{\nu_n, r_1} = \text{gr}_{\nu_n}(g_n)$, where ν_n is the normal to \bar{S}_{w_n} at x_n .

By compactness, $\nu_n \rightarrow \bar{\nu}$ (up to subsequences). For n large enough $C_{\bar{\nu}, r_1/2} \subset C_{\nu_n, r_1} + x_n - x_0$, and there exist functions f_n uniformly bounded in $C^{1, \frac{1}{4}}$ such that $\text{gr}_{\bar{\nu}}(f_n) \cap C_{\bar{\nu}, r_1/2} = (\text{gr}_{\nu_n}(g_n) + x_n - x_0) \cap C_{\bar{\nu}, r_1/2}$. Hence

$$(\bar{S}_{w_n} - x_0) \cap C_{\bar{\nu}, r_1/2} = \text{gr}_{\bar{\nu}}(f_n),$$

and by Ascoli-Arzelà Theorem f_n converges to some function f in $C^{1, \beta}$ for every $\beta < \frac{1}{4}$, with $f(0) = f'(0) = 0$. Using the Kuratowski convergence of \bar{S}_{w_n} to Γ , we deduce that $(\Gamma - x_0) \cap C_{\bar{\nu}, r_1/2} = \text{gr}_{\bar{\nu}}(f)$, and since $f'(0) = 0$ it must be $\bar{\nu} = \nu_\Gamma(x_0)$. \square

From what we have proved it follows that for every $n \in \mathbb{N}$ there exists a diffeomorphism $\Phi_n : \bar{\Omega} \rightarrow \bar{\Omega}$, with $\text{supp}(\Phi_n - Id) \subset\subset (U \setminus \partial_D \Omega)$, such that $\bar{S}_{w_n} = \Phi_n(\Gamma)$ and $\|\Phi_n - Id\|_{C^{1, \alpha}(\Gamma)} \rightarrow 0$.

With this information, we can finally conclude the proof of the isolated local minimality of u . Indeed, since $\mathcal{H}^1(\bar{S}_{w_n} \setminus S_{w_n}) = 0$ by Corollary 7.5, we have that $(\Phi_n(\Gamma), w_n) \in \mathcal{A}(\Omega)$ and $F(\Phi_n(\Gamma), w_n) = \mathcal{F}(w_n)$. Hence for n large enough, using (7.3),

$$F(\Phi_n(\Gamma), w_n) = \mathcal{F}(w_n) \leq \mathcal{F}(u) = F(\Gamma, u),$$

which implies that $\Phi_n = Id$ and $w_n = u$ for all (large) n by Proposition 6.2. Hence u itself is a solution to (7.2), and as a consequence of (7.1) also v_n solves the same minimum problem. We can then repeat all the previous argument for the sequence v_n instead of w_n , which leads, as before, to $v_n = u$ for n sufficiently large. This is the desired contradiction, since we are assuming $v_n \neq u$ for every n .

8. APPLICATIONS AND EXAMPLES

We start this section by showing that any regular critical pair (Γ, u) satisfying (4.18) is strictly stable in a sufficiently small tubular neighborhood $\mathcal{N}_\varepsilon(\Gamma)$ of the discontinuity set. As a consequence of our main result, we deduce the local minimality of (Γ, u) in $\mathcal{N}_\varepsilon(\Gamma)$, and also that (Γ, u) is in fact a global minimizer in a smaller neighborhood. This is in analogy with the result proved in [19], where it is shown, by means of a calibration method, that a critical point is a Dirichlet minimizer in small domains.

Proposition 8.1 (local and global minimality in small tubular neighborhoods). *Let (Γ, u) be a regular critical pair satisfying condition (4.18). Then there exists $\varepsilon_0 > 0$ such that the tubular neighborhood $\mathcal{N}_\varepsilon(\Gamma)$ of Γ is an admissible subdomain and (Γ, u) is strictly stable in $\mathcal{N}_\varepsilon(\Gamma)$ for every $\varepsilon < \varepsilon_0$. In particular, there exists $\delta > 0$ such that $F(\Gamma, u) < F(K, v)$ for every $(K, v) \in \mathcal{A}(\Omega)$ with $0 < \|u - v\|_{L^1(\Omega)} < \delta$ and $v = u$ in $\Omega \setminus \mathcal{N}_\varepsilon(\Gamma)$.*

Moreover, there exists $\varepsilon_1 \in (0, \varepsilon_0)$ such that (Γ, u) is a global minimizer in $\mathcal{N}_\varepsilon(\Gamma)$ for every $\varepsilon < \varepsilon_1$, in the sense that $F(\Gamma, u) \leq F(K, v)$ for every $(K, v) \in \mathcal{A}(\Omega)$ with $v = u$ in $\Omega \setminus \mathcal{N}_\varepsilon(\Gamma)$.

Proof. Clearly $\mathcal{N}_\varepsilon(\Gamma)$ is an admissible subdomain for ε small enough, and in view of Proposition 4.11 we shall prove that

$$\lim_{\varepsilon \rightarrow 0} \mu(\mathcal{N}_\varepsilon(\Gamma)) = +\infty$$

in order to obtain the first part of the statement. Assume by contradiction that there exist $\varepsilon_n \rightarrow 0^+$, $C > 0$ and $v_n \in H_{U_n}^1(\Omega \setminus \Gamma)$ such that $\|\Phi_{v_n}\|_\sim = 1$ and

$$2 \int_{\Omega} |\nabla v_n|^2 \leq C$$

for every n , where we set $U_n := \mathcal{N}_{\varepsilon_n}(\Gamma)$. Then v_n is a bounded sequence in $H_{U_1}^1(\Omega \setminus \Gamma)$, which converges weakly to 0 since the measure of U_n goes to 0. By compactness of the map $v \mapsto \Phi_v$, we have that Φ_{v_n} converge to 0 strongly in $H^1(\Gamma \cap \Omega)$, which is in contradiction to the fact that $\|\Phi_{v_n}\|_\sim = 1$ for every n .

To prove the second part of the statement, let u_ε be a solution to the minimum problem

$$\min\{\mathcal{F}(v) : v \in SBV(\Omega), v = u \text{ in } \Omega \setminus \mathcal{N}_\varepsilon(\Gamma)\}, \quad (8.1)$$

where \mathcal{F} is the relaxed functional introduced at the beginning of Section 7. We remark that, by classical regularity results for minimizers of the Mumford-Shah functional, $\mathcal{H}^1(\overline{S_{u_\varepsilon}} \setminus S_{u_\varepsilon}) = 0$ and thus $\mathcal{F}(u_\varepsilon) = F(\overline{S_{u_\varepsilon}}, u_\varepsilon)$. Hence, since $u_\varepsilon \rightarrow u$ in $L^1(\Omega)$ as $\varepsilon \rightarrow 0$ because the measure of $\mathcal{N}_\varepsilon(\Gamma)$ goes to 0, we conclude that $u_\varepsilon = u$ for ε small enough, as a consequence of the isolated local minimality of (Γ, u) . Then u is a solution to (8.1), and the conclusion follows by Remark 7.1. \square

Remark 8.2. Let (Γ, u) be a regular critical pair, and assume that

$$-2 \int_{\Omega} |\nabla v_\varphi|^2 dx + \int_{\Gamma \cap \Omega} |\nabla_\Gamma \varphi|^2 d\mathcal{H}^1 + \int_{\Gamma \cap \Omega} H^2 \varphi^2 d\mathcal{H}^1 - \int_{\Gamma \cap \partial\Omega} H_{\partial\Omega} \varphi^2 d\mathcal{H}^0 > 0$$

for every $\varphi \in H^1(\Gamma \cap \Omega) \setminus \{0\}$, where $v_\varphi \in H^1(\Omega \setminus \Gamma)$, $v_\varphi = 0$ on $\partial_D \Omega$, solves

$$\int_{\Omega} \nabla v_\varphi \cdot \nabla z dx + \int_{\Gamma \cap \Omega} [z^+ \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^+) - z^- \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^-)] d\mathcal{H}^1 = 0$$

for every $z \in H^1(\Omega \setminus \Gamma)$ with $z = 0$ on $\partial_D \Omega$. Then (Γ, u) is strictly stable in every admissible subdomain U . Hence, under the previous assumptions we can conclude that for every neighborhood $\mathcal{N}_\eta(\mathcal{S})$, where \mathcal{S} is the relative boundary of $\partial_D \Omega$ in $\partial\Omega$, there exists $\delta(\eta) > 0$ such that $F(\Gamma, u) < F(K, v)$ for every $(K, v) \in \mathcal{A}(\Omega)$ with $\|v - u\|_{L^1(\Omega)} < \delta$ and $v = u$ in $\mathcal{N}_\eta(\mathcal{S})$.

We now provide some explicit examples of critical point to which Theorem 3.7 can be applied. In particular, in Example 8.3 we discuss how the stability of constant critical pairs depends on the geometry of the domain Ω , while in Remark 8.4 we discuss how to construct families of (non-constant) critical pairs by a perturbing the Dirichlet data.

Example 8.3. Let Γ be a straight line contained in Ω connecting two points $x_1, x_2 \in \partial\Omega$ of minimal distance, and let u be equal to two different constants in the two connected components of $\Omega \setminus \Gamma$. Assume that Ω is strictly concave at x_1 and x_2 (that is, the curvature $H_{\partial\Omega}$ with respect to the exterior normal is strictly negative at x_1 and x_2). Then (Γ, u) is a regular critical pair such that for every admissible subdomain U

$$\partial^2 F((\Gamma, u); U)[\varphi] = \int_{\Gamma} |\nabla_\Gamma \varphi|^2 - H_{\partial\Omega}(x_1) \varphi^2(x_1) - H_{\partial\Omega}(x_2) \varphi^2(x_2) > 0$$

for every $\varphi \in H^1(\Gamma) \setminus \{0\}$. Hence it follows by Theorem 3.7 that (Γ, u) is an isolated local minimizer for F in every admissible subdomain U .

If the domain Ω is strictly convex, then a straight line connecting two points on $\partial\Omega$ of minimal distance is never a local minimizer: indeed, if U is any admissible subdomain, by evaluating the quadratic form $\partial^2 F((\Gamma, u); U)$ at the constant function $\varphi = 1$ we get

$$\partial^2 F((\Gamma, u); U)[1] = -H_{\partial\Omega}(x_1) - H_{\partial\Omega}(x_2) < 0.$$

We remark that this is not in contradiction to the result of Proposition 8.1, since in the present situation condition (4.18) is not satisfied.

Remark 8.4 (families of stable critical pairs by perturbation of the Dirichlet data). Let (Γ, u) be a strictly stable regular critical pair in an admissible subdomain U , and assume in addition that u^+ and u^- are of class C^2 in a neighborhood of Γ .

We fix a function $\psi_0 \in C_c^\infty(\partial_D \Omega)$ and we consider the perturbed Dirichlet datum $u_\varepsilon := u + \varepsilon \psi_0$ for $\varepsilon > 0$. As an application of the Implicit Function Theorem, one can show that for every ε sufficiently small there exists a strictly stable regular critical pair $(\Gamma_\varepsilon, v_\varepsilon)$ with $v_\varepsilon = u_\varepsilon$ in $(\Omega \setminus U) \cup \partial_D \Omega$.

The idea of the proof is to associate, with every $\varepsilon > 0$ and $\psi \in C^{2,\alpha}(\Gamma)$, the curve Γ_ψ defined as in (5.2) and the function $u_{\varepsilon,\psi}$ which minimizes the Dirichlet integral in $H^1(\Omega \setminus \Gamma_\psi)$ and attains the boundary condition $u_{\varepsilon,\psi} = u_\varepsilon$ in $(\Omega \setminus U) \cup \partial_D \Omega$. Then one considers the map

$$G : \mathbb{R} \times C^{2,\alpha}(\Gamma) \rightarrow C^{0,\alpha}(\Gamma) \times \mathbb{R} \times \mathbb{R}$$

defined by

$$G(\varepsilon, \psi) := \left((H_\psi - |\nabla_{\Gamma_\psi} u_{\varepsilon,\psi}^+|^2 + |\nabla_{\Gamma_\psi} u_{\varepsilon,\psi}^-|^2) \circ \Phi^\psi, (\eta_\psi \cdot \tau_{\partial\Omega})(\Phi^\psi(x_1)), (\eta_\psi \cdot \tau_{\partial\Omega})(\Phi^\psi(x_2)) \right).$$

Here Φ^ψ is a diffeomorphism mapping Γ onto Γ_ψ (see (5.2)), H_ψ denotes the curvature of Γ_ψ , η_ψ is the unit co-normal of $\Gamma_\psi \cap \partial\Omega$, $\{x_1, x_2\} = \Gamma \cap \partial\Omega$, and $\tau_{\partial\Omega}$ is the tangent vector to $\partial\Omega$, oriented in such a way that it coincides with ν in a neighborhood of $\Gamma \cap \partial\Omega$. It can be shown that the map G is of class C^1 in a neighborhood of $(0, 0)$, satisfies $G(0, 0) = 0$ (as (Γ, u) is a critical pair), and the partial derivative $\partial_\psi G(0, 0)$ is an invertible bounded linear operator, thanks to the strict positivity of the second variation at (Γ, u) (see [4] for more details). Hence it is possible to apply the Implicit Function Theorem and to obtain the desired family of critical pairs.

We conclude this section by observing, in the following remark, that our analysis can be extended to the periodic case: more precisely, we assume that the domain is a rectangle, Γ is a curve joining two opposite points on the boundary, and the Neumann boundary conditions are replaced by periodicity conditions on the sides connected by Γ . The remaining pair of sides represents the Dirichlet part of the boundary. We also discuss an explicit example in this different setting. In the remaining part of this section, with a slight abuse of notation we denote the generic point of \mathbb{R}^2 by (x, y) .

Remark 8.5. Let $R := [0, b] \times (-a, a)$, where $a, b > 0$ are positive real numbers. We define the infinite strip $\tilde{R} := \mathbb{R} \times (-a, a)$, the Dirichlet boundary $\partial_D R := [0, b] \times \{-a, a\}$, and the class of admissible pairs

$$\mathcal{A}(R) := \left\{ (K, v) : K \subset \mathbb{R}^2 \text{ closed, } K + (b, 0) = K, v \in H_{\text{loc}}^1(\tilde{R} \setminus K) \cap H^1(R \setminus K), \right. \\ \left. v_x(x+b, y) = v_x(x, y) \text{ for every } (x, y) \in \tilde{R} \setminus K \right\}.$$

We denote by $H_{\text{per}}^1(R \setminus K)$ the class of functions $z \in H_{\text{loc}}^1(\tilde{R} \setminus K) \cap H^1(R \setminus K)$ such that the map $x \mapsto z(x, y)$ is b -periodic for every $y \in (-a, a)$. Finally we consider the functional

$$F(K, v) := \int_{R \setminus K} |\nabla v|^2 + \mathcal{H}^1(K \cap R) \quad \text{for } (K, v) \in \mathcal{A}(R).$$

Similarly to what we did in Section 3, we say that $(\Gamma, u) \in \mathcal{A}(R)$ is a regular critical pair if $\Gamma \subset \tilde{R}$ is a curve of class C^∞ such that $\Gamma \cap R$ connects two opposite points on the $\partial_D R$, u satisfies

$$\int_{R \setminus \Gamma} \nabla u \cdot \nabla z = 0 \quad \text{for every } z \in H_{\text{per}}^1(R \setminus \Gamma) \text{ with } z = 0 \text{ on } \partial_D R,$$

and moreover the transmission condition and the non-vanishing jump condition (see Definition 3.3) hold on Γ . Setting $H_{\text{per}}^1(\Gamma) := \{\varphi \in H_{\text{loc}}^1(\Gamma) : \varphi(x+b, y) = \varphi(x, y) \text{ for every } (x, y) \in \Gamma\}$, we say that a regular critical pair (Γ, u) is strictly stable if

$$\partial^2 F(\Gamma, u)[\varphi] := -2 \int_R |\nabla v_\varphi|^2 + \int_{\Gamma \cap R} |\nabla_\Gamma \varphi|^2 d\mathcal{H}^1 + \int_{\Gamma \cap R} H^2 \varphi^2 d\mathcal{H}^1 > 0$$

for every $\varphi \in H_{\text{per}}^1(\Gamma) \setminus \{0\}$, where $v_\varphi \in H_{\text{per}}^1(R \setminus \Gamma)$, $v_\varphi = 0$ on $\partial_D R$, is the solution to

$$\int_R \nabla v_\varphi \cdot \nabla z + \int_{\Gamma \cap R} \left[z^+ \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^+) - z^- \operatorname{div}_\Gamma(\varphi \nabla_\Gamma u^-) \right] d\mathcal{H}^1 = 0 \quad (8.2)$$

for every $z \in H_{\text{per}}^1(R \setminus \Gamma)$, $z = 0$ on $\partial_D R$.

Then one can prove that every strictly stable regular critical pair (Γ, u) is a local minimizer, in the sense that there exists $\delta > 0$ such that $F(\Gamma, u) < F(K, v)$ for every $(K, v) \in \mathcal{A}(R)$ with $v = u$ on $\partial_D R$ and $0 < \|u - v\|_{L^1(R)} < \delta$. We omit the proof of this result, since it can be obtained by repeating all the arguments which lead to the proof of Theorem 3.7 with the natural modifications (notice that the proof in the present setting is in fact simpler, since by periodicity we can work in the whole strip \tilde{R} avoiding the technical difficulties related to the presence of Neumann boundary conditions).

Example 8.6. Here we adapt to the periodic setting described in Remark 8.5 the example discussed in [6, Section 7]. Setting $R = [0, b) \times (-a, a)$, we consider the regular critical pair $(\Gamma, u) \in \mathcal{A}(R)$ where $\Gamma = \mathbb{R} \times \{0\}$ and $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the function

$$u(x, y) := \begin{cases} x + 1 & \text{for } y \geq 0, \\ -x & \text{for } y < 0. \end{cases}$$

Notice that the energy of (Γ, u) is invariant along vertical translations of the discontinuity set. Nevertheless, we shall prove in fact that if

$$\frac{2b}{\pi} \tanh\left(\frac{2\pi a}{b}\right) < 1, \quad (8.3)$$

then (Γ, u) is an isolated local minimizer up to vertical translations: precisely, there exists $\delta > 0$ such that $F(\Gamma, u) < F(K, v)$ for every $(K, v) \in \mathcal{A}(R)$ with $v = u$ on $\partial_D R$ and $\|u - v\|_{L^1(R)} < \delta$, unless K coincides with a vertical translation of Γ . Moreover, (8.3) is sharp in the sense that if $\frac{2b}{\pi} \tanh\left(\frac{2\pi a}{b}\right) > 1$ then (Γ, u) is unstable.

To this aim, we will test the strict positivity of second variation at (Γ, u) on the subspace $H_0^1(0, b)$ of $H_{\text{per}}^1(\Gamma)$ of the functions vanishing at the endpoints, showing that

$$\partial^2 F(\Gamma, u)[\varphi] \geq C_0 \|\varphi\|_{H^1(0, b)} \quad \text{for every } \varphi \in H_0^1(0, b) \setminus \{0\} \quad \text{iff} \quad \frac{2b}{\pi} \tanh\left(\frac{2\pi a}{b}\right) < 1. \quad (8.4)$$

In turn, setting $\Gamma_\varepsilon := \mathbb{R} \times \{\varepsilon\}$ and

$$u_\varepsilon(x, y) := \begin{cases} x + 1 & \text{for } y \geq \varepsilon, \\ -x & \text{for } y < \varepsilon, \end{cases}$$

we have that $(\Gamma_\varepsilon, u_\varepsilon)$ is still a critical pair with the same energy of (Γ, u) , and, assuming (8.3) and (8.4), there exists $\varepsilon_0 > 0$ such that for every $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$ we have

$$\partial^2 F(\Gamma_\varepsilon, u_\varepsilon)[\varphi] \geq \frac{C_0}{2} \|\varphi\|_{H^1(0, b)}^2 \quad \text{for every } \varphi \in H_0^1(0, b) \setminus \{0\}. \quad (8.5)$$

This can be deduced by comparing the explicit expressions of the second variation at (Γ, u) and at $(\Gamma_\varepsilon, u_\varepsilon)$ and observing that

$$\sup_{\|\varphi\|_{H^1(0, b)}=1} \left| \int_R |\nabla v_\varphi^\varepsilon|^2 - \int_R |\nabla v_\varphi|^2 \right| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

(where v_φ and v_φ^ε are the solutions to (8.2) corresponding to (Γ, u) and $(\Gamma_\varepsilon, u_\varepsilon)$ respectively); this last estimate is obtained by subtracting the equations satisfied by v_φ and v_φ^ε . From (8.5) it follows that any configuration which is close in $W^{2, \infty}$ and coincides with Γ_ε at the endpoints has strictly larger energy than $(\Gamma_\varepsilon, u_\varepsilon)$: more precisely, there exists $\delta_0 > 0$ such that for every $|\varepsilon| < \varepsilon_0$, for every b -periodic function $h \in W_{\text{loc}}^{2, \infty}(\mathbb{R})$ with $0 < \|h - \varepsilon\|_{W^{2, \infty}(0, b)} < \delta_0$, $h(0) = h(b) = \varepsilon$, and for every v such that $(\Gamma_h, v) \in \mathcal{A}(R)$ and $v = u$ on $\partial_D R$, we have $F(\Gamma_h, v) > F(\Gamma_\varepsilon, u_\varepsilon) = F(\Gamma, u)$, where we denoted by Γ_h the graph of h . This can be deduced by repeating the arguments for the proof of Theorem 5.2, paying attention to the fact that the local minimality neighborhood can be chosen uniform with respect to n .

In turn, from this property easily follows the isolated local $W^{2, \infty}$ -minimality of (Γ, u) , since it implies the existence of a positive δ such that for every $(\Gamma_h, v) \in \mathcal{A}(R)$ with $0 < \|h\|_{W^{2, \infty}(0, b)} < \delta$

and $v = u$ on $\partial_D R$ we have $F(\Gamma_h, v) > F(\Gamma, u)$, unless $\Gamma_h = \Gamma_\varepsilon$ for some $\varepsilon > 0$ and $v = u_\varepsilon$. Finally, this property implies also the local L^1 -minimality (up to translations), by the same argument developed in Sections 6 and 7.

We are left with the proof of (8.4). Condition (4.18) is automatically satisfied on the subspace $H_0^1(0, b)$, and we can discuss the sign of $\partial^2 F(\Gamma, u)$ in terms of the eigenvalue λ_1 introduced in (4.20). We will prove that

$$\lambda_1(R) = \frac{2b}{\pi} \tanh \frac{2\pi a}{b}. \quad (8.6)$$

We remark that λ_1 coincides with the greatest λ such that there exists a nontrivial solution $(v, \varphi) \in H_{\text{per}}^1(R \setminus \Gamma) \times \tilde{H}_0^1(0, b)$, $v = 0$ in $\partial_D R$, to the equations

$$\lambda \int_R \nabla v \cdot \nabla z + \int_0^b (\varphi' z^+ + \varphi' z^-) dx = 0, \quad \int_0^b (\varphi' \psi' + 2\psi' v^+ + 2\psi' v^-) dx = 0$$

for every $z \in H_{\text{per}}^1(R \setminus \Gamma)$ with $z = 0$ on $\partial_D R$, and for every $\psi \in \tilde{H}_0^1(0, b)$. By symmetry, $v(x, y) = v(x, -y)$, so that by setting $R^+ := (0, b) \times (0, a)$, we look for a solution to

$$\begin{cases} \Delta v = 0 & \text{in } R^+, \\ v = 0 & \text{on } \partial_D R, \\ \lambda \partial_y v = \varphi' & \text{on } \Gamma, \\ \varphi'' = -4\partial_x v & \text{on } \Gamma. \end{cases}$$

The last two conditions say that

$$\lambda \partial_y v(x, 0) = -4(v(x, 0) - c), \quad c := \frac{1}{b} \int_0^b v(x, 0) dx.$$

Hence we are left with the determination of the greatest λ such that there exists a nontrivial periodic solution v to the system

$$\begin{cases} \Delta v = 0 & \text{in } R^+, \\ v = 0 & \text{on } \partial_D R, \\ \lambda \partial_y v = -4(v - c) & \text{on } \Gamma. \end{cases}$$

We expand $v(\cdot, y)$ in series of cosines:

$$v(x, y) = \sum_{n=0}^{+\infty} c_n(y) \cos\left(\frac{n\pi}{b}x\right),$$

and by the first two condition of the system we have that $c_n(y) = c_n \sinh\left(\frac{n\pi}{b}(a-y)\right)$, with $c_n \in \mathbb{R}$. Hence

$$v(x, y) = \sum_{n=0}^{+\infty} c_n \cos\left(\frac{n\pi}{b}x\right) \sinh\left(\frac{n\pi}{b}(a-y)\right)$$

and by imposing the last condition of the system we have

$$\lambda \sum_{n=0}^{+\infty} c_n \frac{n\pi}{b} \cos\left(\frac{n\pi}{b}x\right) \cosh\left(\frac{n\pi}{b}a\right) = 4 \sum_{n=0}^{+\infty} c_n \cos\left(\frac{n\pi}{b}x\right) \sinh\left(\frac{n\pi}{b}a\right) - 4c.$$

By expanding also c in series of cosines, we deduce from the previous inequality that $c = 0$, and also

$$\lambda c_n \frac{n\pi}{b} \cosh\left(\frac{n\pi a}{b}\right) = 4c_n \sinh\left(\frac{n\pi a}{b}\right)$$

for all $n \geq 1$. Hence, since we are looking for a positive λ , it follows that $\lambda = \frac{4b}{n\pi} \tanh\left(\frac{n\pi a}{b}\right)$ whenever $c_n \neq 0$. Thus only one of the coefficients c_n can be different from 0, and by periodicity it must correspond to an even index (here we used also the fact that the function $t \mapsto \frac{4b}{t\pi} \tanh\left(\frac{t\pi a}{b}\right)$ is monotone decreasing). Hence there exists $\bar{n} \geq 2$ even such that $c_{\bar{n}} \neq 0$ and

$$\lambda = \frac{4b}{\bar{n}\pi} \tanh\left(\frac{\bar{n}\pi a}{b}\right),$$

and clearly the largest value of λ corresponds to $\bar{n} = 2$. This completes the proof of (8.6) and, in turn, of (8.4).

9. APPENDIX

We collect here some technical results which have been used in the paper. In the following lemma we assume to be in the same setting as described at the beginning of Section 5.

Lemma 9.1. *Let $(\Gamma_n)_n$ be a sequence of curves of class $C^{1,\alpha}$, for some $\alpha \in (0,1)$, converging to Γ in $C^{1,\alpha}$, in the sense that there exist diffeomorphisms $\Phi_n : \bar{\Omega} \rightarrow \bar{\Omega}$ of class $C^{1,\alpha}$ such that $\Gamma_n = \Phi_n(\Gamma)$ and $\|\Phi_n - Id\|_{C^{1,\alpha}(\Gamma)} \rightarrow 0$.*

Then there exist $\psi_n \in C^{1,\alpha}(\Gamma)$, with $\psi_n \rightarrow 0$ in $C^{1,\alpha}(\Gamma)$, such that $\Gamma_n = \Gamma_{\psi_n}$, where Γ_{ψ_n} is the set defined according to (5.2).

Moreover, denoting by H_{Γ_n} and H the curvatures of Γ_n and of Γ respectively, if

$$\|H_{\Gamma_n} \circ \Phi_n - H\|_{L^\infty(\Gamma)} \rightarrow 0 \quad (9.1)$$

then ψ_n is of class $W^{2,\infty}$ and $\psi_n \rightarrow 0$ in $W^{2,\infty}(\Gamma)$.

Proof. We first extend each curve Γ_n (and Γ itself) outside $\bar{\Omega}$ as a straight line so that the resulting curves are of class $C^{1,\alpha}$ and still converge to Γ in the $C^{1,\alpha}$ sense. We can then localize in a small square $R = (-\rho, \rho) \times (-\rho, \rho)$ (which we assume for simplicity centered at the origin) in which we can express Γ and Γ_n as graphs of $C^{1,\alpha}$ functions:

$$\Gamma_n \cap R = \{(x, f_n(x)) : x \in (-\rho, \rho)\}, \quad \Gamma \cap R = \{(x, f(x)) : x \in (-\rho, \rho)\}$$

with $f_n \rightarrow f$ in $C^{1,\alpha}$. By a covering argument it is sufficient to prove the result in R (notice that, by our extension of the curves outside $\bar{\Omega}$, in this way we can cover also a neighborhood of the intersection of Γ with $\partial\Omega$).

We recall that in a sufficiently small tubular neighborhood $\mathcal{N}_{\eta_0}(\Gamma)$ of Γ are well defined two maps $\pi : \mathcal{N}_{\eta_0}(\Gamma) \rightarrow \Gamma$, $\tau : \mathcal{N}_{\eta_0}(\Gamma) \rightarrow \mathbb{R}$ of class C^2 (thank to the C^2 regularity of the vector field X generating the flow Ψ) such that $y = \Psi(\tau(y), \pi(y))$ for every y .

Taking $\rho' < \rho$, for n sufficiently large we can define a map $\tilde{\pi}_n : (-\rho', \rho') \rightarrow (-\rho, \rho)$ by setting $\tilde{\pi}_n(x) := \pi_1 \circ \pi(x, f_n(x))$, where $\pi_1(x, y) := x$. Notice that $\tilde{\pi}_n$ tends to the identity in $C^{1,\alpha}$, hence it is invertible and also its inverse converges to the identity in $C^{1,\alpha}$. Defining

$$\phi_n(x) := \tau(\tilde{\pi}_n^{-1}(x), f_n(\tilde{\pi}_n^{-1}(x)))$$

for $x \in (-\rho', \rho')$, since τ is regular and vanishes on Γ we deduce that $\phi_n \rightarrow 0$ in $C^{1,\alpha}(-\rho', \rho')$.

Hence the map $\psi_n(x, f(x)) := \phi_n(x)$, for $|x| < \rho'$, is of class $C^{1,\alpha}$ on $\Gamma \cap ((-\rho', \rho') \times (-\rho, \rho))$, converges to 0 in $C^{1,\alpha}$ and satisfies $\Gamma_{\psi_n} = \Gamma_n$. This proves the first part of the statement.

The second part follows similarly: indeed, since the sets Γ_n are locally one-dimensional graphs, the boundedness in L^∞ of the curvatures of Γ_n yields the $W^{2,\infty}$ -regularity of the functions f_n , and the convergence (9.1) implies in addition that $f_n \rightarrow f$ in $W^{2,\infty}$. Hence the conclusion follows from the explicit expression of ψ_n obtained above. \square

We conclude with two regularity results for the Neumann problem and for the mixed Dirichlet-Neumann problem in planar domains with angles.

Lemma 9.2. *Let A be an open subset of the unit ball B_1 such that $\partial A \cap B_1 = \Gamma_1 \cup \Gamma_2$, where Γ_1 and Γ_2 are two curves of class $C^{1,\beta}$ meeting at the origin with an internal angle $\alpha \in (0, \pi)$. Let $u \in H^1(A)$ be a weak solution to*

$$\begin{cases} \Delta u = 0 & \text{in } A, \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \cup \Gamma_2. \end{cases}$$

Then ∇u has a $C^{0,\gamma}$ extension up to $\Gamma_1 \cup \Gamma_2$, for $\gamma = \min\{\beta, \frac{\pi}{\alpha} - 1\}$, with $C^{0,\gamma}$ -norm bounded by a constant depending only on the $C^{1,\beta}$ -norm of Γ_1 and Γ_2 .

Proof. We consider A as a subset of the complex plane \mathbb{C} (we can assume without loss of generality that the positive real axis coincides with the tangent to Γ_1 at the origin, and that the tangent to Γ_2 at the origin is the line $\{z = \rho e^{i\theta} : \rho > 0, \theta = \alpha\}$). Consider the map $\Phi : \bar{A} \rightarrow \Phi(\bar{A})$ given by $\Phi(z) := z^{\frac{\pi}{\alpha}} = \rho^{\frac{\pi}{\alpha}} e^{i\frac{\pi}{\alpha}\theta}$, where $z = \rho e^{i\theta}$. The map Φ is of class $C^{1, \frac{\pi}{\alpha}-1}(\bar{A})$, and since it is conformal out of the origin, the function $v := u \circ \Phi^{-1}$ is harmonic in $\Phi(A)$ and satisfies a

homogeneous Neumann condition on $\Phi(\Gamma_1 \cup \Gamma_2)$. Moreover $\Phi(\Gamma_1 \cup \Gamma_2)$ is a curve of class $C^{1,\gamma}$, hence by classical regularity results (see, e.g., [2, Theorem 7.49]) ∇v has a $C^{0,\gamma}$ extension up to $\Phi(\Gamma_1 \cup \Gamma_2)$, with $C^{0,\gamma}$ -norm bounded by a constant depending only on the $C^{1,\gamma}$ -norm of $\Phi(\Gamma_2)$. The conclusion follows since $u = v \circ \Phi$, using the regularity of Φ . \square

Lemma 9.3. *Let A be an open subset of the unit ball B_1 such that $\partial A \cap B_1 = \Gamma_1 \cup \Gamma_2$, where Γ_1 and Γ_2 are two curves of class $C^{1,\beta}$ meeting at the origin with an internal angle equal to $\frac{\pi}{2}$. Let $u \in H^1(A)$ be a weak solution to*

$$\begin{cases} \Delta u = f & \text{in } A \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \\ u = u_0 & \text{on } \Gamma_2 \end{cases} \quad \text{or to} \quad \begin{cases} \Delta u = f & \text{in } A \\ \partial_\nu u = 0 & \text{on } \Gamma_1 \cup \Gamma_2 \end{cases}$$

where $f \in L^\infty(A)$, and $u_0 \in C^2(\bar{A})$ is such that $\partial_\nu u_0 = 0$ on Γ_1 . Then ∇u has a $C^{0,\beta}$ extension up to $\Gamma_1 \cup \Gamma_2$, with $C^{0,\beta}$ -norm bounded by a constant depending only on $\|f\|_\infty$, on the $C^{1,\beta}$ -norm of Γ_1 and Γ_2 , and on $\|u_0\|_{C^2(\bar{A})}$ in the first case.

Proof. Let u solve the first problem, and let $\tilde{u} := u - u_0$. Then \tilde{u} is a solution to

$$\begin{cases} \Delta \tilde{u} = \tilde{f} & \text{in } A, \\ \partial_\nu \tilde{u} = 0 & \text{on } \Gamma_1, \\ \tilde{u} = 0 & \text{on } \Gamma_2, \end{cases}$$

where $\tilde{f} := f - \Delta u_0$. We can find a radius $\rho > 0$ and a $C^{1,\beta}$ conformal mapping Φ in $\bar{A} \cap \bar{B}_\rho$ such that $\Phi(\Gamma_1)$ is a straight line meeting $\Phi(\Gamma_2)$ orthogonally. Then the function $v := \tilde{u} \circ \Phi^{-1}$ solves

$$\begin{cases} \Delta v = g & \text{in } \Phi(A), \\ \partial_\nu v = 0 & \text{on } \Phi(\Gamma_1), \\ v = 0 & \text{on } \Phi(\Gamma_2), \end{cases}$$

where $g := (\tilde{f} \circ \Phi^{-1})|\det \nabla \Phi^{-1}|$. By even reflection across $\Phi(\Gamma_1)$ and by applying classical regularity results, we can conclude that ∇v has a $C^{0,\beta}$ extension up to $\Phi(\Gamma_1 \cup \Gamma_2)$, with $C^{0,\beta}$ -norm bounded by a constant depending only on $\|g\|_\infty$ and on the $C^{1,\beta}$ -norm of $\Phi(\Gamma_1 \cup \Gamma_2)$. Now the conclusion follows by using the regularity of the map Φ .

The regularity for the solution to the second problem can be obtained by a similar (and, in fact, simpler) argument. \square

9.1. Proof of the density lower bound. This concluding subsection is entirely devoted to the proof of Theorem 2.4. Most of the proofs are classical and very similar to those contained in [3] (which in turn follow the approach of [11]), and for this reason we will just sketch them by describing only the main changes needed, referring the interested reader to [4] for details.

We start by observing that, if w satisfies the hypotheses of the theorem, the following energy upper bound holds in every ball $B_\rho(x)$ with $\rho \leq R_0$ (it can be easily deduced by comparing the energies of w and of $w\chi_{\Omega' \setminus (B_\rho(x) \cap \Omega)}$):

$$\mathcal{F}(w; B_\rho(x) \cap \Omega') \leq c_0 \rho, \tag{9.2}$$

where c_0 depends only on R_0 , ω , u and Ω . In the following, C will always denote a positive constant depending only on the previous quantities. We now show that we can replace the Dirichlet condition in $\Omega' \setminus \Omega$ by a homogeneous boundary condition.

Lemma 9.4. *Set $\tilde{w} := w - u$. Then $\tilde{w} \in SBV(\Omega')$, $\tilde{w} = 0$ in $\Omega' \setminus \Omega$, and there exist $\eta > 0$, $\tilde{\omega} > 0$ (depending only on Ω , ω and u) such that for every $x \in \bar{\Omega} \cap \mathcal{N}_\eta(\partial_D \Omega)$ and for every $\rho < \eta$*

$$\mathcal{F}(\tilde{w}; B_\rho(x) \cap \Omega') \leq \mathcal{F}(v; B_\rho(x) \cap \Omega') + \tilde{\omega} \rho^{\frac{3}{2}}$$

whenever $v \in SBV(\Omega')$ is such that $v = 0$ in $\Omega' \setminus \Omega$ and $\{v \neq \tilde{w}\} \subset\subset B_\rho(x)$.

Proof. By choosing η sufficiently small, we can guarantee that $\bar{S}_u \cap B_\rho(x) = \emptyset$ for each ball $B_\rho(x)$ as in the statement, hence $S_{\tilde{w}} \cap B_\rho(x) = S_w \cap B_\rho(x)$. By comparing the energies of w and $v + u$ we obtain

$$\mathcal{F}(\tilde{w}; B_\rho(x) \cap \Omega') \leq \mathcal{F}(v; B_\rho(x) \cap \Omega') + 2 \int_{B_\rho(x) \cap \Omega'} \nabla u \cdot (\nabla v - \nabla w) + 2 \int_{B_\rho(x) \cap \Omega'} |\nabla u|^2 + \omega \rho^2.$$

Now, using the fact that $\nabla u \in L^\infty$ and the upper bound (9.2), we have

$$2 \int_{B_\rho(x) \cap \Omega'} |\nabla u|^2 \leq C\rho^2, \quad -2 \int_{B_\rho(x) \cap \Omega'} \nabla w \cdot \nabla u \leq C\rho^{\frac{3}{2}},$$

while for every $\varepsilon > 0$ we have

$$2 \int_{B_\rho(x) \cap \Omega'} \nabla v \cdot \nabla u \leq \varepsilon^2 \|\nabla v\|_{L^2}^2 + \frac{1}{\varepsilon^2} \|\nabla u\|_{L^2}^2 \leq \varepsilon^2 \mathcal{F}(v; B_\rho(x) \cap \Omega') + \frac{C}{\varepsilon^2} \rho^2.$$

It follows that

$$\mathcal{F}(\tilde{w}; B_\rho(x) \cap \Omega') \leq (1 + \varepsilon^2) \mathcal{F}(v; B_\rho(x) \cap \Omega') + C \left(1 + \frac{1}{\varepsilon^2}\right) \rho^2 + C\rho^{\frac{3}{2}}.$$

Defining the deviation from minimality of \tilde{w} in a Borel set B as

$$\text{Dev}(\tilde{w}; B) := \mathcal{F}(\tilde{w}; B \cap \Omega') - \inf \{ \mathcal{F}(v; B \cap \Omega') : v \in SBV(\Omega'), v = 0 \text{ in } \Omega' \setminus \Omega, \{v \neq \tilde{w}\} \subset\subset B \}, \quad (9.3)$$

from the previous inequality we obtain, by taking the infimum over all v ,

$$\begin{aligned} \text{Dev}(\tilde{w}; B_\rho(x)) &\leq \varepsilon^2 \mathcal{F}(\tilde{w}; B_\rho(x) \cap \Omega') + C \left(1 + \frac{1}{\varepsilon^2}\right) \rho^2 + C\rho^{\frac{3}{2}} \\ &\leq c_0 \varepsilon^2 \rho + C \left(1 + \frac{1}{\varepsilon^2}\right) \rho^2 + C\rho^{\frac{3}{2}} \leq \tilde{\omega} \rho^{\frac{3}{2}}, \end{aligned}$$

where we used (9.2) in the second inequality and we choose $\varepsilon = \rho^{\frac{1}{4}}$ in the last inequality. \square

In the proof of the main decay property in Lemma 9.8 we will need to consider the blow-up in a sequence of balls whose centers converge to a point in $\overline{\partial_D \Omega} \cap \overline{\partial_N \Omega}$. This situation is examined in the following lemma.

Lemma 9.5. *Let $x_n \in \bar{\Omega}$, $x_n \rightarrow x_0 \in \overline{\partial_D \Omega} \cap \overline{\partial_N \Omega}$, and $r_n \rightarrow 0^+$. Setting*

$$\Omega_n := \frac{\Omega' - x_n}{r_n} \cap B_1, \quad D_n := \frac{(\Omega' \setminus \Omega) - x_n}{r_n} \cap B_1, \quad (9.4)$$

there exist $\delta_1, \delta_2 \in [0, 1]$ and a coordinate system such that (up to subsequences)

$$\Omega_n \rightarrow \Omega_0 := \{(\xi, \zeta) \in B_1 : \xi < \delta_1\}, \quad D_n \rightarrow D_0 := \{(\xi, \zeta) \in B_1 : \xi < \delta_1, \zeta > \delta_2\}$$

in L^1 . Moreover, the relative isoperimetric inequality holds in Ω_n with a constant which can be chosen independently of n (and which will be denoted by γ). Finally, assuming $\delta_2 < 1$, given $v \in W^{1,2}(\Omega_0)$ with $v = 0$ in D_0 there exists a sequence $v_n \in W^{1,2}(B_1)$ such that $v_n \rightarrow v$ in $W^{1,2}(\Omega_0)$ and $v_n = 0$ in D_n .

Proof. The first part of the lemma states an intuitive geometric fact that can be proved rigorously arguing as in [3, Lemma 6.4]. The fact that the constant in the relative isoperimetric inequality can be chosen uniformly for all the sets Ω_n follows from the fact that the boundaries of the sets Ω_n are close to the boundary of Ω_0 in the C^1 sense.

Finally, we prove the last part of the statement, under the assumption $\delta_2 < 1$. We extend v to the set $\tilde{\Omega} = \Omega_0 \cup \{\zeta > \delta_2\}$ by setting $v = 0$ outside Ω_0 , and since $\tilde{\Omega}$ satisfies the exterior cone condition we can find $\tilde{v} \in W^{1,2}(\mathbb{R}^2)$ such that $\tilde{v}|_{\tilde{\Omega}} = v$. Setting, for $(\xi, \zeta) \in B_1$,

$$v_n(\xi, \zeta) := \tilde{v}(\xi, \zeta + a_n), \quad a_n := \sup_{(\xi, \zeta) \in \partial D_n \cap \Omega_n} |\zeta - \delta_2| \rightarrow 0,$$

we obtain a sequence with the desired properties. \square

In the following compactness property, which is a consequence of the Poincaré inequality, we adapt [2, Proposition 7.5] to our context.

Lemma 9.6. *Let x_n and r_n be as in Lemma 9.5, and assume that $|D_n| \geq d_0 > 0$ for every n . Let $u_n \in SBV(\Omega_n)$, with $u_n = 0$ a.e. in D_n , be such that*

$$\sup_n \int_{\Omega_n} |\nabla u_n|^2 dx < \infty, \quad \lim_{n \rightarrow \infty} \mathcal{H}^1(S_{u_n}) = 0.$$

Setting $\bar{u}_n := (u_n \wedge \tau_n^+) \vee \tau_n^-$, where

$$\begin{aligned} \tau_n^+ &:= \inf\{t \in [-\infty, +\infty] : |\{u_n < t\}| \geq |\Omega_n| - (2\gamma\mathcal{H}^1(S_{u_n}))^2\}, \\ \tau_n^- &:= \inf\{t \in [-\infty, +\infty] : |\{u_n < t\}| \geq (2\gamma\mathcal{H}^1(S_{u_n}))^2\}, \end{aligned}$$

(here γ is the constant introduced in Lemma 9.5), one has that $\bar{u}_n = 0$ in D_n for n large, and (up to subsequences) $\bar{u}_n \rightarrow v \in W^{1,2}(\Omega_0)$ in $L^2_{\text{loc}}(\Omega_0)$, $u_n \rightarrow v$ a.e. in Ω_0 , and for every $\rho \leq 1$

$$\int_{\Omega_0 \cap B_\rho} |\nabla v|^2 dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega_n \cap B_\rho} |\nabla \bar{u}_n|^2 dx. \quad (9.5)$$

Proof. The proof can be obtained by repeating word by word the proof of [3, Lemma 6.1]. We have only to be careful about the fact that in our context also the domain Ω_n depends on n and is not fixed along the sequence. The essential remark here is that the isoperimetric inequality holds in the sets Ω_n with a constant which can be chosen independent of n , as observed in Lemma 9.5. \square

The following lemma is a variant of [2, Theorem 7.7]. For $B \subset \mathbb{R}^2$ Borel set and $c > 0$ we set

$$\mathcal{F}(v, c; B) := \int_B |\nabla v|^2 dx + c\mathcal{H}^1(S_v \cap B).$$

Lemma 9.7. *Let x_n and r_n be as in Lemma 9.5, and assume that $|D_n| \geq d_0 > 0$ for every n . Let $c_n > 0$, $u_n \in SBV(\Omega_n)$, with $u_n = 0$ in D_n , be such that*

$$\begin{aligned} \sup_n \mathcal{F}(u_n, c_n; \Omega_n) &< +\infty, \quad \lim_{n \rightarrow +\infty} \text{Dev}_{D_n}(u_n, c_n; B_1) = 0, \\ \lim_{n \rightarrow +\infty} \mathcal{H}^1(S_{u_n}) &= 0, \quad u_n \rightarrow v \in W^{1,2}(\Omega_0) \text{ a.e. in } \Omega_0, \end{aligned}$$

where

$$\begin{aligned} \text{Dev}_{D_n}(v, c; B) &:= \mathcal{F}(v, c; B \cap \Omega_n) \\ &- \inf\{\mathcal{F}(w, c; B \cap \Omega_n) : w \in SBV(\Omega_n), w = 0 \text{ in } D_n, \{w \neq v\} \subset\subset B\}. \end{aligned}$$

Then

$$\int_{\Omega_0} |\nabla v|^2 dx \leq \int_{\Omega_0} |\nabla w|^2 dx$$

for every $w \in W^{1,2}(\Omega_0)$ such that $w = 0$ in D and $\{v \neq w\} \subset\subset B_1$, and

$$\lim_{n \rightarrow +\infty} \mathcal{F}(u_n, c_n; \Omega_n \cap B_\rho) = \int_{\Omega_0 \cap B_\rho} |\nabla v|^2 dx \quad \text{for every } \rho \in (0, 1).$$

Proof. Also in this case the proof is analogous to the one of [3, Proposition 6.2], and uses the auxiliary result contained in Lemma 9.6 in order to overcome the technical difficulties due to the fact of working in a variable domain. We remark that in the proof it is essential to use the last part of Lemma 9.5, which allows us to approximate any competitor w for v , vanishing in the limit domain D , with functions w_n vanishing in D_n , for which the quasi-minimality of v_n can be exploited. Taking into account these observations, it is straightforward to check that the proof of [3, Proposition 6.2] yields the conclusion also in our case. \square

The following lemma contains the main decay property used to prove Theorem 2.4.

Lemma 9.8. *There exists a positive constant C such that for every $\tau \in (0, 1)$ there exist $\varepsilon(\tau) > 0$, $\theta(\tau) > 0$ and $r(\tau) > 0$ with the property that for every $x \in \bar{\Omega}$ and $\rho \leq r(\tau)$, whenever $v \in SBV(\Omega' \cap B_\rho(x))$ is such that $v = 0$ in $(\Omega' \setminus \Omega) \cap B_\rho(x)$,*

$$\mathcal{H}^1(S_v \cap B_\rho(x) \cap \Omega') < \varepsilon(\tau)\rho, \quad \text{Dev}(v; B_\rho(x)) < \theta(\tau)\mathcal{F}(v; B_\rho(x) \cap \Omega')$$

(the deviation from minimality is defined as in (9.3)) then

$$\mathcal{F}(v; B_{\tau\rho}(x) \cap \Omega') \leq C\tau^2 \mathcal{F}(v; B_\rho(x) \cap \Omega').$$

Proof. By choosing C large enough, we can assume without loss of generality that $\tau < \frac{1}{4}$. The proof is by a contradiction argument: let $\varepsilon_n \rightarrow 0$, $\theta_n \rightarrow 0$, $r_n \rightarrow 0$, $x_n \in \overline{\Omega}$, $v_n \in SBV(B_{r_n}(x_n) \cap \Omega')$, $v_n = 0$ in $(\Omega' \setminus \Omega) \cap B_{r_n}(x_n)$, be such that

$$\mathcal{H}^1(S_{v_n} \cap B_{r_n}(x_n) \cap \Omega') = \varepsilon_n r_n, \quad \text{Dev}(v_n; B_{r_n}(x_n)) = \theta_n \mathcal{F}(v_n; B_{r_n}(x_n) \cap \Omega'),$$

and

$$\mathcal{F}(v_n; B_{\tau r_n}(x_n) \cap \Omega') > C\tau^2 \mathcal{F}(v_n; B_{r_n}(x_n) \cap \Omega'),$$

where C will be chosen later. By a change of variables, we set

$$w_n(y) := r_n^{-\frac{1}{2}} c_n^{\frac{1}{2}} v_n(x_n + r_n y), \quad c_n := \frac{r_n}{\mathcal{F}(v_n; B_{r_n}(x_n) \cap \Omega')}.$$

We obtain a sequence $w_n \in SBV(\Omega_n)$ such that $\mathcal{F}(w_n, c_n; \Omega_n) = 1$, $\text{Dev}_{D_n}(w_n, c_n; B_1) = \theta_n$, $\mathcal{H}^1(S_{w_n} \cap \Omega_n) = \varepsilon_n$, and

$$\mathcal{F}(w_n, c_n; B_\tau \cap \Omega_n) > C\tau^2$$

(here Ω_n and D_n are defined as in (9.4)). Up to subsequences, $x_n \rightarrow x_0$, and we are in one of the following cases:

- $x_0 \in \Omega$: in this case the balls $B_{r_n}(x_n)$ are contained in Ω for n large, hence the boundary does not play any role and the contradiction follows from [2, Lemma 7.14];
- $x_0 \in \partial_D \Omega$: the balls $B_{r_n}(x_n)$ intersect only the Dirichlet part of the boundary for n large, and the contradiction follows from [3, Lemma 6.6];
- $x_0 \in \partial_N \Omega$: we have that $\Omega_n \rightarrow \Omega_0 = \{(\xi, \zeta) \in B_1 : \xi < \delta_1\}$ for some $\delta_1 \in [0, 1]$ (in a suitable coordinate system) and $D_n = \emptyset$ for n large enough. Adapting Lemma 9.6 and Lemma 9.7 to this situation (in which the Dirichlet condition does not play any role) we have that, up to further subsequences, $w_n - m_n \rightarrow w$ almost everywhere in Ω_0 , where m_n are medians of w_n in Ω_n and $w \in W^{1,2}(\Omega_0)$, with

$$\int_{\Omega_0} |\nabla w|^2 \leq \liminf_n \int_{\Omega_n} |\nabla w_n|^2 \leq 1.$$

In addition, w is harmonic in Ω_0 and satisfies a homogeneous Neumann condition on $\{(\xi, \zeta) : \xi = \delta_1\}$, and hence (by the decay properties of harmonic functions)

$$C\tau^2 \leq \lim_{n \rightarrow +\infty} \mathcal{F}(w_n, c_n; B_\tau \cap \Omega_n) = \int_{B_\tau \cap \Omega_0} |\nabla w|^2 \leq 8\tau^2 \int_{B_{\frac{1}{2}} \cap \Omega_0} |\nabla w|^2 \leq 8\tau^2$$

which is a contradiction if we take $C > 8$.

- $x_0 \in \partial_D \Omega \cap \partial_N \Omega$: in this case we are under the assumptions of Lemma 9.5. If $\delta_2 \in (\frac{1}{2}, 1]$, then $B_{\frac{1}{2}} \cap D_n = \emptyset$ for n large enough, and we can argue exactly as in the previous case, in the ball $B_{\frac{1}{2}}$. It remains only to deal with the case $\delta_2 \in [0, \frac{1}{2}]$.

To get a contradiction also in the case $\delta_2 \in [0, \frac{1}{2}]$, observe first that $|D_n| \geq d_0 > 0$. We can apply Lemma 9.6 and Lemma 9.7 to deduce that, up to subsequences, $w_n \rightarrow w_\infty \in W^{1,2}(\Omega_0)$ a.e. in Ω_0 , with $w_\infty = 0$ in D ,

$$\int_{\Omega_0} |\nabla w_\infty|^2 \leq \liminf_{n \rightarrow \infty} \int_{\Omega_n} |\nabla w_n|^2 \leq 1.$$

Moreover for every $w \in W^{1,2}(\Omega_0)$ such that $w = 0$ in D and $\{w \neq w_\infty\} \subset \subset B_1$

$$\int_{\Omega_0} |\nabla w_\infty|^2 \leq \int_{\Omega_0} |\nabla w|^2,$$

and

$$\mathcal{F}(w_n, c_n; B_r \cap \Omega_n) \rightarrow \int_{B_r \cap \Omega_0} |\nabla w_\infty|^2 \quad \text{for every } r \in (0, 1).$$

If \tilde{w}_∞ is the harmonic function in B_1 obtained by applying firstly an even reflection of w_∞ across $\{(\xi, \zeta) : \xi = \delta_1\}$, and then an odd reflection across $\{(\xi, \zeta) : \zeta = \delta_2\}$, we conclude, by using the decay properties of harmonic functions, that

$$\begin{aligned} C\tau^2 &\leq \lim_{n \rightarrow \infty} \mathcal{F}(w_n, c_n; B_\tau \cap \Omega_n) = \int_{B_\tau \cap \Omega_0} |\nabla w_\infty|^2 \leq \int_{B_\tau} |\nabla \tilde{w}_\infty|^2 \\ &\leq (2\tau)^2 \int_{B_{\frac{1}{2}}} |\nabla \tilde{w}_\infty|^2 \leq 4(2\tau)^2 \int_{B_{\frac{1}{2}} \cap \Omega_0} |\nabla w_\infty|^2 \leq 16\tau^2, \end{aligned}$$

and this is a contradiction if we chose $C > 16$. \square

We have now all the ingredients to conclude the proof of Theorem 2.4.

Proof of Theorem 2.4. Let η be given by Lemma 9.4. We first observe that the density lower bound holds in any ball $B_\rho(x)$ with $x \in \overline{\Omega} \setminus \mathcal{N}_\eta(\partial_D \Omega)$ and $\rho \leq \rho_0$ (for some $\rho_0 < \eta$ depending only on ω , u and Ω): indeed, in this case the Dirichlet boundary condition does not play any role, and the result is classical. It is then sufficient to prove the lower bound for the function \tilde{w} defined in Lemma 9.4 in balls $B_\rho(x)$ centered at points $x \in \overline{S_\omega} \cap \mathcal{N}_\eta(\partial_D \Omega)$, since in such balls $S_w \cap B_\rho(x) = S_{\tilde{w}} \cap B_\rho(x)$ if $\rho < \eta$.

In turn, in this case the conclusion follows by repeating exactly the proof of [3, Theorem 3.4] (with the particular values of the parameters $N = 2$, $s = \frac{1}{2}$ and $\beta = 1$), using the decay property proved in our context in Lemma 9.8. \square

REFERENCES

- [1] E. ACERBI, N. FUSCO, M. MORINI, *Minimality via second variation for a nonlocal isoperimetric problem*. Comm. Math. Phys. **322** (2013), 515-557.
- [2] L. AMBROSIO, N. FUSCO, D. PALLARA, *Functions of bounded variation and free discontinuity problems*. Oxford University Press, New York, 2000.
- [3] J.-F. BABADJIAN, A. GIACOMINI, *Existence of strong solutions for quasi-static evolution in brittle fracture*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) (to appear).
- [4] M. BONACINI, *Minimality and stability results for a class of free-discontinuity and nonlocal isoperimetric problems*. Ph.D. thesis (available online at <https://digitalibrary.sissa.it/handle/1963/7191>).
- [5] A. BRAIDES, *Approximation of free-discontinuity problems*. Lecture notes in Mathematics, 1694. Springer-Verlag, Berlin, 1998.
- [6] F. CAGNETTI, M.G. MORA, M. MORINI, *A second order minimality condition for the Mumford-Shah functional*. Calc. Var. Partial Differential Equations **33** (2008), 37-74.
- [7] M. CARRIERO, A. LEACI, *Existence theorem for a Dirichlet problem with free discontinuity set*. Nonlinear Anal. **15** (1990), 661-677.
- [8] M. CICALESE, G.P. LEONARDI, *A selection principle for the sharp quantitative isoperimetric inequality*. Arch. Ration. Mech. Anal. **206** (2012), 617-643.
- [9] G. DAL MASO, M.G. MORA, M. MORINI, *Local calibrations for minimizers of the Mumford-Shah functional with rectilinear discontinuity sets*. J. Math. Pures Appl. **79** (2000), 141-162.
- [10] G. DAVID, *Singular sets of minimizers for the Mumford-Shah functional*. Progress in Mathematics, 233. Birkhäuser Verlag, Basel, 2005.
- [11] E. DE GIORGI, M. CARRIERO, A. LEACI, *Existence theorem for a minimum problem with free discontinuity set*. Arch. Ration. Mech. Anal. **108** (1989), 195-218.
- [12] N. FUSCO, M. MORINI, *Equilibrium configurations of epitaxially strained elastic films: second order minimality conditions and qualitative properties of solutions*. Arch. Ration. Mech. Anal. **203** (2012), 247-327.
- [13] P. GRISVARD, *Elliptic problems in nonsmooth domains*. Monographs and Studies in Mathematics, 24. Pitman (Advanced Publishing Program), Boston, 1985.
- [14] M. GRÜTER, *Boundary regularity for solutions of a partitioning problem*. Arch. Rational Mech. Anal. **97** (1987), 261-270.
- [15] V. JULIN, G. PISANTE, *Minimality via second variation for microphase separation of diblock copolymer melts*. Preprint (2013).
- [16] H. KOCH, G. LEONI, M. MORINI, *On optimal regularity of free boundary problems and a conjecture of De Giorgi*. Comm. Pure Appl. Math. **58** (2005), 1051-1076.
- [17] R.V. KOHN, P. STERNBERG, *Local minimisers and singular perturbations*. Proc. Roy. Soc. Edinburgh Sect. A **111** (1989), 69-84.
- [18] F. MADDALENA, S. SOLIMINI, *Blow-up techniques and regularity near the boundary for free discontinuity problems*. Adv. Nonlinear Stud. **1** (2001), 1-41.

- [19] M.G. MORA, M. MORINI, *Local calibrations for minimizers of the Mumford-Shah functional with a regular discontinuity set*. Ann. Inst. H. Poincaré Anal. Nonlin. **18** (2001), 403-436.
- [20] M. MORINI, *Free discontinuity problems: calibration and approximation of solutions*. Ph.D. Thesis (available online at <https://digitallibrary.sissa.it/handle/1963/5398>).
- [21] M. MORINI, *Global calibrations for the non-homogeneous Mumford-Shah functional*. Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **1** (2002), 603-648.
- [22] D. MUMFORD, J. SHAH, *Boundary detection by minimizing functionals, I*. Proc. IEEE Conf. on Computer Vision and Pattern Recognition (San Francisco, 1985).
- [23] D. MUMFORD, J. SHAH, *Optimal approximation by piecewise smooth functions and associated variational problems*. Comm. Pure Appl. Math **17** (1989), 577-685.
- [24] L. SIMON, *Lectures on geometric measure theory*. Proceedings of the Centre for Mathematical Analysis, Australian National University, 3. Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [25] P. STERNBERG, K. ZUMBRUN, *A Poincaré inequality with applications to volume-constrained area-minimizing surfaces*. J. Reine Angew. Math. **503** (1998), 63-85.

(M. Bonacini) SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY
Email address: marco.bonacini@sissa.it

(M. Morini) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI PARMA, PARMA, ITALY
Email address: massimiliano.morini@unipr.it