A NOTE ON CODAZZI TENSORS

GIOVANNI CATINO, CARLO MANTEGAZZA, AND LORENZO MAZZIERI

ABSTRACT. We discuss a gap in Besse's book [1], recently pointed out by Merton in [7], which concerns the classification of Riemannian manifolds admitting a Codazzi tensors with exactly two distinct eigenvalues. For such manifolds, we prove a structure theorem, without adding extra hypotheses and then we conclude with some application of this theory to the classification of three–dimensional gradient Ricci solitons.

CONTENTS

1.	Introduction	1
2.	Codazzi Tensors with Two Distinct Eigenvalues	2
3.	An Example	4
4.	Three–Dimensional Gradient Ricci Solitons	6
Ret	ferences	9

1. INTRODUCTION

For $n \ge 3$, let (M^n, g) be a smooth Riemannian manifold and consider a Codazzi tensor T on M^n , i. e., a symmetric bilinear form satisfying the Codazzi equation

$$(\nabla_X \mathbf{T})(Y, Z) = (\nabla_Y \mathbf{T})(X, Z),$$

for every tangent vectors X, Y, Z.

In the book *Einstein Manifolds* [1], by Besse, it is proved that if a Riemannian manifold (M^n, g) admits a Codazzi tensor T such that at every point of M^n , T has exactly two distinct eigenvalues, then

- if the constant multiplicities of the two eigenspaces are larger than one, (M^n, g) is locally a Riemannian product,
- if the above multiplicities are respectively 1 and n-1 and the trace of T is constant, then (M^n, g) is locally a warped product of an (n-1)-dimensional Riemannian manifolds on an interval of \mathbb{R} .

For more details, we refer the reader to discussion 16.12 in [1].

Before showing this result, Besse states "... a similar argument works without this hypothesis [that trace of T is constant]". Recently in [7], G. Merton provided a counterexample to the local warping structure, showing that the last Besse's statement is false. In [7], he also discusses some possible extra hypotheses, weaker than *trace of* T *constant*, under which the local warped structure can be obtained.

Date: May 23, 2018.

Our goal here is to describe, without adding extra hypotheses to Besse's statement, what is the local geometric structure of a Riemannian manifold admitting a Codazzi tensor with exactly two distinct eigenvalues. Essentially, one has that the manifold may present zones where it is a warped product on a interval and zones where it is not. In this latter case, it turns out that the manifold admits a local totally geodesics foliation. This is the content of our Theorem 2.1.

In Section 3, we will give an example of a Riemannian manifold where both the situations (local warped product structure and local totally geodesics foliation) described in our structure theorem are present at the same time.

Finally, in the last section, we will show how this Codazzi tensors theory can be applied to the classification of gradient Ricci solitons.

Acknowledgments. The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). They are supported by the GNAMPA project "Equazioni di evoluzione geometriche e strutture di tipo Einstein".

2. CODAZZI TENSORS WITH TWO DISTINCT EIGENVALUES

In this section we present the statement and the proof of our main theorem.

Theorem 2.1. Let T be a Codazzi tensor on (M^n, g) , with $n \ge 3$. Suppose that at every point of M^n , the tensor T has exactly two distinct eigenvalues ρ and σ of multiplicity 1 and n - 1, respectively. Finally, we let $W = \{p \in M^n | d\sigma(p) \neq 0\}$. Then, we have that

- (1) The closed set $\overline{W} = W \cup \partial W$ with the metric $g|_{\overline{W}}$ is locally isometric to the warped product of some (n-1)-dimensional Riemannian manifold on an interval of \mathbb{R} and σ is constant along the "leaves" of the warped product.
- (2) The boundary of W, if present, is given by the disjoint union of connected totally geodesic hypersurfaces where σ is constant.
- (3) Each connected component of the complement of \overline{W} in M, if present, has σ constant and it is foliated by totally geodesic hypersurfaces.

The (n-1)-dimensional tangent subspaces to the above warping hypersurfaces at point (1) and to the totally geodesic hypersurfaces at points (2) and (3) are the eigenspaces of T with respect to σ .

Proof. Since the Codazzi tensor T has exactly two distinct eigenvalues ρ and σ of multiplicity 1 and n - 1, respectively, we have by Proposition 16.11 in [1] that the tangent bundle TM of M splits as the orthogonal direct sum of two *integrable* eigendistributions: a line field V_{ρ} and a codimension one distribution V_{σ} with totally *umbilical* leaves, which means that the second fundamental form h of each leaf is proportional to the metric g^{σ} , induced by g on V_{σ} .

To fix the notations, we will denote by ∇ the Levi–Civita connection of the metric g on M^n and we recall that the (scalar) second fundamental form of a leaf L of the codimension one distribution V_{σ} can be defined as

$$h(X,Y) = -g(\nabla_X Y,\nu),$$

where *X* and *Y* are vector fields along *L* and ν is a choice of a unit normal vector field to *L*. The fact that *L* is umbilical means that, for every couple of vector fields *X*, *Y* tangent to *L*,

we have

$$h(X,Y) = \frac{\mathrm{H}}{n-1} g^{\sigma}(X,Y) \, ,$$

where H, the mean curvature of L, is defined as the trace of h with respect to g^{σ} .

Since $n \ge 3$, we have that the codimension one distribution V_{σ} has dimension strictly bigger than one. Thus, we infer from Proposition 16.11 in [1] that the eigenfunction σ must be constant along the leaves of V_{σ} . In particular, whenever $d\sigma \neq 0$, the leaves of V_{σ} are locally regular level sets of σ .

To proceed, we fix a point $p \in M$ and we consider a local coordinate system (x^0, \ldots, x^{n-1}) adapted to the leaves of V_{σ} on a neighborhood U of p. This means that $\partial/\partial x^0 \in V_{\rho}$ and $\partial/\partial x^j \in V_{\sigma}$, for $j = 1, \ldots, n-1$. In this chart, the unit vector field $\nu = (\partial/\partial x^0)/\sqrt{g_{00}}$ is normal to any leaf of the distribution V_{σ} and since the two eigendistributions are mutually orthogonal we immediately get $g_{0j} = 0$ and $T_{0j} = 0$, for $j = 1, \ldots, n-1$. If L is the leaf of V_{σ} through the point p, the second fundamental form of L about p and the umbilicity condition can be written as

$$h_{ij} = -\left\langle \nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^j}, \nu \right\rangle = -\left\langle \nabla_{\frac{\partial}{\partial x^i}} \frac{\partial}{\partial x^j}, \frac{\partial}{\partial x^0} \right\rangle / \sqrt{g_{00}} = -\Gamma_{ij}^0 \sqrt{g_{00}} = \frac{\mathrm{H}}{n-1} g_{ij}^{\sigma}, \qquad (2.1)$$

for i, j = 1, ..., n - 1.

Denoting by ∇^{σ} the Levi–Civita connection of the induced metric g^{σ} , the Codazzi–Mainardi equations (see Theorem 1.72 in [1]) read

$$\left(\nabla^{\sigma}_{\frac{\partial}{\partial x^{i}}}h\right)\left(\frac{\partial}{\partial x^{j}},\frac{\partial}{\partial x^{k}}\right) - \left(\nabla^{\sigma}_{\frac{\partial}{\partial x^{j}}}h\right)\left(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{k}}\right) = \left\langle \operatorname{Rm}\left(\frac{\partial}{\partial x^{i}},\frac{\partial}{\partial x^{j}}\right)\frac{\partial}{\partial x^{k}},\nu\right\rangle.$$
(2.2)

Using the umbilicity property (2.1) of *L* and tracing the left hand side of equation (2.2) with the inverse of the metric $(g^{\sigma})^{ik}_{\sigma} = g^{ik}$, we get

$$g^{ik} \bigg[\left(\nabla^{\sigma}_{\frac{\partial}{\partial x^{i}}} h \right) \left(\frac{\partial}{\partial x^{j}}, \frac{\partial}{\partial x^{k}} \right) - \left(\nabla^{\sigma}_{\frac{\partial}{\partial x^{j}}} h \right) \left(\frac{\partial}{\partial x^{i}}, \frac{\partial}{\partial x^{k}} \right) \bigg] = \frac{1}{n-1} \partial_{j} \mathbf{H} - \partial_{j} \mathbf{H} = -\frac{n-2}{n-1} \partial_{j} \mathbf{H}.$$

Tracing also the right hand side, we get

$$-\frac{n-2}{n-1}\partial_{j}\mathbf{H} = g^{ik} \left\langle \operatorname{Rm}\left(\frac{\partial}{\partial x^{i}}, \frac{\partial}{\partial x^{j}}\right) \frac{\partial}{\partial x^{k}}, \nu \right\rangle = \operatorname{Ric}\left(\frac{\partial}{\partial x^{j}}, \nu\right) = \operatorname{Ric}_{0j}/\sqrt{g_{00}}, \quad (2.3)$$

as $g^{i0} = 0$ when $i \ge 1$ and $\langle \operatorname{Rm}(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}) \frac{\partial}{\partial x^k}, \nu \rangle$ is equal to zero if i = k = 0. Now, it is a general fact (see Corollary 16.17 in [1]) that every Codazzi tensor T commutes with the Ricci tensor, that is, $g^{kl} T_{ik} \operatorname{Ric}_{lj} = g^{kl} \operatorname{Ric}_{ik} T_{lj}$. In particular,

$$\rho \operatorname{Ric}_{0j} = g^{kl} \operatorname{T}_{0k} \operatorname{Ric}_{lj} = g^{kl} \operatorname{Ric}_{0k} \operatorname{T}_{lj} = \sigma g^{kl} \operatorname{Ric}_{0k} g_{lj} = \sigma \operatorname{Ric}_{0j},$$

hence, $\operatorname{Ric}_{0j} = 0$ for every j = 1, ..., n - 1, as $\rho \neq \sigma$ in *U*. We conclude by equation (2.3) that the mean curvature H is constant along every connected component of *L*, hence, the same conclusion holds for any leaf of V_{σ} .

Next, we recall from Proposition 16.11 (ii) in [1] that the eigenvalue σ is constant along the leaves of V_{σ} , thus, in our local chart, it only depends on the x^0 variable. Moreover, by the same proposition, one has that

$$\mathbf{H} = \frac{1}{\rho - \sigma} \frac{\partial \sigma}{\partial x^0}.$$
 (2.4)

From this we deduce that the connected component of the V_{σ} -leaves through critical points of σ are minimal and by the umbilicity they are also totally geodesic. This gives the description at the point (3) of the (possibly non present) interior of the set where $d\sigma = 0$.

We pass now to consider the open set $W \subset M$ given by the complement of the critical points of σ in M. We are going to prove that ρ is locally constant on the connected component of the V_{σ} -leaves which are sitting in W. To see this, it is sufficient to take the (coordinate) derivative of both sides of relation (2.4) with respect to x^{j} , for j = 1, ..., n - 1. This gives

$$0 = \partial_j \mathbf{H} = -\frac{1}{(\rho - \sigma)^2} \frac{\partial \rho}{\partial x^j} \frac{\partial \sigma}{\partial x^0} + \frac{1}{\rho - \sigma} \frac{\partial^2 \sigma}{\partial x^j \partial x^0} = -\frac{1}{(\rho - \sigma)^2} \frac{\partial \rho}{\partial x^j} \frac{\partial \sigma}{\partial x^0},$$

where we used the symmetry of the second derivative together with the constancy of σ along the V_{σ} -leaves. Since in our coordinates $d\sigma = \partial_0 \sigma dx^0$ and $d\sigma \neq 0$ in W, the claim follows. To conclude, we observe that the boundary of W (if any) can be described as a suitable union of connected component of level sets of σ . By continuity the eigenvalue ρ must be locally constant also on ∂W .

To show that g has a warped product structure on $\overline{W} = W \cup \partial W$, we first observe that the condition $\partial_j \rho = 0$, for j = 1, ..., n - 1, combined with [1, Proposition 16.11–(ii)], implies that V_{ρ} is a *geodesic* line distribution in \overline{W} . This means that $\nabla_{\nu}\nu = 0$, which easily implies $\Gamma_{00}^j = 0$, hence $\partial_j g_{00} = 0$, for every j = 1, ..., n - 1. Equation (2.1) then yields

$$\frac{\partial g_{ij}}{\partial x^0} = -2\Gamma^0_{ij} = \frac{2(\mathrm{H}/\sqrt{g_{00}})}{n-1}g_{ij}$$

Since H and g_{00} are constant along V_{σ} , one has that

$$\frac{\partial g_{ij}}{\partial x^0}(x^0,\ldots,x^{n-1}) = \varphi(x^0) g_{ij}(x^0,\ldots,x^{n-1})$$

for some function φ depending only on the x^0 variable. Setting $\psi(x^0) = d\varphi/dx^0$, one has that $e^{-\psi}g_{ij}$ does not depend on the x^0 variable. Thus, for every i, j = 1, ..., n-1, we can write

$$g_{ij}(x^0,\ldots,x^{n-1}) = e^{\psi(x^0)} G_{ij}(x^1,\ldots,x^{n-1}),$$

for some suitable functions G_{ij} . This prove that g has a local warped product structure in \overline{W} and the proof is complete.

Remark 2.2. If the metric is analytic and the Riemannian manifold (M^n, g) is connected, the presence of an open set where σ is constant implies that everywhere σ is constant and $d\sigma = 0$, hence $W = \emptyset$. In the opposite case $\overline{W} = M^n$ and the totally geodesic hypersurfaces (where σ is constant) whose union gives ∂W are locally finite.

Hence, in the analytic case we have a dichotomy: either the whole manifold is locally a warped product or it is globally foliated by totally geodesic hypersurfaces.

We show now that actually the two situations described in Theorem 2.1 can be both present in a Riemannian manifold if the metric is only smooth but not analytic.

We follow the line of Merton [7].

Let $M = \mathbb{R} \times \mathbb{S}^1 \times \mathbb{S}^1$ be endowed with the Riemannian metric

$$g(t, x, y) = \left(\sigma(t) - \rho(t, x, y)\right)^{-2} dt^2 + \sigma dx^2 + \sigma dy^2,$$

where $\sigma : \mathbb{R} \to \mathbb{R}^+$ and $\rho : M \to \mathbb{R}$ are smooth functions, such that:

• The function σ is monotone increasing from 1 to 2, with $\sigma' > 0$, in the interval $(-\infty, -1)$, constant equal to 2 in the interval [-1, 1] and again monotone increasing from 2 to 3, with $\sigma' > 0$, in the interval $(1, +\infty)$.

- The function ρ is equal to 3σ when $t \in (-\infty, -1]$ or $t \in [1, +\infty)$, for every $(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1$.
- For $t \in (-1,1)$ and every $(x, y) \in \mathbb{S}^1 \times \mathbb{S}^1$, the function ρ is nonconstant on the leaves $\{t\} \times \mathbb{S}^1 \times \mathbb{S}^1$, in particular it cannot be three times the function σ .

We then define the (1,1)-tensor T as follows

$$T(\partial_t) = \rho(t, x, y)\partial_t$$

$$T(\partial_x) = \sigma(t)\partial_x$$

$$T(\partial_y) = \sigma(t)\partial_y$$

and we will show that T is a Codazzi tensor. The (0,2)-version of T reads

$$T_{tt} = T_t^t g_{tt} = \frac{\rho}{(\sigma - \rho)^2}$$
$$T_{xx} = T_x^x g_{xx} = \sigma^2$$
$$T_{yy} = T_y^y g_{yy} = \sigma^2$$

and all the other components are null.

The Christoffel symbols of the metric g are given by

$$\begin{split} \Gamma^t_{tt} &= -\left(\sigma - \rho\right)^{-1} (\sigma' - \partial_t \rho) \\ \Gamma^i_{tt} &= -\sigma^{-1} (\sigma - \rho)^{-3} \partial_i \rho \\ \Gamma^t_{it} &= (\sigma - \rho)^{-1} \partial_i \rho \\ \Gamma^i_{jt} &= \sigma^{-1} \sigma' \delta^i_j / 2 \\ \Gamma^t_{ij} &= - (\sigma - \rho)^2 \sigma' \delta_{ij} / 2 \\ \Gamma^k_{ij} &= 0 \,, \end{split}$$

where the indices i, j, k can only be x and y. Thus we compute (we skip the trivial checks)

$$\nabla_y \mathbf{T}_{xx} - \nabla_x \mathbf{T}_{yx} = \partial_y \sigma^2 - 2\mathbf{T}_{xp} \Gamma_{xy}^p + \mathbf{T}_{yp} \Gamma_{xx}^p + \mathbf{T}_{xp} \Gamma_{xy}^p$$
$$= -\sigma^2 \Gamma_{xy}^x + \sigma^2 \Gamma_{xx}^y$$
$$= 0$$
$$\nabla_t \mathbf{T}_{xx} - \nabla_x \mathbf{T}_{tx} = \partial_t \sigma^2 - 2\mathbf{T}_{xp} \Gamma_{xt}^p + \mathbf{T}_{tp} \Gamma_{xx}^p + \mathbf{T}_{xp} \Gamma_{xt}^p$$
$$= 2\sigma \sigma' - \sigma^2 \Gamma_{xt}^x + \frac{\rho}{(\sigma - \rho)^2} \Gamma_{xx}^t$$

$$= 2\sigma\sigma' - \sigma^2 \Gamma_{xt}^{*} + \frac{1}{(\sigma - \rho)^2} \Gamma_{xx}^{*}$$
$$= 2\sigma\sigma' - \sigma\sigma'/2 - \sigma'\rho/2$$
$$= (3\sigma - \rho)\sigma'/2$$

$$\nabla_{x} T_{tt} - \nabla_{t} T_{xt} = \partial_{x} \left(\frac{\rho}{(\sigma - \rho)^{2}} \right) - 2 T_{tp} \Gamma_{tx}^{p} + T_{xp} \Gamma_{tt}^{p} + T_{tp} \Gamma_{xt}^{p}$$

$$= \frac{\partial_{x} \rho}{(\sigma - \rho)^{2}} + \frac{2\rho \partial_{x} \rho}{(\sigma - \rho)^{3}} - \frac{\rho}{(\sigma - \rho)^{2}} \Gamma_{tx}^{t} + \sigma^{2} \Gamma_{tt}^{x}$$

$$= \frac{\partial_{x} \rho}{(\sigma - \rho)^{2}} + \frac{2\rho \partial_{x} \rho}{(\sigma - \rho)^{3}} - \frac{\rho \partial_{x} \rho}{(\sigma - \rho)^{3}} - \frac{\sigma \partial_{x} \rho}{(\sigma - \rho)^{3}}$$

$$= 0$$

$$\nabla_{t} T_{xy} - \nabla_{x} T_{ty} = - T_{xp} \Gamma_{ty}^{p} - T_{yp} \Gamma_{tx}^{p} + T_{yp} \Gamma_{tx}^{p} + T_{tp} \Gamma_{xy}^{p}$$

$$= -\sigma^{2} \Gamma_{ty}^{x} + \frac{\rho}{(\sigma - \rho)^{2}} \Gamma_{ty}^{t}$$

$$= 0$$

$$\nabla_{x} T_{yt} - \nabla_{y} T_{xt} = - T_{yp} \Gamma_{tx}^{p} - T_{tp} \Gamma_{xy}^{p} + T_{xp} \Gamma_{ty}^{p} + T_{tp} \Gamma_{xy}^{p}$$

$$= -\sigma^{2} \Gamma_{tx}^{y} + \sigma^{2} \Gamma_{ty}^{x}$$

$$= 0.$$

Hence, by our choices for the functions σ and ρ , the tensor T is a Codazzi tensor.

It is easy to see that in the zone where σ is nonconstant, the manifold is a warped product on a interval of \mathbb{R} , instead, in the zone $(-1,1) \times \mathbb{S}^1 \times \mathbb{S}^1$, if the function ρ is suitably chosen nonconstant on the leaves $\{t\} \times \mathbb{S}^1 \times \mathbb{S}^1$, it can be checked that (M, g) is not a warped product on an interval (actually, in this example, it is incidentally a warped product on $\mathbb{S}^1 \times \mathbb{S}^1$), see the careful analysis in [7].

Hence, the two situations described in Theorem 2.1 are both present in this example.

4. THREE-DIMENSIONAL GRADIENT RICCI SOLITONS

Let (M^3, g) be a three–dimensional gradient Ricci soliton, that is a Riemannian manifold satisfying the equation

$$\operatorname{Ric} + \nabla^2 f = \lambda g \tag{4.1}$$

for some smooth function $f: M^3 \to \mathbb{R}$ and some constant $\lambda \in \mathbb{R}$.

Lemma 4.1. On every three–dimensional gradient Ricci soliton the tensor

$$\Gamma = \left(\operatorname{Ric} - \frac{1}{2} \operatorname{R} g \right) e^{-f}$$

is a Codazzi tensor.

Proof. Let (M^3, g) be a three dimensional gradient Ricci soliton satisfying equation (4.1) and let

$$\mathbf{T}_{ij} = \left(\mathbf{R}_{ij} - \frac{1}{2}\mathbf{R}\,g_{ij}\right)e^{-f} \,.$$

We want to prove that T is a Codazzi tensor, i.e. we have to show that

$$\nabla_k \mathbf{T}_{ij} = \nabla_j \mathbf{T}_{ik} \,,$$

for every i, j, k = 1, 2, 3. One has

$$\nabla_k \mathbf{T}_{ij} - \nabla_j \mathbf{T}_{ik} = \left[\nabla_k \mathbf{R}_{ij} - \nabla_j \mathbf{R}_{ik} - \frac{1}{2} (\nabla_k \mathbf{R} g_{ij} - \nabla_j \mathbf{R} g_{ik}) \right] e^{-f} + \left[\frac{1}{2} \mathbf{R} (\nabla_k f g_{ij} - \nabla_j f g_{ik}) - \nabla_k f \mathbf{R}_{ij} + \nabla_j f \mathbf{R}_{ik} \right] e^{-f} .$$
(4.2)

On the other hand, the following two identities hold on any gradient Ricci soliton (for a proof, see [5], for instance)

$$\nabla_k \mathbf{R} = 2\nabla_p f \,\mathbf{R}_{pk} \tag{4.3}$$

$$\nabla_k \mathbf{R}_{ij} - \nabla_j \mathbf{R}_{ik} = -\mathbf{R}_{kjip} \nabla_p f \,. \tag{4.4}$$

Moreover, since we are in dimension three, one has the decomposition of the Riemann tensor

$$R_{kjip} = R_{ik}g_{jp} - R_{kp}g_{ij} + R_{jp}g_{ik} - R_{ij}g_{kp} - \frac{1}{2}R(g_{ik}g_{jp} - g_{ij}g_{kp}).$$

Combining with equation (4.4), we obtain

$$\nabla_k \mathbf{R}_{ij} - \nabla_j \mathbf{R}_{ik} = -\nabla_j f \mathbf{R}_{ik} + \nabla_p f \mathbf{R}_{kp} g_{ij} - \nabla_p f \mathbf{R}_{jp} g_{ik} + \nabla_k f \mathbf{R}_{ij} - \frac{1}{2} \mathbf{R} (\nabla_k f g_{ij} - \nabla_j f g_{ik}).$$

Hence, substituting this in equation (4.2) and using relation (4.3), we immediately get

$$\nabla_k \mathbf{T}_{ij} - \nabla_j \mathbf{T}_{ik} = 0.$$

As an application of this lemma and the results of the previous sections, we have the following theorem.

Theorem 4.2. Let (M^3, g) be a complete, three–dimensional, simply connected Riemannian manifold, which is a steady, gradient Ricci soliton and assume that there exists an open subset $U \subset M$, where the Ricci tensor of g has at most two distinct eigenvalues. Then, either the manifold splits a line or it is locally conformally flat.

Proof. By [9], as (M^3, g) is complete, this gradient Ricci soliton generates an ancient Ricci flow. Then, by the result [3, Corollary 2.4] the evolving manifold, hence the Ricci soliton, must have nonnegative sectional curvatures. Moreover, it is well known, by the properties of the parabolic equations, that the metric g must be analytic (see [4, Chapter 3, Section 2]).

If at least one sectional curvature is zero at some point, then the manifold (M^3, g) "splits a line" (see [4]), that is, it is isometric to the Riemannian product of \mathbb{R} with a surface. Hence, we will assume in the rest of the proof that all the sectional curvatures are strictly positive everywhere.

The analyticity of the metric implies that either at every point of the open subset U all of the three eigenvalues of the Ricci tensor coincide, or there is another, possibly smaller, open subset W of M^3 such that the Ricci tensor has everywhere in W exactly two distinct eigenvalues.

The first case cannot occur, since (U, g) would be locally isometric to an Einstein manifold with positive curvature, then, by analyticity, (M^3, g) must be isometric to the sphere \mathbb{S}^3 , and this would contradict the fact that every compact steady Ricci soliton is Ricci flat. Thus, we assume from now on that there exists an open subset W where the Ricci tensor has exactly two distinct eigenvalues. This implies that on W the Codazzi tensor $T = (Ric - Rg/2)e^{-f}$ has two distinct eigenvalues σ , with multiplicity 2, and ρ , with multiplicity 1.

As Remark 2.2 applies to this case, by Theorem 2.1, we have that two possible subcases: either around every point of W the manifold is locally isometric to a warped product of a surface on an interval, or the eigenvalue σ of the Codazzi tensor T is constant on W, hence on the whole M^3 by analyticity. On the other hand, since the curvature of (M^3, g) is strictly positive, (M^3, g) cannot admit equidistant totally geodesic submanifolds of dimension greater than one by the second Rauch comparison theorem (see [6]) and this latter subcase is excluded.

In the former subcase, the leaves of the distribution V_{σ} are umbilical and the two eigenvalues of T are constant on every leaf. Let *L* be a connected component of a leaf of V_{σ} . By Gauss formula, one has that the scalar curvature of the induced metric g^{σ} is given by

$$R^{\sigma} = R - 2R_0^0 + H^2/2, \qquad (4.5)$$

where H denotes the mean curvature of *L*. Since *g* has positive sectional curvature, one has $Rg - 2 \operatorname{Ric} > 0$ and we obtain that R^{σ} is positive. We want to prove that *g* is locally a warped product on an interval of \mathbb{R} of two–dimensional fibers with constant positive curvature. Hence, we have to show that R^{σ} is constant on *L*.

First of all, we observe that by the same reasoning as in the proof of Theorem 2.1, we know that H is constant on *L*. Thus, it remains to show that also the quantity $R - 2R_0^0$ is constant on *L*. The fact that all the eigenvalues of the tensor T are constant on *L*, implies that the trace of T

$$tr(T) = -Re^{-f}/2$$

is constant on *L*. We claim that also *f* has to be constant on *L*.

Using the adapted coordinate system as in the proof of Theorem 2.1, we assume by contradiction that $\partial_j f \neq 0$, for some $j \in \{1, 2\}$ at some point of *L*. As $\partial_j tr(T) = 0$ it follows that ∇f and ∇R are parallel and

$$\mathbf{R}\nabla f = \nabla \mathbf{R} = 2\mathrm{Ric}(\nabla f, \cdot),$$

hence, ∇f is an eigenvalue of the Ricci tensor. Being $\partial_j f \neq 0$, then it must be

$$\mathbf{R}\nabla f = 2\mathbf{Ric}_{j}^{j}\nabla f$$
,

which is a contradiction, as $R = 2Ric_j^j + Ric_0^0$ and Ric_0^0 is positive by assumption (notice that we have used the fact that the Ricci tensor has exactly two distinct eigenvalues with the same eigendistributions as the Codazzi tensor T). Thus, we have proved that f is constant on L which implies that R is constant on L too. Then, it follows, by the definition of T and the fact that its eigenvalues are constant on L, that also the eigenvalues of the Ricci tensor are constant on L. In particular, R_0^0 is constant on L.

By relation (4.5), we conclude that *L* has positive constant scalar curvature \mathbb{R}^{σ} . Hence, the leaf *L* is locally isometric to \mathbb{S}^2 and the metric *g* in *W* is locally a warped product of an interval with two–dimensional spherical fibers. In particular it is locally conformally flat.

Using once again the analyticity, we can conclude that since $(W, g|_W)$ is a locally conformally flat open subset of (M^3, g) , then the whole (M^3, g) must be locally conformally flat. This completes the proof.

Remark 4.3. By the same argument, the conclusion of this theorem also holds for complete, three–dimensional, simply connected, *expanding*, gradient Ricci solitons with nonnegative sectional curvatures.

Remark 4.4. Three–dimensional, locally conformally flat, gradient steady Ricci solitons were classified by Cao–Chen [2]. In particular, under the assumptions of Theorem 4.2 we have that (M^3, g) is isometric to \mathbb{R}^3 , the *Bryant* soliton or the Riemannian product of \mathbb{R} with *Hamilton's cigar*. Note that if a three–dimensional, gradient steady Ricci soliton splits a line, then it must be the Riemannian product of \mathbb{R} with a two–dimensional complete simply connected gradient steady soliton, that is, \mathbb{R}^2 or Hamilton's cigar.

A NOTE ON CODAZZI TENSORS

REFERENCES

- 1. A. L. Besse, Einstein manifolds, Springer–Verlag, Berlin, 2008.
- 2. H.-D. Cao and Q. Chen, On locally conformally flat gradient steady Ricci solitons, Trans. Amer. Math. Soc. 364 (2012), 2377–2391.
- 3. B.-L. Chen, Strong uniqueness of the Ricci flow, J. Diff. Geom. 82 (2009), 363-382.
- 4. B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo, and L. Ni, *The Ricci flow: techniques and applications. Part II. Analytic aspects*, Mathematical Surveys and Monographs, vol. 144, American Mathematical Society, Providence, RI, 2008.
- 5. M. Eminenti, G. La Nave, and C. Mantegazza, *Ricci solitons: the equation point of view*, Manuscripta Math. **127** (2008), no. 3, 345–367.
- 6. D. Gromoll, W. Klingenberg, and W. Meyer, *Riemannsche Geometrie im Großen*, Lecture Notes in Mathematics, Vol. 55, Springer–Verlag, Berlin, 1975.
- 7. G. Merton, Codazzi tensors with two eigenvalue functions, ArXiv Preprint Server http://arxiv.org, 2011.
- 8. L. Ni and N. Wallach, On a classification of gradient shrinking solitons, Math. Res. Lett. 15 (2008), no. 5, 941–955.
- 9. Z.-H. Zhang, On the completeness of gradient Ricci solitons, Proc. Amer. Math. Soc. 137 (2009), no. 8, 2755–2759.

(Giovanni Catino) DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, MILANO, ITALY, 20133

E-mail address, G. Catino: giovanni.catino@polimi.it

(Carlo Mantegazza) SCUOLA NORMALE SUPERIORE DI PISA, PIAZZA DEI CAVALIERI7, PISA, ITALY, 56126 *E-mail address*, C. Mantegazza: c.mantegazza@sns.it

(Lorenzo Mazzieri) SCUOLA NORMALE SUPERIORE DI PISA, PIAZZA DEI CAVALIERI7, PISA, ITALY, 56126 *E-mail address*, L. Mazzieri: 1.mazzieri@sns.it