

# ERROR ANALYSIS FOR A FINITE DIFFERENCE SCHEME FOR AXISYMMETRIC MEAN CURVATURE FLOW OF GENUS-0 SURFACES

KLAUS DECKELNICK<sup>†</sup> AND ROBERT NÜRNBERG<sup>‡</sup>

**Abstract.** We consider a finite difference approximation of mean curvature flow for axisymmetric surfaces of genus zero. A careful treatment of the degeneracy at the axis of rotation for the one dimensional partial differential equation for a parameterization of the generating curve allows us to prove error bounds with respect to discrete  $L^2$ - and  $H^1$ -norms for a fully discrete approximation. The theoretical results are confirmed with the help of numerical convergence experiments. We also present numerical simulations for some genus-0 surfaces, including for a non-embedded self-shrinker for mean curvature flow.

**Key words.** mean curvature flow, axisymmetry, finite differences, error analysis, self-shrinker

**AMS subject classifications.** 65M60, 65M12, 65M15, 53C44, 35K55

**1. Introduction.** Consider a family of surfaces  $(\mathcal{S}(t))_{t \in [0, T]} \subset \mathbb{R}^3$  evolving by mean curvature flow, i.e.

$$(1.1) \quad \mathcal{V}_{\mathcal{S}} = k_m \quad \text{on } \mathcal{S}(t).$$

Here,  $\mathcal{V}_{\mathcal{S}}$  denotes the normal velocity of  $\mathcal{S}(t)$  in the direction of the normal  $\vec{\nu}_{\mathcal{S}(t)}$ , and  $k_m$  is the mean curvature of  $\mathcal{S}(t)$ , i.e. the sum of its principal curvatures. As the  $L^2$ -gradient flow for the area functional, (1.1) is one of the most important geometric evolution equations with applications in materials science and image processing. We refer the reader to [12, 17] for an introduction and important results of mean curvature flow.

In this paper we are concerned with the numerical approximation of solutions of (1.1) using a parametric approach. If  $\vec{X} : \mathcal{M} \times [0, T] \rightarrow \mathbb{R}^3$  is a family of embeddings such that  $\mathcal{S}(t) = \vec{X}(\mathcal{M}, t)$ , then (1.1) is satisfied if  $\vec{X}_t \circ \vec{X}^{-1} = k_m \vec{\nu}_{\mathcal{S}(t)}$  on  $\mathcal{S}(t)$ . Making use of the fact that the mean curvature vector  $k_m \vec{\nu}_{\mathcal{S}(t)}$  can be written as  $\Delta_{\mathcal{S}(t)} \vec{\text{id}}$ , where  $\Delta_{\mathcal{S}(t)}$  denotes the Laplace–Beltrami operator on  $\mathcal{S}(t)$ , Dziuk [10] suggested a finite element method in order to approximate solutions of (1.1). While this approach has been widely used in the following years, the numerical analysis of the method remained open. Only recently, Kovács, Li and Lubich [16] obtained error estimates for a parametric approach that uses not only the position  $\vec{X}$ , but also the mean curvature and the normal as variables. Both approaches are based on evolution equations in which the velocity vector points purely in normal direction, which may lead to degenerate meshes at the discrete level. A way to tackle this issue is to introduce a suitable additional tangential motion in such a way, that mesh points are better distributed on the approximate surface. Corresponding schemes have been suggested by Barrett, Garcke and Nürnberg [4], as well as by Elliott and Fritz [13], using DeTurck’s trick. For the approach from [13], error bounds for a finite difference scheme in the case of surfaces of torus type have recently been obtained in [18]. For more details on the numerical approximation of geometric evolution equations we refer to the review articles [8, 6].

---

<sup>†</sup>Institut für Analysis und Numerik, Otto-von-Guericke-Universität Magdeburg, 39106 Magdeburg, Germany

<sup>‡</sup>Department of Mathematics, University of Trento, Trento, Italy

In what follows, we are interested in the case that the evolving surfaces are axisymmetric with respect to the  $x_2$ -axis, i.e. we assume that there exists a mapping  $\vec{x}(\cdot, t) : [0, 1] \rightarrow \mathbb{R}_{\geq 0} \times \mathbb{R}$  such that

$$\mathcal{S}(t) = \left\{ (\vec{x}(\rho, t) \cdot \vec{e}_1 \cos \theta, \vec{x}(\rho, t) \cdot \vec{e}_2, \vec{x}(\rho, t) \cdot \vec{e}_1 \sin \theta)^T : \rho \in [0, 1], \theta \in [0, 2\pi] \right\}.$$

As shown in [5, 2], the law (1.1) translates into the following evolution equation for the curves  $(\Gamma(t))_{t \in [0, T]}$  parameterised by  $\vec{x}(\cdot, t)$ :

$$(1.2) \quad \vec{x}_t \cdot \vec{\nu} = \varkappa - \frac{\vec{\nu} \cdot \vec{e}_1}{\vec{x} \cdot \vec{e}_1},$$

where  $\vec{\nu}$  is a unit normal to  $\Gamma(t)$  and  $\varkappa = \vec{\varkappa} \cdot \vec{\nu}$  denotes curvature, with  $\vec{\varkappa} = \frac{1}{|\vec{x}_\rho|} \left( \frac{\vec{x}_\rho}{|\vec{x}_\rho|} \right)_\rho$  the curvature vector. We note that without the last term on the right hand side of (1.2), the problem collapses to curve shortening flow,

$$(1.3) \quad \vec{x}_t \cdot \vec{\nu} = \varkappa,$$

which is the analogue of (1.1) for curves. Since the relations (1.2) and (1.3) only prescribe the normal velocity, there is a certain freedom in choosing the tangential part of the velocity vector. Setting the tangential velocity to zero for (1.3) leads to the formulation  $\vec{x}_t = \frac{1}{|\vec{x}_\rho|} \left( \frac{\vec{x}_\rho}{|\vec{x}_\rho|} \right)_\rho$ , and optimal error bounds for a semidiscrete continuous-in-time finite element approximation of it have been obtained by Dziuk [11]. On the other hand, an application of DeTurck's trick gives rise to the formulation  $\vec{x}_t = \frac{\vec{x}_{\rho\rho}}{|\vec{x}_\rho|^2}$  for classical curve shortening flow. An error analysis for a corresponding semidiscrete finite element scheme has been first presented in [7], and this was later extended in [13] to the family of problems  $\alpha \vec{x}_t + (1 - \alpha)(\vec{x}_t \cdot \vec{\nu})\vec{\nu} = \frac{\vec{x}_{\rho\rho}}{|\vec{x}_\rho|^2}$ ,  $\alpha \in (0, 1]$ . Inspired by the ideas in [7], the present authors in [2] applied DeTurck's trick to the flow (1.2) to obtain the system

$$(1.4) \quad \vec{x}_t = \frac{\vec{x}_{\rho\rho}}{|\vec{x}_\rho|^2} - \frac{\vec{\nu} \cdot \vec{e}_1}{\vec{x} \cdot \vec{e}_1} \vec{\nu},$$

cf. [2, (1.7)]. Note that (1.4) is strictly parabolic and that a solution of (1.4) satisfies (1.2). The difference to curve shortening flow consists in the presence of the term  $\frac{\vec{\nu} \cdot \vec{e}_1}{\vec{x} \cdot \vec{e}_1}$ , which is the principal curvature related to the parallels of  $\mathcal{S}(t)$ . It is possible to rewrite (1.4) in the following divergence form

$$(1.5) \quad \vec{x} \cdot \vec{e}_1 |\vec{x}_\rho|^2 \vec{x}_t = ((\vec{x} \cdot \vec{e}_1) \vec{x}_\rho)_\rho - |\vec{x}_\rho|^2 \vec{e}_1,$$

giving rise to a natural variational formulation. On the basis of this weak formulation, a semi-implicit scheme using piecewise linear finite elements in space and a backward Euler method in time was suggested by the authors in [2]. In particular, in [2, Theorem 2.2] optimal error bounds both in  $H^1$  and  $L^2$  are obtained in the case of genus-1 surfaces. While the numerical method still performs well also for genus-0 surfaces, it is however not possible to apply the employed analysis to genus-0 surfaces. The reason for the additional difficulties in the genus-0 case comes from the different properties of the curves  $\Gamma(t)$ : for genus-1 surfaces,  $\Gamma(t)$  is a closed curve satisfying  $\vec{x} \cdot \vec{e}_1 > 0$  on  $[0, 1]$  so that this term is bounded strictly from below on compact time intervals, thus simplifying the analysis. In contrast, a description of a genus-0 surface

in our setting requires  $\Gamma(t)$  to be open with its endpoints lying on the  $x_2$ -axis, which means that  $\vec{x} \cdot \vec{e}_1 = 0$  at the endpoints of the interval  $[0, 1]$ . Furthermore, in order to guarantee smoothness of the surface  $\mathcal{S}(t)$ , the curve  $\Gamma(t)$  has to meet the  $x_2$ -axis at a right angle. In order to formulate the resulting initial-boundary problem, it is convenient to rewrite (1.4). To do so, we choose  $\vec{\nu} = \vec{\tau}^\perp$  with the unit tangent  $\vec{\tau} = \frac{\vec{x}_\rho}{|\vec{x}_\rho|}$  and  $\cdot^\perp$  denoting clockwise rotation by  $\frac{\pi}{2}$ . Observing that

$$(\vec{\nu} \cdot \vec{e}_1) \vec{\nu} = \frac{1}{|\vec{x}_\rho|^2} (\vec{x}_\rho^\perp \cdot \vec{e}_1) \vec{x}_\rho^\perp = \frac{1}{|\vec{x}_\rho|^2} (\vec{x}_\rho \cdot \vec{e}_2) \vec{x}_\rho^\perp,$$

we are led to the following system

$$(1.6a) \quad \vec{x}_t = \frac{\vec{x}_{\rho\rho}}{|\vec{x}_\rho|^2} - \frac{1}{|\vec{x}_\rho|^2} \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \vec{x}_\rho^\perp \quad \text{in } (0, 1) \times (0, T],$$

$$(1.6b) \quad \vec{x} \cdot \vec{e}_1 = 0, \quad \vec{x}_\rho \cdot \vec{e}_2 = 0 \quad \text{on } \{0, 1\} \times [0, T].$$

Since  $\vec{x}(\rho, t) \cdot \vec{e}_1 \rightarrow 0$ , as  $\rho \rightarrow \rho_0 \in \{0, 1\}$ , the last term in (1.6a) needs to be treated with care. Using the boundary conditions (1.6b), it is shown in (A.3) in Appendix A, with the help of L'Hospital's rule, that

$$\lim_{\rho \searrow 0} \left[ -\frac{1}{|\vec{x}_\rho(\rho, t)|^2} \frac{\vec{x}_\rho(\rho, t) \cdot \vec{e}_2}{\vec{x}(\rho, t) \cdot \vec{e}_1} \vec{x}_\rho^\perp(\rho, t) \right] = \frac{\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_2}{|\vec{x}_\rho(0, t)|^2} \vec{e}_2,$$

so that the expression acts like a second order operator close to the boundary without affecting the parabolicity of the problem. Nevertheless, the different behaviour of  $\frac{\vec{\nu} \cdot \vec{e}_1}{\vec{x} \cdot \vec{e}_1}$  in the interior and close to the boundary is a major problem for the analysis of a numerical scheme. Rather than using the variational form (1.5) that worked well for genus-1 surfaces, we shall introduce a scheme which directly discretises (1.6a) with the help of finite differences. Our main result are optimal error bounds measuring the error in discrete versions of the usual integral norms.

The paper is organised as follows. In Section 2, we formulate our assumptions on the solution of (1.6) and derive a number of properties that will be used in the error analysis. In the second part, we introduce our numerical scheme and provide an estimate for the consistency error. Section 3 is devoted to the proof of our main error estimates, which include an  $\mathcal{O}(h^2 + \Delta t)$  bound for a discrete  $H^1$ -norm. Finally, in Section 4 we present the results of several numerical simulations.

We end this section with a few comments about notation. Throughout,  $C$  denotes a generic positive constant independent of the mesh parameter  $h$  and the time step size  $\Delta t$ . At times  $\varepsilon$  will play the role of a (small) positive parameter, with  $C_\varepsilon > 0$  depending on  $\varepsilon$ , but independent of  $h$  and  $\Delta t$ .

## 2. Finite difference discretization.

**ASSUMPTION 2.1.** *Let  $\vec{x} : [0, 1] \times [0, T] \rightarrow \mathbb{R}_{\geq 0} \times \mathbb{R}$  be a solution of (1.6) such that  $\partial_t^i \partial_\rho^j \vec{x}$  exist and are continuous on  $[0, 1] \times [0, T]$  for all  $i, j \in \mathbb{N}_0$  with  $2i + j \leq 4$ . Furthermore, we assume that  $\vec{x}_\rho(\rho, t) \neq 0$  for all  $(\rho, t) \in [0, 1] \times [0, T]$ , as well as*

$$(2.1) \quad \vec{x} \cdot \vec{e}_1 > 0 \quad \text{in } (0, 1) \times [0, T].$$

It is beyond the scope of this paper to prove the existence of a solution to (1.6) with the above regularity. We note, however, that the well-posedness of the corresponding problem, in the case that the curves  $\Gamma(t)$  can be written as a graph, was recently studied in [14].

Let us collect a few properties of the solution which will be used in the error analysis. To begin, there exist constants  $0 < c_0 \leq C_0$  such that

$$(2.2) \quad c_0 \leq |\vec{x}_\rho| \leq C_0 \quad \text{in } [0, 1] \times [0, T].$$

Recalling (1.6b), we infer that  $\vec{x}_\rho(0, t) \cdot \vec{e}_1 \geq c_0$ ,  $\vec{x}_\rho(1, t) \cdot \vec{e}_1 \leq -c_0$  which together with (2.1) implies that there exist  $c_1 > 0, \delta > 0$  with

$$(2.3a) \quad \vec{x}_\rho \cdot \vec{e}_1 \geq \frac{1}{2}c_0 \quad \text{in } [0, \delta] \times [0, T],$$

$$(2.3b) \quad \vec{x}_\rho \cdot \vec{e}_1 \leq -\frac{1}{2}c_0 \quad \text{in } [1 - \delta, 1] \times [0, T],$$

$$(2.3c) \quad \vec{x} \cdot \vec{e}_1 \geq c_1 \quad \text{in } [\frac{1}{2}\delta, 1 - \frac{1}{2}\delta] \times [0, T].$$

Let us formally describe how this observation can be translated into an estimate on the solution. If we multiply (1.6a) by  $-\vec{x}_{\rho\rho}$  and integrate over  $[0, 1]$ , we find upon integration by parts and observing from (1.6b) that  $\vec{x}_t \cdot \vec{x}_\rho = 0$  on  $\{0, 1\} \times [0, T]$ , that

$$(2.4) \quad \frac{1}{2} \frac{d}{dt} \int_0^1 |\vec{x}_\rho|^2 d\rho + \int_0^1 \frac{|\vec{x}_{\rho\rho}|^2}{|\vec{x}_\rho|^2} d\rho - \int_0^1 \frac{1}{|\vec{x}_\rho|^2} \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \vec{x}_\rho^\perp \cdot \vec{x}_{\rho\rho} d\rho = 0.$$

Since (1.6b) implies  $\frac{\vec{x}_\rho^\perp(0, t)}{|\vec{x}_\rho(0, t)|} \approx -\vec{e}_2$  on  $[0, \delta]$ , we can rewrite the third term on  $[0, \delta]$ , on noting (2.2) and (2.3a), as

$$(2.5) \quad \begin{aligned} & - \int_0^\delta \frac{1}{|\vec{x}_\rho|^2} \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \vec{x}_\rho^\perp \cdot \vec{x}_{\rho\rho} d\rho \\ & \approx \int_0^\delta \frac{1}{|\vec{x}_\rho|} \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \vec{x}_{\rho\rho} \cdot \vec{e}_2 d\rho = \frac{1}{2} \int_0^\delta \frac{1}{|\vec{x}_\rho|} \frac{1}{\vec{x} \cdot \vec{e}_1} [(\vec{x}_\rho \cdot \vec{e}_2)^2]_\rho d\rho \\ & = \frac{1}{2} \left[ \frac{1}{|\vec{x}_\rho|} \frac{1}{\vec{x} \cdot \vec{e}_1} (\vec{x}_\rho \cdot \vec{e}_2)^2 \right]_0^\delta + \frac{1}{2} \int_0^\delta \frac{\vec{x}_\rho \cdot \vec{e}_1}{|\vec{x}_\rho|} \frac{(\vec{x}_\rho \cdot \vec{e}_2)^2}{(\vec{x} \cdot \vec{e}_1)^2} d\rho + \frac{1}{2} \int_0^\delta \frac{\vec{x}_\rho \cdot \vec{x}_{\rho\rho}}{|\vec{x}_\rho|^3} \frac{(\vec{x}_\rho \cdot \vec{e}_2)^2}{\vec{x} \cdot \vec{e}_1} d\rho \\ & \geq \frac{1}{4} \frac{c_0}{C_0} \int_0^\delta \frac{(\vec{x}_\rho \cdot \vec{e}_2)^2}{(\vec{x} \cdot \vec{e}_1)^2} d\rho + \frac{1}{2} \int_0^\delta \frac{\vec{x}_\rho \cdot \vec{x}_{\rho\rho}}{|\vec{x}_\rho|^3} \frac{(\vec{x}_\rho \cdot \vec{e}_2)^2}{\vec{x} \cdot \vec{e}_1} d\rho, \end{aligned}$$

so that we obtain  $L^2$ -control of  $\frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1}$  close to 0. A similar calculation applies close to 1, while the denominator  $\vec{x} \cdot \vec{e}_1$  is bounded away from 0 on  $[\delta, 1 - \delta]$  in view of (2.3c). Our aim is to mimic this argument within the error analysis (cf. Lemma 3.4). To do so, we will directly discretise (1.6a) using a finite difference scheme, and the discrete analogue of the above estimate is then obtained by multiplying with a suitable second order finite difference.

In order to define our finite difference scheme, let us introduce the set of grid points  $\mathcal{G}_h := \{q_0, q_1, \dots, q_J\}$ , where  $q_j = jh$  and  $h = \frac{1}{J}, j = 0, \dots, J$ . For a grid function  $\vec{v} : \mathcal{G}_h \rightarrow \mathbb{R}^2$  we write  $\vec{v}_j := \vec{v}(q_j), j = 0, \dots, J$ . Furthermore we associate with  $\vec{v}$  the following finite difference operators:

$$(2.6a) \quad \delta^- \vec{v}_j := \frac{\vec{v}_j - \vec{v}_{j-1}}{h}, \quad j = 1, \dots, J;$$

$$(2.6b) \quad \delta^+ \vec{v}_j := \delta^- \vec{v}_{j+1} = \frac{\vec{v}_{j+1} - \vec{v}_j}{h}, \quad j = 0, \dots, J-1;$$

$$(2.6c) \quad \delta^1 \vec{v}_j := \frac{1}{2}(\delta^+ \vec{v}_j + \delta^- \vec{v}_j) = \frac{\vec{v}_{j+1} - \vec{v}_{j-1}}{2h}, \quad j = 1, \dots, J-1.$$

$$(2.6d) \quad \delta^2 \vec{v}_j := \frac{\delta^+ \vec{v}_j - \delta^- \vec{v}_j}{h} = \frac{\vec{v}_{j+1} - 2\vec{v}_j + \vec{v}_{j-1}}{h^2}, \quad j = 1, \dots, J-1.$$

Two grid functions  $\vec{v}$  and  $\vec{w}$  satisfy the following summation by parts formula:

$$(2.7) \quad h \sum_{j=1}^J \delta^- \vec{v}_j \cdot \delta^- \vec{w}_j = -h \sum_{j=1}^{J-1} \vec{v}_j \cdot \delta^2 \vec{w}_j + \vec{v}_J \cdot \delta^- \vec{w}_J - \vec{v}_0 \cdot \delta^+ \vec{w}_0.$$

In addition, we introduce the following discrete norms and seminorms

$$(2.8) \quad |\vec{v}|_{0,h}^2 := \frac{1}{2}h|\vec{v}_0|^2 + h \sum_{j=1}^{J-1} |\vec{v}_j|^2 + \frac{1}{2}h|\vec{v}_J|^2; \quad |\vec{v}|_{1,h}^2 := h \sum_{j=1}^J |\delta^- \vec{v}_j|^2;$$

$$\|\vec{v}\|_{1,h}^2 := |\vec{v}|_{0,h}^2 + |\vec{v}|_{1,h}^2; \quad |\vec{v}|_{2,h}^2 := h \sum_{j=1}^{J-1} |\delta^2 \vec{v}_j|^2.$$

We also recall the following inverse inequality, as well as a discrete version of a well-known Sobolev type inequality.

LEMMA 2.2. *Let  $\vec{v} : \mathcal{G}_h \rightarrow \mathbb{R}^2$  be an arbitrary grid function. Then*

$$(2.9) \quad \max_{1 \leq k \leq J} |\delta^- \vec{v}_k| \leq h^{-\frac{1}{2}} |\vec{v}|_{1,h},$$

$$(2.10) \quad \max_{0 \leq k \leq J} |\vec{v}_k|^2 \leq |\vec{v}|_{0,h}^2 + 2|\vec{v}|_{0,h} |\vec{v}|_{1,h},$$

$$(2.11) \quad \max_{1 \leq k \leq J} |\delta^- \vec{v}_k|^2 \leq |\vec{v}|_{1,h}^2 + 2|\vec{v}|_{1,h} |\vec{v}|_{2,h}.$$

In addition, if  $\vec{v}_0 \cdot \vec{e}_1 = \vec{v}_J \cdot \vec{e}_1 = 0$ , then

$$(2.12) \quad |\vec{v}_j \cdot \vec{e}_1| \leq 2q_j(1 - q_j) \max_{1 \leq k \leq J} |\delta^- \vec{v}_k|, \quad 0 \leq j \leq J.$$

*Proof.* The inverse inequality (2.9) follows immediately from the definition (2.8). Let  $0 \leq k \leq J$ . For  $0 \leq j \leq k$  it follows from (2.6b), the elementary inequality

$$(a + b)^2 \leq 2(a^2 + b^2), \quad a, b \in \mathbb{R}$$

and (2.8) that

$$(2.13) \quad |\vec{v}_k|^2 = |\vec{v}_j|^2 + \sum_{\ell=j}^{k-1} (|\vec{v}_{\ell+1}|^2 - |\vec{v}_\ell|^2) = |\vec{v}_j|^2 + h \sum_{\ell=j}^{k-1} (\vec{v}_{\ell+1} + \vec{v}_\ell) \cdot \delta^+ \vec{v}_\ell$$

$$\leq |\vec{v}_j|^2 + \sqrt{2} \left( h \sum_{\ell=j}^{k-1} (|\vec{v}_{\ell+1}|^2 + |\vec{v}_\ell|^2) \right)^{\frac{1}{2}} |\vec{v}|_{1,h} \leq |\vec{v}_j|^2 + 2|\vec{v}|_{0,h} |\vec{v}|_{1,h}.$$

Similarly, for  $k+1 \leq j \leq J$ , we have

$$(2.14) \quad |\vec{v}_k|^2 = |\vec{v}_j|^2 - \sum_{\ell=k}^{j-1} (|\vec{v}_{\ell+1}|^2 - |\vec{v}_\ell|^2) \leq |\vec{v}_j|^2 + 2|\vec{v}|_{0,h} |\vec{v}|_{1,h}.$$

Combining (2.13) and (2.14) yields that  $\max_{0 \leq k \leq J} |\vec{v}_k|^2 \leq |\vec{v}_j|^2 + 2|\vec{v}|_{0,h} |\vec{v}|_{1,h}$ , for  $0 \leq j \leq J$ . Multiplication by  $\frac{h}{2}$  for  $j = 0, J$ , and by  $h$  for  $1 \leq j \leq J-1$ , followed by summation over  $j = 0, \dots, J$ , yields (2.10). The inequality (2.11) is obtained in an analogous manner, taking into account that  $\delta^+ \delta^- \vec{v}_j = \delta^2 \vec{v}_j$ .

In order to prove (2.12), we observe that  $\vec{v}_0 \cdot \vec{e}_1 = \vec{v}_J \cdot \vec{e}_1 = 0$  implies

$$|\vec{v}_j \cdot \vec{e}_1| \leq h \sum_{k=1}^j |\delta^- \vec{v}_k| \leq jh \max_{1 \leq k \leq j} |\delta^- \vec{v}_k| \quad \text{and} \quad |\vec{v}_j \cdot \vec{e}_1| \leq (J-j)h \max_{1 \leq k \leq J} |\delta^- \vec{v}_k|,$$

so that

$$|\vec{v}_j \cdot \vec{e}_1| \leq \min\{q_j, 1 - q_j\} \max_{1 \leq k \leq J} |\delta^- \vec{v}_k| \leq 2q_j(1 - q_j) \max_{1 \leq k \leq J} |\delta^- \vec{v}_k|, \quad 0 \leq j \leq J. \quad \square$$

We consider the following fully discrete approximation, where in order to discretise in time, we let  $t_m = m \Delta t$ ,  $m = 0, \dots, M$ , with the uniform time step  $\Delta t = \frac{T}{M} > 0$ . Let  $\vec{X}_j^0 = \vec{x}_0(q_j)$ ,  $j = 0, \dots, J$ . Then, for  $m = 0, \dots, M - 1$  find  $\vec{X}^m : \mathcal{G}_h \rightarrow \mathbb{R}^2$  such that for  $j = 1, \dots, J - 1$

$$(2.15a) \quad \frac{\vec{X}_j^{m+1} - \vec{X}_j^m}{\Delta t} = \frac{\delta^2 \vec{X}_j^{m+1}}{|\delta^1 \vec{X}_j^m|^2} - \frac{1}{|\delta^1 \vec{X}_j^m|^2} \frac{\delta^1 \vec{X}_j^{m+1} \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} (\delta^1 \vec{X}_j^m)^\perp$$

together with the boundary conditions

$$(2.15b) \quad \vec{X}_0^{m+1} \cdot \vec{e}_1 = 0; \quad \delta^+ \vec{X}_0^{m+1} \cdot \vec{e}_2 = \frac{1}{4} h |\delta^+ \vec{X}_0^m|^2 \frac{\vec{X}_0^{m+1} - \vec{X}_0^m}{\Delta t} \cdot \vec{e}_2,$$

$$(2.15c) \quad \vec{X}_J^{m+1} \cdot \vec{e}_1 = 0; \quad \delta^- \vec{X}_J^{m+1} \cdot \vec{e}_2 = -\frac{1}{4} h |\delta^- \vec{X}_J^m|^2 \frac{\vec{X}_J^{m+1} - \vec{X}_J^m}{\Delta t} \cdot \vec{e}_2.$$

The above scheme requires the solution of a linear system in each time step. We will address the existence and uniqueness of this system in Section 3, within the error analysis. Furthermore, we remark that (2.15b) and (2.15c) are obtained from inserting (1.6a), (1.6b) into a Taylor expansion at  $\rho \in \{0, 1\}$ , yielding a consistency error that is small enough to derive optimal error bounds. At the same time, the form of these conditions turns out to be crucial in order to handle the degeneracy of the equation close to the axis of rotation.

LEMMA 2.3 (Consistency). *Suppose that  $\vec{x} : [0, 1] \times [0, T] \rightarrow \mathbb{R}^2$  satisfies Assumption 2.1. Let  $\vec{x}_j^m := \vec{x}(q_j, t_m)$  for  $j = 0, \dots, J$  and  $m = 0, \dots, M$ . Define the consistency errors of the finite difference scheme (2.15) by*

$$(2.16a) \quad \vec{R}_j^{m+1} := \frac{\vec{x}_j^{m+1} - \vec{x}_j^m}{\Delta t} - \frac{\delta^2 \vec{x}_j^{m+1}}{|\delta^1 \vec{x}_j^m|^2} + \frac{1}{|\delta^1 \vec{x}_j^m|^2} \frac{\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2}{\vec{x}_j^m \cdot \vec{e}_1} (\delta^1 \vec{x}_j^m)^\perp, \quad 1 \leq j \leq J - 1,$$

as well as

$$(2.16b) \quad \vec{R}_0^{m+1} := \delta^+ \vec{x}_0^{m+1} \cdot \vec{e}_2 - \frac{1}{4} h |\delta^+ \vec{x}_0^m|^2 \frac{\vec{x}_0^{m+1} - \vec{x}_0^m}{\Delta t} \cdot \vec{e}_2,$$

$$(2.16c) \quad \vec{R}_J^{m+1} := \delta^- \vec{x}_J^{m+1} \cdot \vec{e}_2 + \frac{1}{4} h |\delta^- \vec{x}_J^m|^2 \frac{\vec{x}_J^{m+1} - \vec{x}_J^m}{\Delta t} \cdot \vec{e}_2.$$

Then there exists a constant  $C > 0$  such that, for  $m = 0, \dots, M - 1$ ,

$$(2.17) \quad |\vec{R}_j^{m+1}| \leq C (h^2 + \Delta t), \quad j = 1, \dots, J - 1, \quad \text{and} \quad |\vec{R}_0^{m+1}| + |\vec{R}_J^{m+1}| \leq Ch(h^2 + \Delta t).$$

*Proof.* Simple Taylor expansions yield the well-known results

$$(2.18a) \quad \left| \frac{\bar{x}_j^{m+1} - \bar{x}_j^m}{\Delta t} - \bar{x}_t(q_j, t_m) \right| \leq C\Delta t, \quad 0 \leq j \leq J, \quad 0 \leq m \leq M-1,$$

$$(2.18b) \quad |\delta^- \bar{x}_j^m - \bar{x}_\rho(q_j, t_m)| \leq Ch, \quad 1 \leq j \leq J, \quad 0 \leq m \leq M,$$

$$(2.18c)$$

$$|\delta^1 \bar{x}_j^m - \bar{x}_\rho(q_j, t_m)| + |\delta^2 \bar{x}_j^m - \bar{x}_{\rho\rho}(q_j, t_m)| \leq Ch^2, \quad 1 \leq j \leq J-1, \quad 0 \leq m \leq M,$$

$$(2.18d)$$

$$\delta^1 \bar{x}_j^m - \bar{x}_\rho(q_j, t_m) - \frac{1}{6}h^2 \bar{x}_{\rho\rho\rho}(q_j, t_m) = \mathcal{O}(h^3), \quad 1 \leq j \leq J-1, \quad 0 \leq m \leq M,$$

where we have observed that  $\bar{x}(q_j, \cdot) \in C^2([0, T])$  and  $\bar{x}(\cdot, t_m) \in C^4([0, 1])$ . Evaluating (1.6a) at  $(\rho, t) = (q_j, t_m)$ ,  $j = 1, \dots, J-1$ ,  $m = 0, \dots, M-1$ , we find that

$$(2.19) \quad \bar{x}_t(q_j, t_m) = \frac{\bar{x}_{\rho\rho}(q_j, t_m)}{|\bar{x}_\rho(q_j, t_m)|^2} - \frac{1}{|\bar{x}_\rho(q_j, t_m)|^2} \frac{\bar{x}_\rho(q_j, t_m) \cdot \bar{e}_2}{\bar{x}_j^m \cdot \bar{e}_1} \bar{x}_\rho^\perp(q_j, t_m),$$

where the assumed regularity of  $\bar{x}$  allows us to use (1.6a) also at time  $t = 0$ . If we combine (2.16a) with (2.19), and note (2.18a) as well as (2.18c), we obtain

$$(2.20) \quad \begin{aligned} |\bar{R}_j^{m+1}| &\leq \left| \frac{\bar{x}_j^{m+1} - \bar{x}_j^m}{\Delta t} - \bar{x}_t(q_j, t_m) \right| + \left| \frac{\delta^2 \bar{x}_j^{m+1}}{|\delta^1 \bar{x}_j^m|^2} - \frac{\bar{x}_{\rho\rho}(q_j, t_m)}{|\bar{x}_\rho(q_j, t_m)|^2} \right| \\ &\quad + \frac{|\bar{x}_\rho(q_j, t_m) \cdot \bar{e}_2|}{\bar{x}_j^m \cdot \bar{e}_1} \left| \frac{(\delta^1 \bar{x}_j^m)^\perp}{|\delta^1 \bar{x}_j^m|^2} - \frac{\bar{x}_\rho^\perp(q_j, t_m)}{|\bar{x}_\rho(q_j, t_m)|^2} \right| \\ &\quad + \frac{1}{|\delta^1 \bar{x}_j^m|} \frac{|(\delta^1 \bar{x}_j^{m+1} - \bar{x}_\rho(q_j, t_m)) \cdot \bar{e}_2|}{\bar{x}_j^m \cdot \bar{e}_1} \\ &\leq C(h^2 + \Delta t) + C \frac{|(\delta^1 \bar{x}_j^{m+1} - \bar{x}_\rho(q_j, t_m)) \cdot \bar{e}_2|}{\bar{x}_j^m \cdot \bar{e}_1}. \end{aligned}$$

In addition, it follows from (2.18d), (A.1b), (A.9) and (A.8) that

$$\begin{aligned} &|(\delta^1 \bar{x}_j^{m+1} - \bar{x}_\rho(q_j, t_m)) \cdot \bar{e}_2| \\ &\leq |(\delta^1 \bar{x}_j^{m+1} - \delta^1 \bar{x}_j^m) \cdot \bar{e}_2| + |(\delta^1 \bar{x}_j^m - \bar{x}_\rho(q_j, t_m)) \cdot \bar{e}_2| \\ &\leq \Delta t \sup_{t_m \leq t \leq t_{m+1}} |\delta^1 \bar{x}_t(q_j, t) \cdot \bar{e}_2| + Ch^2 \min_{q \in \{0,1\}} |\bar{x}_{\rho\rho\rho}(q_j, t_m) \cdot \bar{e}_2 - \bar{x}_{\rho\rho\rho}(q, t_m) \cdot \bar{e}_2| + Ch^3 \\ &\leq K\Delta t \sup_{t_m \leq t \leq t_{m+1}} |\bar{x}(q_j, t) \cdot \bar{e}_1| + Ch^2 \min_{q \in \{0,1\}} |\bar{x}_{\rho\rho\rho}(q_j, t_m) - \bar{x}_{\rho\rho\rho}(q, t_m)| + Ch^3 \\ &\leq C\Delta tq_j(1-q_j) + Ch^2 \min_{q \in \{0,1\}} |q_j - q| + Ch^3 \leq Cq_j(1-q_j)(\Delta t + h^2). \end{aligned}$$

If we insert this bound into (2.20) and note that  $\bar{x}_j^m \cdot \bar{e}_1 \geq c_2 q_j(1-q_j)$ ,  $0 \leq j \leq J$ , in view of (A.8), we obtain (2.17) for  $\bar{R}_j^{m+1}$ ,  $j = 1, \dots, J-1$ . Let us next examine  $R_0^{m+1}$ . A Taylor expansion yields

$$\delta^+ \bar{x}_0^{m+1} = \frac{\bar{x}_1^{m+1} - \bar{x}_0^{m+1}}{h} = \bar{x}_\rho(0, t_{m+1}) + \frac{1}{2}h \bar{x}_{\rho\rho}(0, t_{m+1}) + \frac{1}{6}h^2 \bar{x}_{\rho\rho\rho}(0, t_{m+1}) + \mathcal{O}(h^3),$$

which together with (1.6b), (A.1b),  $\vec{x}_{\rho\rho}(0, \cdot) \in C^1([0, T])$  and (A.1c) implies that

$$\begin{aligned} \delta^+ \vec{x}_0^{m+1} \cdot \vec{e}_2 &= \frac{1}{2} h \vec{x}_{\rho\rho}(0, t_{m+1}) \cdot \vec{e}_2 + \mathcal{O}(h^3) = \frac{1}{2} h \vec{x}_{\rho\rho}(0, t_m) \cdot \vec{e}_2 + \mathcal{O}(h(h^2 + \Delta t)) \\ &= \frac{1}{4} h (\vec{x}_t(0, t_m) \cdot \vec{e}_2) |\vec{x}_\rho(0, t_m)|^2 + \mathcal{O}(h(h^2 + \Delta t)) \\ &= \frac{1}{4} h \frac{\vec{x}_0^{m+1} - \vec{x}_0^m}{\Delta t} \cdot \vec{e}_2 |\delta^+ \vec{x}_0^m|^2 + \mathcal{O}(h(h^2 + \Delta t)), \end{aligned}$$

where we have used (2.18a) as well as

$$\begin{aligned} |\delta^+ \vec{x}_0^m|^2 - |\vec{x}_\rho(0, t_m)|^2 &= (\delta^+ \vec{x}_0^m - \vec{x}_\rho(0, t_m)) \cdot (\delta^+ \vec{x}_0^m + \vec{x}_\rho(0, t_m)) \\ &= \left( \frac{1}{2} h \vec{x}_{\rho\rho}(0, t_m) + \mathcal{O}(h^2) \right) \cdot (2\vec{x}_\rho(0, t_m) + \mathcal{O}(h)) \\ &= h \vec{x}_{\rho\rho}(0, t_m) \cdot \vec{x}_\rho(0, t_m) + \mathcal{O}(h^2) = \mathcal{O}(h^2), \end{aligned}$$

recall (A.1a). The bound for  $R_J^{m+1}$  is obtained in a similar way.  $\square$

**THEOREM 2.4.** *Suppose that  $\vec{x} : [0, 1] \times [0, T] \rightarrow \mathbb{R}^2$  satisfies Assumption 2.1. Then there exist  $h_0 > 0, \gamma > 0$  such that the discrete solution  $(\vec{X}^m)_{m=1, \dots, M}$  to (2.15) exists, and the error*

$$(2.21) \quad \vec{E}_j^m := \vec{x}_j^m - \vec{X}_j^m, \quad j = 0, \dots, J; \quad m = 0, \dots, M$$

satisfies:

$$(2.22) \quad \max_{1 \leq m \leq M} \left[ \|\vec{E}^m\|_{1,h}^2 + \max_{0 \leq j \leq J} |\vec{E}_j^m|^2 \right] \leq C(h^4 + (\Delta t)^2),$$

$$(2.23) \quad \Delta t \sum_{m=1}^M \left[ \|\vec{E}^m\|_{2,h}^2 + \left| \frac{\vec{E}^m - \vec{E}^{m-1}}{\Delta t} \right|_{0,h}^2 \right] \leq C(h^4 + (\Delta t)^2),$$

provided that  $0 < h \leq h_0$  and  $\Delta t \leq \gamma h$ .

**3. Proof of Theorem 2.4.** Assumption 2.1 assures the existence of positive constants  $c_0, C_0, c_1, \delta$  such that (2.2) and (2.3) hold. Let  $h \leq \delta$ . We set  $J_1 := \lfloor \frac{\delta}{h} \rfloor \in \mathbb{Z}_{\geq 1}$ , so that  $q_{J_1} = J_1 h \in [\frac{1}{2}\delta, \delta]$ . We shall prove Theorem 2.4 with the help of an induction argument. In particular, we will prove that there exist  $h_0 > 0, 0 < \gamma \leq 1$  and  $\mu > 0$  such that if  $0 < h \leq h_0$  and  $\Delta t \leq \gamma h$ , then for  $m \in \{0, \dots, M\}$  the discrete solution  $\vec{X}^m$  exists and satisfies

$$(3.1) \quad \|\vec{E}^m\|_{1,h}^2 \leq (h^4 + (\Delta t)^2) e^{\mu t_m}.$$

The assertion (3.1) clearly holds for  $m = 0$ , for arbitrary  $h_0 \leq \delta, 0 < \gamma \leq 1$  and  $\mu > 0$ . On assuming that (3.1) holds for a fixed  $m \in \{0, \dots, M-1\}$ , we will now show that it also holds for  $m+1$ .

To begin, let us choose  $0 < h_0 \leq \delta$  and  $0 < \gamma \leq 1$  so small that

$$(h_0^2 + \gamma^2) e^{\mu T} \leq 1.$$

Then, since  $\Delta t \leq \gamma h$ , (3.1) implies that

$$\|\vec{E}^m\|_{1,h}^2 \leq h^2 (h^2 + \gamma^2) e^{\mu t_m} \leq h^2, \quad 0 < h \leq h_0.$$

In particular, we infer from Lemma 2.2 that

$$(3.2) \quad \max_{0 \leq j \leq J} |\vec{E}_j^m| + \max_{1 \leq j \leq J} |\delta^- \vec{E}_j^m| + \max_{1 \leq j \leq J-1} |\delta^1 \vec{E}_j^m| \leq Ch^{\frac{1}{2}}.$$



This implies for  $1 \leq j \leq J$ , on recalling (2.18b) and (2.2), that

$$|\delta^- \vec{X}_j^m| \leq |\delta^- \vec{x}_j^m| + |\delta^- \vec{E}_j^m| \leq |\vec{x}_\rho(q_j, t_m)| + Ch^{\frac{1}{2}} \leq C_0 + Ch^{\frac{1}{2}},$$

and similarly  $|\delta^- \vec{X}_j^m| \geq c_0 - Ch^{\frac{1}{2}}$ . Arguing in the same way for  $\delta^1 \vec{X}_j^m$ , we infer that

$$(3.3) \quad \frac{1}{2}c_0 \leq |\delta^- \vec{X}_j^m| \leq 2C_0, \quad 1 \leq j \leq J; \quad \frac{1}{2}c_0 \leq |\delta^1 \vec{X}_j^m| \leq 2C_0, \quad 1 \leq j \leq J-1,$$

provided that  $0 < h \leq h_0$  and  $h_0 > 0$  is chosen smaller if necessary. A similar argument together with (2.3a), (2.3b) shows that

$$(3.4) \quad \delta^- \vec{X}_j^m \cdot \vec{e}_1 \geq \frac{1}{4}c_0, \quad 1 \leq j \leq J_1; \quad \delta^- \vec{X}_j^m \cdot \vec{e}_1 \leq -\frac{1}{4}c_0, \quad J - J_1 \leq j \leq J.$$

Next, since  $\vec{x}_\rho(0, t_m) \cdot \vec{e}_2 = 0$ , recall (1.6b), we have from (3.2) and (2.15b) that

$$\begin{aligned} h|\delta^- \vec{X}_1^m| &\leq h|\delta^- \vec{x}_1^m| + h|\delta^- \vec{E}_1^m| \leq h\delta^- \vec{x}_1^m \cdot \vec{e}_1 + h|\delta^- \vec{x}_1^m \cdot \vec{e}_2| + Ch^{\frac{3}{2}} \\ &\leq h\delta^- \vec{X}_1^m \cdot \vec{e}_1 + Ch^{\frac{3}{2}} = \vec{X}_1^m \cdot \vec{e}_1 + Ch^{\frac{3}{2}} \leq (1 + Ch^{\frac{1}{2}})\vec{x}_1^m \cdot \vec{e}_1 + Ch^{\frac{3}{2}}, \end{aligned}$$

where in the last step we have observed that

$$\vec{X}_1^m \cdot \vec{e}_1 - \vec{x}_1^m \cdot \vec{e}_1 = -h\delta^- \vec{E}_1^m \cdot \vec{e}_1 \leq Ch^{\frac{3}{2}} \leq Ch^{\frac{1}{2}}\vec{x}_1^m \cdot \vec{e}_1,$$

on noting  $\vec{x}_1^m \cdot \vec{e}_1 \geq \frac{1}{2}c_0h$ . Arguing in the same way at the right boundary, we obtain

$$(3.5) \quad \frac{3}{4}h|\delta^- \vec{X}_J^m| \leq \vec{X}_J^m \cdot \vec{e}_1 \leq \frac{4}{3}\vec{x}_J^m \cdot \vec{e}_1; \quad \frac{3}{4}h|\delta^- \vec{X}_J^m| \leq \vec{X}_{J-1}^m \cdot \vec{e}_1 \leq \frac{4}{3}\vec{x}_{J-1}^m \cdot \vec{e}_1,$$

for a possibly smaller  $h_0 > 0$ . Next, (3.2), (2.3c) and the fact that  $J_1h \geq \frac{1}{2}\delta$  imply that

$$(3.6) \quad \vec{X}_j^m \cdot \vec{e}_1 \geq c_1 - Ch^{\frac{1}{2}} \geq \frac{1}{2}c_1, \quad J_1 \leq j \leq J - J_1,$$

after choosing  $h_0$  again smaller if required. In addition, there exists  $c_3 > 0$  such that

$$(3.7) \quad \vec{X}_j^m \cdot \vec{e}_1 \geq c_3q_j(1 - q_j), \quad 0 \leq j \leq J.$$

To see this, note that  $\vec{X}_0^m \cdot \vec{e}_1 = 0$  and (3.4) imply that

$$(3.8a) \quad \vec{X}_j^m \cdot \vec{e}_1 \geq \frac{1}{4}c_0jh \geq \frac{1}{4}c_0q_j(1 - q_j), \quad 0 \leq j \leq J_1,$$

and similarly

$$(3.8b) \quad \vec{X}_j^m \cdot \vec{e}_1 \geq \frac{1}{4}c_0q_j(1 - q_j), \quad J - J_1 \leq j \leq J.$$

Combining these estimates with (3.6) proves the bound (3.7). If we combine (3.7) with (2.12) and (3.3), we obtain

$$(3.9) \quad \frac{\vec{X}_{j\pm 1}^m \cdot \vec{e}_1}{\vec{X}_j^m \cdot \vec{e}_1} \leq \frac{4C_0}{c_3} \frac{q_{j\pm 1}(1 - q_{j\pm 1})}{q_j(1 - q_j)} \leq \frac{8C_0}{c_3}, \quad 1 \leq j \leq J-1.$$

Finally, (2.2) and (3.3) imply that

$$(3.10a) \quad \left| \frac{1}{|\delta^- \vec{x}_j^m|^2} - \frac{1}{|\delta^- \vec{X}_j^m|^2} \right| \leq C|\delta^- \vec{E}_j^m|, \quad 1 \leq j \leq J,$$

$$(3.10b) \quad \left| \frac{1}{|\delta^1 \vec{x}_j^m|^2} - \frac{1}{|\delta^1 \vec{X}_j^m|^2} \right| \leq C|\delta^1 \vec{E}_j^m|, \quad 1 \leq j \leq J-1.$$

LEMMA 3.1 (Existence and uniqueness). *Let  $\vec{X}^m : \mathcal{G}_h \rightarrow \mathbb{R}^2$  be as above. Then (2.15) has a unique solution  $\vec{X}^{m+1} : \mathcal{G}_h \rightarrow \mathbb{R}^2$ , provided that  $h_0$  is small enough.*

*Proof.* The relations (2.15) form a linear system with  $2(J+1)$  unknowns for the  $2(J+1)$  values of  $\vec{X}^{m+1}$  at the nodes  $q_j$ ,  $j = 0, \dots, J$ . It is therefore sufficient to show that the corresponding homogeneous system

$$(3.11a) \quad \frac{1}{\Delta t} \vec{X}_j - \frac{1}{|\delta^1 \vec{X}_j^m|^2} \delta^2 \vec{X}_j = -\frac{1}{|\delta^1 \vec{X}_j^m|^2} \frac{\delta^1 \vec{X}_j \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} (\delta^1 \vec{X}_j^m)^\perp, \quad j = 1, \dots, J-1;$$

$$(3.11b) \quad \vec{X}_0 \cdot \vec{e}_1 = 0; \quad \delta^+ \vec{X}_0 \cdot \vec{e}_2 = \frac{1}{4} \frac{h}{\Delta t} (\vec{X}_0 \cdot \vec{e}_2) |\delta^+ \vec{X}_0^m|^2;$$

$$(3.11c) \quad \vec{X}_J \cdot \vec{e}_1 = 0; \quad \delta^- \vec{X}_J \cdot \vec{e}_2 = -\frac{1}{4} \frac{h}{\Delta t} (\vec{X}_J \cdot \vec{e}_2) |\delta^- \vec{X}_J^m|^2$$

only has the trivial solution  $\vec{X} = \vec{0}$ . If we multiply (3.11a) with  $-h\delta^2 \vec{X}_j$  and sum from  $j = 1, \dots, J-1$  we obtain with the help of (2.7) that

$$\begin{aligned} & \frac{1}{\Delta t} |\vec{X}|_{1,h}^2 + \frac{1}{\Delta t} (\vec{X}_0 \cdot \delta^+ \vec{X}_0 - \vec{X}_J \cdot \delta^- \vec{X}_J) + h \sum_{j=1}^{J-1} \frac{1}{|\delta^1 \vec{X}_j^m|^2} |\delta^2 \vec{X}_j|^2 \\ &= h \sum_{j=1}^{J-1} \frac{1}{|\delta^1 \vec{X}_j^m|^2} \frac{\delta^1 \vec{X}_j \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} (\delta^1 \vec{X}_j^m)^\perp \cdot \delta^2 \vec{X}_j. \end{aligned}$$

In view of (3.11b) and (3.5) we have

$$\frac{1}{\Delta t} \vec{X}_0 \cdot \delta^+ \vec{X}_0 = \frac{1}{\Delta t} (\vec{X}_0 \cdot \vec{e}_2) (\delta^+ \vec{X}_0 \cdot \vec{e}_2) = \frac{4}{h} \frac{(\delta^+ \vec{X}_0 \cdot \vec{e}_2)^2}{|\delta^+ \vec{X}_0^m|^2} \geq \frac{9}{4} h \frac{(\delta^- \vec{X}_1 \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2}.$$

Using a similar argument at the right end point, as well as (3.3), we deduce

$$\begin{aligned} (3.12) \quad & \frac{1}{\Delta t} |\vec{X}|_{1,h}^2 + \frac{1}{4C_0^2} |\vec{X}|_{2,h}^2 + \frac{9}{4} h \left( \frac{(\delta^- \vec{X}_1 \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^+ \vec{X}_{J-1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right) \\ & \leq h \sum_{j=1}^{J-1} \frac{\delta^1 \vec{X}_j \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} \left( \frac{(\delta^1 \vec{X}_j^m)^\perp}{|\delta^1 \vec{X}_j^m|^2} - \frac{(\delta^1 \vec{x}_j^m)^\perp}{|\delta^1 \vec{x}_j^m|^2} \right) \cdot \delta^2 \vec{X}_j \\ & \quad + h \sum_{j=1}^{J-1} \frac{1}{|\delta^1 \vec{x}_j^m|^2} \frac{\delta^1 \vec{X}_j \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} (\delta^1 \vec{x}_j^m)^\perp \cdot \delta^2 \vec{X}_j \\ & =: h \sum_{j=1}^{J-1} \vec{S}_j^1 \cdot \delta^2 \vec{X}_j + h \sum_{j=1}^{J-1} \vec{S}_j^2 \cdot \delta^2 \vec{X}_j. \end{aligned}$$

Using (3.10b) and (3.2) we infer that

$$|\vec{S}_j^1| \leq C \frac{|\delta^1 \vec{X}_j \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1} |\delta^1 \vec{E}_j^m| |\delta^2 \vec{X}_j| \leq Ch^{\frac{1}{2}} \frac{|\delta^1 \vec{X}_j \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1} |\delta^2 \vec{X}_j|$$

and hence

$$h \sum_{j=1}^{J-1} \vec{S}_j^1 \cdot \delta^2 \vec{X}_j \leq \frac{1}{8C_0^2} |\vec{X}|_{2,h}^2 + Ch^2 \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{X}_j \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2}.$$

The term  $\vec{S}_j^2$  corresponds exactly to  $-\vec{T}_j^{m,3}$  in (3.13) below, if we replace  $\vec{E}^{m+1}$  by  $\vec{X}$ . We may therefore deduce from Lemma 3.4 that

$$\begin{aligned} h \sum_{j=1}^{J-1} \vec{S}_j^2 \cdot \delta^2 \vec{X}_j &\leq -c_4 h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{X}_j \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} + \frac{5}{3} h \left( \frac{(\delta^- \vec{X}_1 \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^+ \vec{X}_{J-1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right) \\ &\quad + \frac{1}{8C_0^2} |\vec{X}|_{2,h}^2 + C |\vec{X}|_{1,h}^2. \end{aligned}$$

If we insert the above bounds into (3.12) and recall that  $\Delta t \leq \gamma h \leq h$  we infer that

$$\left( \frac{1}{h} - C \right) |\vec{X}|_{1,h}^2 + (c_4 - Ch) h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{X}_j \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} \leq 0,$$

which implies that  $\vec{X} \equiv \vec{X}_0$  provided that  $0 < h \leq h_0$ , where  $h_0$  is chosen smaller if necessary. The boundary conditions (3.11b), on noting (3.3), then yield  $\vec{X} \equiv 0$ .  $\square$

We begin our error analysis by combining (2.21), (2.15a) and (2.16a), in order to derive the following error relation:

$$\begin{aligned} (3.13) \quad \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \frac{\delta^2 \vec{E}_j^{m+1}}{|\delta^1 \vec{X}_j^m|^2} &= \left( \frac{1}{|\delta^1 \vec{x}_j^m|^2} - \frac{1}{|\delta^1 \vec{X}_j^m|^2} \right) \delta^2 \vec{x}_j^{m+1} \\ &\quad + \frac{\delta^1 \vec{X}_j^{m+1} \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} \left[ \left( \frac{1}{|\delta^1 \vec{X}_j^m|^2} - \frac{1}{|\delta^1 \vec{x}_j^m|^2} \right) (\delta^1 \vec{X}_j^m)^\perp - \frac{1}{|\delta^1 \vec{x}_j^m|^2} (\delta^1 \vec{E}_j^m)^\perp \right] \\ &\quad - \frac{1}{|\delta^1 \vec{x}_j^m|^2} \frac{\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} (\delta^1 \vec{x}_j^m)^\perp \\ &\quad + \frac{1}{|\delta^1 \vec{x}_j^m|^2} \left( \frac{1}{\vec{X}_j^m \cdot \vec{e}_1} - \frac{1}{\vec{x}_j^m \cdot \vec{e}_1} \right) (\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2) (\delta^1 \vec{x}_j^m)^\perp + \vec{R}_j^{m+1} \\ &=: \sum_{i=1}^5 \vec{T}_j^{m,i}, \quad 1 \leq j \leq J-1. \end{aligned}$$

Furthermore, for the boundary points we have in view of (1.6b), (2.15b), (2.15c), (2.16b) and (2.16c) that

$$(3.14a) \quad (\vec{E}_0^{m+1} - \vec{E}_0^m) \cdot \vec{e}_1 = (\vec{E}_J^{m+1} - \vec{E}_J^m) \cdot \vec{e}_1 = 0,$$

$$(3.14b) \quad \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} = \frac{4 \delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2 - R_0^{m+1}}{|\delta^+ \vec{X}_0^m|^2} \vec{e}_2 + \left( 1 - \frac{|\delta^+ \vec{x}_0^m|^2}{|\delta^+ \vec{X}_0^m|^2} \right) \frac{\vec{x}_0^{m+1} - \vec{x}_0^m}{\Delta t},$$

$$(3.14c) \quad \frac{\vec{E}_J^{m+1} - \vec{E}_J^m}{\Delta t} = -\frac{4 \delta^- \vec{E}_J^{m+1} \cdot \vec{e}_2 - R_J^{m+1}}{|\delta^- \vec{X}_J^m|^2} \vec{e}_2 + \left( 1 - \frac{|\delta^- \vec{x}_J^m|^2}{|\delta^- \vec{X}_J^m|^2} \right) \frac{\vec{x}_J^{m+1} - \vec{x}_J^m}{\Delta t}.$$

Our strategy for the proof of (3.1) with  $m$  replaced by  $m+1$  is now as follows. In a discrete analogue to the formal procedure in (2.4), we are going to multiply (3.13) with a second order difference of the error  $\vec{E}^{m+1}$ . The ensuing analysis is technical, and so we split it into three steps. In a first step, we control the terms generated on the left hand side of (3.13), in order to obtain Lemma 3.2. Next we estimate four of the five terms generated by the right hand side of (3.13), see Lemma 3.3. The remaining

term, which is generated by  $\vec{T}^{m,3}$  and loosely corresponds to the last integral in (2.4), requires a particularly careful analysis. We present the derived estimate in Lemma 3.4, where in the proof we will mimic the formal calculations from (2.5).

The induction step is then completed by combining the three lemmas.

LEMMA 3.2. *There exists  $C_1 > 0$  such that for all  $0 < \lambda \leq 1$*

(3.15)

$$\begin{aligned} & \frac{1+\lambda}{2\Delta t} (|\vec{E}^{m+1}|_{1,h}^2 - |\vec{E}^m|_{1,h}^2) + \frac{1}{2\Delta t} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}^2 + \frac{1}{4C_0^2} |\vec{E}^{m+1}|_{2,h}^2 \\ & + \frac{1}{4} c_0^2 \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + (2 - C_1 \lambda) h \left[ \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^+ \vec{E}_{J-1}^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right] \\ & \leq Ch(h^4 + (\Delta t)^2) + C|\vec{E}^m|_{1,h}^2 \\ & + h \sum_{i=1}^5 \sum_{j=1}^{J-1} \vec{T}_j^{m,i} \cdot (\lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1}). \end{aligned}$$

*Proof.* Fix  $0 < \lambda \leq 1$ . If we multiply (3.13) by  $h(\lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1})$  and sum over  $j = 1, \dots, J-1$ , we obtain

$$\begin{aligned} (3.16) \quad & - (1 + \lambda) \frac{h}{\Delta t} \sum_{j=1}^{J-1} (\vec{E}_j^{m+1} - \vec{E}_j^m) \cdot \delta^2 \vec{E}_j^{m+1} + h \sum_{j=1}^{J-1} \frac{|\delta^2 \vec{E}_j^{m+1}|^2}{|\delta^1 \vec{X}_j^m|^2} \\ & + h \lambda \sum_{j=1}^{J-1} |\delta^1 \vec{X}_j^m|^2 \left| \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} \right|^2 \\ & = h \sum_{i=1}^5 \sum_{j=1}^{J-1} \vec{T}_j^{m,i} \cdot (\lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1}). \end{aligned}$$

Applying summation by parts, (2.7), to the first term in (3.16), and noting (2.8) and  $2(a-b)a = a^2 - b^2 + (a-b)^2$ , yields

$$\begin{aligned} (3.17) \quad & - \frac{h}{\Delta t} \sum_{j=1}^{J-1} (\vec{E}_j^{m+1} - \vec{E}_j^m) \cdot \delta^2 \vec{E}_j^{m+1} \\ & = \frac{h}{\Delta t} \sum_{j=1}^J \delta^- (\vec{E}_j^{m+1} - \vec{E}_j^m) \cdot \delta^- \vec{E}_j^{m+1} \\ & \quad - \frac{\vec{E}_J^{m+1} - \vec{E}_J^m}{\Delta t} \cdot \delta^- \vec{E}_J^{m+1} + \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} \cdot \delta^+ \vec{E}_0^{m+1} \\ & = \frac{1}{2\Delta t} (|\vec{E}^{m+1}|_{1,h}^2 - |\vec{E}^m|_{1,h}^2) + \frac{1}{2\Delta t} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}^2 \\ & \quad - \frac{\vec{E}_J^{m+1} - \vec{E}_J^m}{\Delta t} \cdot \delta^- \vec{E}_J^{m+1} + \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} \cdot \delta^+ \vec{E}_0^{m+1}. \end{aligned}$$

On noting (3.14b), (2.6b), Young's inequality, (2.17), (3.3), (3.10a) and (3.5), we can

estimate the last term on the right hand side of (3.17) as

$$\begin{aligned}
 \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} \cdot \delta^+ \vec{E}_0^{m+1} &= \frac{4}{h} \frac{(\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2)^2}{|\delta^+ \vec{X}_0^m|^2} - \frac{4}{h} \frac{\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2}{|\delta^+ \vec{X}_0^m|^2} R_0^{m+1} \\
 &\quad + \left(1 - \frac{|\delta^+ \vec{x}_0^m|^2}{|\delta^+ \vec{X}_0^m|^2}\right) \frac{\vec{x}_0^{m+1} - \vec{x}_0^m}{\Delta t} \cdot \vec{e}_2 (\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2) \\
 &\geq \frac{4 - \varepsilon}{h} \frac{(\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2)^2}{|\delta^+ \vec{X}_1^m|^2} - C_\varepsilon h (h^4 + (\Delta t)^2) - C_\varepsilon h |\delta^+ \vec{E}_0^m|^2 \\
 &\geq \frac{9}{16} (4 - \varepsilon) h \frac{(\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} - C_\varepsilon h (h^4 + (\Delta t)^2) - C_\varepsilon h |\delta^+ \vec{E}_0^m|^2.
 \end{aligned}$$

On choosing  $\varepsilon$  sufficiently small, and arguing similarly for  $\frac{\vec{E}_J^{m+1} - \vec{E}_J^m}{\Delta t} \cdot \delta^- \vec{E}_J^{m+1}$ , we find that (3.17) implies

$$\begin{aligned}
 (3.18) \quad &-(1 + \lambda) \frac{h}{\Delta t} \sum_{j=1}^{J-1} (\vec{E}_j^{m+1} - \vec{E}_j^m) \cdot \delta^2 \vec{E}_j^{m+1} \geq \frac{1 + \lambda}{2\Delta t} (|\vec{E}^{m+1}|_{1,h}^2 - |\vec{E}^m|_{1,h}^2) \\
 &+ \frac{1}{2\Delta t} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}^2 + 2h \left[ \frac{(\delta^+ \vec{E}_0^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^- \vec{E}_J^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right] \\
 &- Ch (|\delta^- \vec{E}_1^m|^2 + |\delta^- \vec{E}_J^m|^2) - Ch (h^4 + (\Delta t)^2).
 \end{aligned}$$

In addition, we deduce from (3.14b), (3.10a), (3.3) and the fact that thanks to (3.7) we have  $h|\delta^+ \vec{X}_0^m|^2 \geq h^{-1} (\vec{X}_1^m \cdot \vec{e}_1)^2 \geq C \vec{X}_1^m \cdot \vec{e}_1$  that

$$\left| \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} \right| \leq C \frac{|\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2|}{\vec{X}_1^m \cdot \vec{e}_1} + C \frac{1}{h} |R_0^{m+1}| + C |\delta^- \vec{E}_1^m|,$$

so that (2.17) yields

$$(3.19) \quad \frac{h}{2} \left| \frac{\vec{E}_0^{m+1} - \vec{E}_0^m}{\Delta t} \right|^2 \leq Ch \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + Ch (h^2 + \Delta t)^2 + Ch |\delta^- \vec{E}_1^m|^2.$$

Inserting (3.18) into (3.16) and using (2.8), (3.3), (2.6b), as well as (3.19) and a corresponding estimate at the right boundary, we obtain the desired result (3.15).  $\square$

LEMMA 3.3. *Let  $0 < \lambda \leq 1$  and  $\Delta t \leq \gamma h$ . Then*

$$\begin{aligned}
 (3.20) \quad &h \sum_{i=1}^5 \sum_{j=1}^{J-1} \vec{T}_j^{m,i} \cdot (\lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1}) + h \sum_{j=1}^{J-1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1} \\
 &\leq \frac{1}{8C_0^2} |\vec{E}^{m+1}|_{2,h}^2 + C (|\vec{E}^m|_{1,h}^2 + |\vec{E}^{m+1}|_{1,h}^2) + C \frac{\gamma}{\Delta t} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}^2 \\
 &+ \frac{1}{8} c_0^2 \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + C (h^4 + (\Delta t)^2) + C (\lambda + h) h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2}.
 \end{aligned}$$

*Proof.* We recall the definitions of the terms  $\vec{T}^{m,i}$  in (3.13). Then we note from (3.10b), (3.3), (3.7), (A.8) and (3.2) that

$$\begin{aligned} |\vec{T}_j^{m,1}| + |\vec{T}_j^{m,2}| &\leq C(|\delta^2 \vec{x}_j^{m+1}| + \frac{|\delta^1 \vec{X}_j^{m+1} \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1}) |\delta^1 \vec{E}_j^m| \\ &\leq C(1 + \frac{|\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2|}{\vec{x}_j^{m+1} \cdot \vec{e}_1} \frac{|\vec{x}_j^{m+1} \cdot \vec{e}_1|}{\vec{X}_j^m \cdot \vec{e}_1} + \frac{|\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1}) |\delta^1 \vec{E}_j^m| \\ &\leq C|\delta^1 \vec{E}_j^m| + Ch^{\frac{1}{2}} \frac{|\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1}, \quad 1 \leq j \leq J-1, \end{aligned}$$

so that (2.6c), (2.6b), (3.3) and (2.8) imply that

$$\begin{aligned} (3.21) \quad &h \sum_{i=1}^2 \sum_{j=1}^{J-1} \vec{T}_j^{m,i} \cdot (\lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1}) \\ &\leq Ch \sum_{j=1}^{J-1} \left( |\delta^- \vec{E}_j^m| + |\delta^- \vec{E}_{j+1}^m| + h^{\frac{1}{2}} \frac{|\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1} \right) \left( \lambda \left| \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} \right| + |\delta^2 \vec{E}_j^{m+1}| \right) \\ &\leq \varepsilon \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + \varepsilon |\vec{E}^{m+1}|_{2,h}^2 + C_\varepsilon |\vec{E}^m|_{1,h}^2 + C_\varepsilon h^2 \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2}. \end{aligned}$$

The term involving the product of  $\vec{T}^{m,3}$  with  $\delta^2 \vec{E}^{m+1}$  is not estimated, while

$$\begin{aligned} (3.22) \quad &h \sum_{j=1}^{J-1} \vec{T}_j^{m,3} \cdot \lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} \\ &\leq C\lambda \left( h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} \right)^{\frac{1}{2}} \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h} \\ &\leq \varepsilon \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + C_\varepsilon \lambda h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2}. \end{aligned}$$

Next, we have

$$(3.23) \quad |\vec{T}_j^{m,4}| \leq C \frac{|\vec{E}_j^m \cdot \vec{e}_1|}{(\vec{x}_j^m \cdot \vec{e}_1)(\vec{X}_j^m \cdot \vec{e}_1)} |\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2|, \quad 1 \leq j \leq J-1.$$

In addition, (A.8) and (A.9) yield

$$|\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2| = \frac{|\delta^1 \vec{x}_j^{m+1} \cdot \vec{e}_2|}{\vec{x}_j^{m+1} \cdot \vec{e}_1} \vec{x}_j^{m+1} \cdot \vec{e}_1 \leq Cq_j(1-q_j), \quad 1 \leq j \leq J-1,$$

so that (3.23), (2.12), (A.8), (3.7) and (2.9) imply

$$\begin{aligned}
 |\vec{T}_j^{m,4}| &\leq C \frac{q_j^2(1-q_j)^2}{c_2c_3q_j^2(1-q_j)^2} \max_{1 \leq k \leq J} |\delta^- \vec{E}_k^m| \leq C \max_{1 \leq k \leq J} |\delta^- \vec{E}_k^m| \\
 &\leq C \left( \max_{1 \leq k \leq J} |\delta^- \vec{E}_k^{m+1}| + \max_{1 \leq k \leq J} |\delta^- (\vec{E}_k^{m+1} - \vec{E}_k^m)| \right) \\
 &\leq C \max_{1 \leq k \leq J} |\delta^- \vec{E}_k^{m+1}| + Ch^{-\frac{1}{2}} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}, \quad 1 \leq j \leq J-1.
 \end{aligned}$$

Hence we obtain with the help of (2.11) and the fact that  $\Delta t \leq \gamma h$

(3.24)

$$\begin{aligned}
 &h \sum_{j=1}^{J-1} \vec{T}_j^{m,4} \cdot \left( \lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1} \right) \\
 &\leq C \left( \max_{1 \leq j \leq J} |\delta^- \vec{E}_j^{m+1}| + h^{-\frac{1}{2}} |\vec{E}^{m+1} - \vec{E}^m|_{1,h} \right) \left( \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h} + |\vec{E}^{m+1}|_{2,h} \right) \\
 &\leq \varepsilon |\vec{E}^{m+1}|_{2,h}^2 + \varepsilon \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + C_\varepsilon |\vec{E}^{m+1}|_{1,h}^2 + C_\varepsilon \frac{\gamma}{\Delta t} |\vec{E}^{m+1} - \vec{E}^m|_{1,h}^2.
 \end{aligned}$$

Finally, we infer from (2.17) that

$$\begin{aligned}
 (3.25) \quad &h \sum_{j=1}^{J-1} \vec{T}_j^{m,5} \cdot \left( \lambda |\delta^1 \vec{X}_j^m|^2 \frac{\vec{E}_j^{m+1} - \vec{E}_j^m}{\Delta t} - \delta^2 \vec{E}_j^{m+1} \right) \\
 &\leq C(h^2 + \Delta t) \left( \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h} + |\vec{E}^{m+1}|_{2,h} \right) \\
 &\leq \varepsilon |\vec{E}^{m+1}|_{2,h}^2 + \varepsilon \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + C_\varepsilon (h^4 + (\Delta t)^2).
 \end{aligned}$$

If we add (3.21), (3.22), (3.24) and (3.25), and then choose  $\varepsilon$  sufficiently small, the bound (3.20) follows.  $\square$

In the next lemma we mimic the formal calculations in (2.5), thereby closing our estimates.

LEMMA 3.4. *There exists a constant  $c_4 > 0$  such that*

$$\begin{aligned}
 (3.26) \quad &h \sum_{j=1}^{J-1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1} \\
 &\geq c_4 h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} - \frac{5}{3} h \left( \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^+ \vec{E}_{J-1}^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right) \\
 &\quad - \varepsilon |\vec{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\vec{E}^{m+1}|_{1,h}^2.
 \end{aligned}$$

*Proof.* Let us start by writing

(3.27)

$$\sum_{j=1}^{J-1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1} = \sum_{j=1}^{J_1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1} + \sum_{j=J_1+1}^{J-J_1-1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1} + \sum_{j=J-J_1}^{J-1} \vec{T}_j^{m,3} \cdot \delta^2 \vec{E}_j^{m+1},$$

and we begin by estimating the first sum on the right hand side of (3.27). On recalling (3.13), we can write

$$(3.28) \quad \vec{T}_j^{m,3} = \frac{1}{|\delta^1 \vec{x}_j^m|} \frac{\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} \vec{e}_2 - \frac{1}{|\delta^1 \vec{x}_j^m|} \frac{\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2}{\vec{X}_j^m \cdot \vec{e}_1} \left( \frac{(\delta^1 \vec{x}_j^m)^\perp}{|\delta^1 \vec{x}_j^m|} + \vec{e}_2 \right) =: \vec{S}_j^1 + \vec{S}_j^2.$$

Observing from (2.6) that

$$\begin{aligned} (\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2) \delta^2 \vec{E}_j^{m+1} \cdot \vec{e}_2 &= \frac{1}{h} (\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2) (\delta^+ \vec{E}_j^{m+1} - \delta^- \vec{E}_j^{m+1}) \cdot \vec{e}_2 \\ &= \frac{1}{2h} ((\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2 - (\delta^- \vec{E}_j^{m+1} \cdot \vec{e}_2)^2), \end{aligned}$$

we find that

$$(3.29) \quad \begin{aligned} h \sum_{j=1}^{J_1} \vec{S}_j^1 \cdot \delta^2 \vec{E}_j^{m+1} &= \frac{1}{2} \sum_{j=1}^{J_1} \frac{1}{|\delta^1 \vec{x}_j^m|} \frac{1}{\vec{X}_j^m \cdot \vec{e}_1} ((\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2 - (\delta^- \vec{E}_j^{m+1} \cdot \vec{e}_2)^2) \\ &= \frac{1}{2} \sum_{j=1}^{J_1} \frac{(\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2}{|\delta^1 \vec{x}_j^m| \vec{X}_j^m \cdot \vec{e}_1} - \frac{1}{2} \sum_{j=0}^{J_1-1} \frac{(\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2}{|\delta^1 \vec{x}_{j+1}^m| \vec{X}_{j+1}^m \cdot \vec{e}_1} \\ &= \frac{1}{2} \sum_{j=1}^{J_1-1} \left( \frac{1}{|\delta^1 \vec{x}_j^m|} \frac{1}{\vec{X}_j^m \cdot \vec{e}_1} - \frac{1}{|\delta^1 \vec{x}_{j+1}^m|} \frac{1}{\vec{X}_{j+1}^m \cdot \vec{e}_1} \right) (\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2 \\ &\quad - \frac{1}{2} \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{|\delta^1 \vec{x}_1^m| \vec{X}_1^m \cdot \vec{e}_1} + \frac{1}{2} \frac{(\delta^- \vec{E}_{J_1+1}^{m+1} \cdot \vec{e}_2)^2}{|\delta^1 \vec{x}_{J_1}^m| \vec{X}_{J_1}^m \cdot \vec{e}_1}. \end{aligned}$$

In order to estimate the sum on the right hand side, we observe that  $|\delta^1 \vec{x}_{j+1}^m| \leq \frac{1}{2h} \int_{q_j}^{q_{j+2}} |\vec{x}_\rho^m| d\rho \leq C_0$ , recall (2.2), and  $\vec{X}_j^m \cdot \vec{e}_1 \leq \vec{X}_{j+1}^m \cdot \vec{e}_1 \leq C \vec{X}_j^m \cdot \vec{e}_1$  for  $1 \leq j \leq J_1 - 1$ , recall (3.4) and (3.9). Hence we obtain with the help of (3.4) that

$$\begin{aligned} &\frac{1}{|\delta^1 \vec{x}_j^m|} \frac{1}{\vec{X}_j^m \cdot \vec{e}_1} - \frac{1}{|\delta^1 \vec{x}_{j+1}^m|} \frac{1}{\vec{X}_{j+1}^m \cdot \vec{e}_1} \\ &= h \frac{1}{|\delta^1 \vec{x}_{j+1}^m|} \frac{\delta^- \vec{X}_{j+1}^m \cdot \vec{e}_1}{(\vec{X}_{j+1}^m \cdot \vec{e}_1) \vec{X}_j^m \cdot \vec{e}_1} - \frac{1}{\vec{X}_j^m \cdot \vec{e}_1} \left( \frac{1}{|\delta^1 \vec{x}_{j+1}^m|} - \frac{1}{|\delta^1 \vec{x}_j^m|} \right) \\ &\geq h \frac{1}{C_0} \frac{\frac{c_0}{4}}{(\vec{X}_{j+1}^m \cdot \vec{e}_1)^2} - Ch \frac{1}{\vec{X}_{j+1}^m \cdot \vec{e}_1} \geq h \frac{c_0}{8C_0} \frac{1}{(\vec{X}_{j+1}^m \cdot \vec{e}_1)^2} - Ch, \quad 1 \leq j \leq J_1 - 1. \end{aligned}$$

Furthermore, on noting  $|\delta^1 \vec{x}_1^m| \geq \delta^1 \vec{x}_1^m \cdot \vec{e}_1 = \frac{1}{2h} \vec{x}_2^m \cdot \vec{e}_1 \geq \frac{1}{2h} \vec{x}_1^m \cdot \vec{e}_1 \geq \frac{3}{8} \frac{1}{h} \vec{X}_1^m \cdot \vec{e}_1$ , recall (3.5), we have that

$$\frac{1}{2} \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{|\delta^1 \vec{x}_1^m| \vec{X}_1^m \cdot \vec{e}_1} \leq \frac{4}{3} h \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2}.$$

Inserting the above two estimates into (3.29) yields, on recalling (2.8), that

$$(3.30) \quad \begin{aligned} h \sum_{j=1}^{J_1} \vec{S}_j^1 \cdot \delta^2 \vec{E}_j^{m+1} &\geq h \frac{c_0}{16C_0} \sum_{j=1}^{J_1-1} \frac{(\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_{j+1}^m \cdot \vec{e}_1)^2} - \frac{4}{3} h \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} \\ &\quad - C |\vec{E}^{m+1}|_{1,h}^2. \end{aligned}$$



Note that in view of (1.6b), (2.2) and (2.3a) we have  $\frac{\bar{x}_\rho(0,t)}{|\bar{x}_\rho(0,t)|} = \bar{e}_1$ , so that  $\frac{\bar{x}_\rho^\perp(0,t)}{|\bar{x}_\rho(0,t)|} = \bar{e}_1^\perp = -\bar{e}_2$ . Hence (2.18c) and the smoothness of  $\bar{x}$  imply

$$\begin{aligned} \left| \frac{(\delta^1 \bar{x}_j^m)^\perp}{|\delta^1 \bar{x}_j^m|} + \bar{e}_2 \right| &\leq \left| \frac{(\delta^1 \bar{x}_j^m)^\perp}{|\delta^1 \bar{x}_j^m|} - \frac{\bar{x}_\rho^\perp(q_j, t_m)}{|\bar{x}_\rho(q_j, t_m)|} \right| + \left| \frac{\bar{x}_\rho^\perp(q_j, t_m)}{|\bar{x}_\rho(q_j, t_m)|} - \frac{\bar{x}_\rho^\perp(0, t_m)}{|\bar{x}_\rho(0, t_m)|} \right| \\ &\leq C(h + q_j) \leq Cq_j, \end{aligned}$$

for  $1 \leq j \leq J_1$ , which means that with the help of (3.8a) we obtain

$$(3.31) \quad \begin{aligned} h \sum_{j=1}^{J_1} \bar{S}_j^2 \cdot \delta^2 \bar{E}_j^{m+1} &\geq -Ch \sum_{j=1}^{J_1} (|\delta^- \bar{E}_j^{m+1}| + |\delta^+ \bar{E}_j^{m+1}|) |\delta^2 \bar{E}_j^{m+1}| \\ &\geq -\varepsilon |\bar{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\bar{E}^{m+1}|_{1,h}^2. \end{aligned}$$

Combining (3.30) and (3.31) with (3.28) we obtain

$$(3.32) \quad \begin{aligned} h \sum_{j=1}^{J_1} \bar{T}_j^{m,3} \cdot \delta^2 \bar{E}_j^{m+1} &\geq h \frac{c_0}{16C_0} \sum_{j=2}^{J_1} \frac{(\delta^- \bar{E}_j^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_j^m \cdot \bar{e}_1)^2} - \frac{4}{3} h \frac{(\delta^- \bar{E}_1^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_1^m \cdot \bar{e}_1)^2} \\ &\quad - \varepsilon |\bar{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\bar{E}^{m+1}|_{1,h}^2. \end{aligned}$$

In order to estimate the third sum on the right hand side of (3.27), we start from

$$\bar{T}_j^{m,3} = -\frac{1}{|\delta^1 \bar{x}_j^m|} \frac{\delta^1 \bar{E}_j^{m+1} \cdot \bar{e}_2}{\bar{X}_j^m \cdot \bar{e}_1} \bar{e}_2 - \frac{1}{|\delta^1 \bar{x}_j^m|} \frac{\delta^1 \bar{E}_j^{m+1} \cdot \bar{e}_2}{\bar{X}_j^m \cdot \bar{e}_1} \left( \frac{(\delta^1 \bar{x}_j^m)^\perp}{|\delta^1 \bar{x}_j^m|} - \bar{e}_2 \right),$$

and use similar arguments as above to obtain

$$(3.33) \quad \begin{aligned} h \sum_{j=J-J_1}^{J-1} \bar{T}_j^{m,3} \cdot \delta^2 \bar{E}_j^{m+1} &\geq h \frac{c_0}{16C_0} \sum_{j=J-J_1+1}^{J-1} \frac{(\delta^- \bar{E}_j^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_j^m \cdot \bar{e}_1)^2} - \frac{4}{3} h \frac{(\delta^- \bar{E}_J^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_{J-1}^m \cdot \bar{e}_1)^2} \\ &\quad - \varepsilon |\bar{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\bar{E}^{m+1}|_{1,h}^2. \end{aligned}$$

Moreover, it follows from (2.2), (3.3) and (3.6), that

$$(3.34) \quad \begin{aligned} h \sum_{j=J_1+1}^{J-J_1-1} \bar{T}_j^{m,3} \cdot \delta^2 \bar{E}_j^{m+1} &- h \frac{c_0}{16C_0} \sum_{j=J_1+1}^{J-J_1} \frac{(\delta^- \bar{E}_j^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_j^m \cdot \bar{e}_1)^2} \\ &\geq -Ch \sum_{j=1}^{J-1} (|\delta^- \bar{E}_j^{m+1}| + |\delta^- \bar{E}_{j+1}^{m+1}|) |\delta^2 \bar{E}_j^{m+1}| - Ch \sum_{j=1}^{J-1} |\delta^- \bar{E}_j^{m+1}|^2 \\ &\geq -\varepsilon |\bar{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\bar{E}^{m+1}|_{1,h}^2. \end{aligned}$$

If we combine (3.32), (3.33) and (3.34) we obtain

$$(3.35) \quad \begin{aligned} h \sum_{j=1}^{J-1} \bar{T}_j^{m,3} \cdot \delta^2 \bar{E}_j^{m+1} &\geq h \frac{c_0}{16C_0} \sum_{j=2}^{J-1} \frac{(\delta^- \bar{E}_j^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_j^m \cdot \bar{e}_1)^2} \\ &\quad - \frac{4}{3} h \left( \frac{(\delta^- \bar{E}_1^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_1^m \cdot \bar{e}_1)^2} + \frac{(\delta^+ \bar{E}_{J-1}^{m+1} \cdot \bar{e}_2)^2}{(\bar{X}_{J-1}^m \cdot \bar{e}_1)^2} \right) - \varepsilon |\bar{E}^{m+1}|_{2,h}^2 - C_\varepsilon |\bar{E}^{m+1}|_{1,h}^2. \end{aligned}$$

Observing that in view of (2.6b) and (3.9)

$$\frac{|\delta^+ \vec{E}_j^{m+1} \cdot \vec{e}_2|}{\vec{X}_j^m \cdot \vec{e}_1} \leq \frac{8C_0}{c_3} \frac{|\delta^- \vec{E}_{j+1}^{m+1} \cdot \vec{e}_2|}{\vec{X}_{j+1}^m \cdot \vec{e}_1}, \quad 1 \leq j \leq J-2,$$

we have

$$\begin{aligned} h \sum_{j=1}^{J-1} \frac{(\delta^1 \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} &\leq \frac{h}{2} \sum_{j=1}^{J-1} \frac{(\delta^+ \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} + \frac{h}{2} \sum_{j=1}^{J-1} \frac{(\delta^- \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2} \\ &\leq \frac{h}{2} \left( \frac{(\delta^- \vec{E}_1^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_1^m \cdot \vec{e}_1)^2} + \frac{(\delta^+ \vec{E}_{J-1}^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_{J-1}^m \cdot \vec{e}_1)^2} \right) + Ch \sum_{j=2}^{J-1} \frac{(\delta^- \vec{E}_j^{m+1} \cdot \vec{e}_2)^2}{(\vec{X}_j^m \cdot \vec{e}_1)^2}. \end{aligned}$$

If we insert this bound into (3.35), we deduce (3.26) provided that  $c_4$  is small enough.  $\square$

Combining Lemmas 3.2, 3.3 and 3.4 we obtain after choosing  $\varepsilon$ ,  $\gamma$  and  $\lambda$  sufficiently small

$$(3.36) \quad \frac{1+\lambda}{2\Delta t} (|\vec{E}^{m+1}|_{1,h}^2 - |\vec{E}^m|_{1,h}^2) + \frac{1}{16C_0^2} |\vec{E}^{m+1}|_{2,h}^2 + \frac{1}{8} c_0^2 \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 \\ \leq C (|\vec{E}^m|_{1,h}^2 + |\vec{E}^{m+1}|_{1,h}^2) + C(h^4 + (\Delta t)^2).$$

Furthermore, we have

$$(3.37) \quad \frac{1}{2\Delta t} (|\vec{E}^{m+1}|_{0,h}^2 - |\vec{E}^m|_{0,h}^2) \leq \frac{1}{2} \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h} (|\vec{E}^{m+1}|_{0,h} + |\vec{E}^m|_{0,h}) \\ \leq \frac{1}{16} c_0^2 \lambda \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 + C (|\vec{E}^{m+1}|_{0,h}^2 + |\vec{E}^m|_{0,h}^2).$$

On inserting (3.37) into (3.36), divided by  $(1+\lambda)$ , we obtain that there exist constants  $c_6 > 0$  and  $C_2 > 0$  such that

$$(3.38) \quad \frac{1}{\Delta t} (\|\vec{E}^{m+1}\|_{1,h}^2 - \|\vec{E}^m\|_{1,h}^2) + c_6 \left( |\vec{E}^{m+1}|_{2,h}^2 + \left| \frac{\vec{E}^{m+1} - \vec{E}^m}{\Delta t} \right|_{0,h}^2 \right) \\ \leq C_2 (\|\vec{E}^{m+1}\|_{1,h}^2 + \|\vec{E}^m\|_{1,h}^2) + C_2 (h^4 + (\Delta t)^2).$$

Combining (3.38) with the induction hypothesis (3.1) completes the proof of Theorem 2.4. In fact, if we choose  $h_0$  so small that  $C_2 \Delta t \leq \frac{1}{2}$  for  $\Delta t \leq \gamma h_0$ , then  $0 < (1 - C_2 \Delta t)^{-1} \leq 1 + 2C_2 \Delta t$ , and so it follows from (3.38) and (3.1) that

$$\begin{aligned} \|\vec{E}^{m+1}\|_{1,h}^2 &\leq (1 - C_2 \Delta t)^{-1} \left[ (1 + C_2 \Delta t) \|\vec{E}^m\|_{1,h}^2 + C_2 \Delta t (h^4 + (\Delta t)^2) \right] \\ &\leq (1 + 2C_2 \Delta t)^2 \|\vec{E}^m\|_{1,h}^2 + C_2 (1 + 2C_2 \Delta t) \Delta t (h^4 + (\Delta t)^2) \\ &\leq (1 + 2C_2 \Delta t)^2 (h^4 + (\Delta t)^2) e^{\mu t_m} + 2C_2 \Delta t (h^4 + (\Delta t)^2) \\ &\leq (1 + 3C_2 \Delta t)^2 (h^4 + (\Delta t)^2) e^{\mu t_m} \\ &\leq (h^4 + (\Delta t)^2) e^{6C_2 \Delta t} e^{\mu t_m} = (h^4 + (\Delta t)^2) e^{\mu t_{m+1}}, \end{aligned}$$

TABLE 1

Errors for the convergence test for (4.1) over the time interval  $[0, 0.125]$  with  $\Delta t = h$ .

$J$	$\max_{m=0,\dots,M}  \vec{x}^m - \vec{X}^m _{0,h}$	EOC	$\max_{m=0,\dots,M}  \vec{x}^m - \vec{X}^m _{1,h}$	EOC
32	3.5744e-02	—	1.1225e-01	—
64	2.0034e-02	0.84	6.2934e-02	0.83
128	1.0690e-02	0.91	3.3582e-02	0.91
256	5.5352e-03	0.95	1.7389e-02	0.95
512	2.8185e-03	0.97	8.8546e-03	0.97

TABLE 2

Errors for the convergence test for (4.1) over the time interval  $[0, 0.125]$  with  $\Delta t = h^2$ .

$J$	$\max_{m=0,\dots,M}  \vec{x}^m - \vec{X}^m _{0,h}$	EOC	$\max_{m=0,\dots,M}  \vec{x}^m - \vec{X}^m _{1,h}$	EOC
32	1.0024e-03	—	3.1480e-03	—
64	2.5201e-04	1.99	7.9165e-04	1.99
128	6.3093e-05	2.00	1.9821e-04	2.00
256	1.5779e-05	2.00	4.9571e-05	2.00
512	3.9451e-06	2.00	1.2394e-05	2.00

if we choose  $\mu = 6C_2$ . Since  $\mu$ , as well as  $\gamma$ , were chosen independently of  $h$  and  $\Delta t$ , we have shown (3.1) by induction. Together with (2.10) this proves the inequality (2.22). Finally, multiplying (3.38) by  $\Delta t$  and summing for  $m = 0, \dots, M - 1$  yields the bound (2.23).

**4. Numerical results.** It is easy to show that a shrinking sphere with radius  $[1 - 4t]^{\frac{1}{2}}$  is a solution to (1.1). In fact, the parameterization

$$(4.1) \quad \vec{x}(\rho, t) = [1 - 4t]^{\frac{1}{2}} \begin{pmatrix} \sin(\pi \rho) \\ \cos(\pi \rho) \end{pmatrix}$$

solves (1.6). On letting  $\vec{x}_j^m = \vec{x}(q_j, t_m)$ ,  $j = 0, \dots, J$ , we compare (4.1) to the discrete solutions  $(\vec{X}^m)_{m=0,\dots,M}$  of (2.15) and perform two convergence experiments. In particular, we choose either  $\Delta t = h$  or  $\Delta t = h^2$ , for  $h = J^{-1} = 2^{-k}$ ,  $k = 5, \dots, 9$ .

The results in Tables 1 and 2 confirm the theoretical results proved in Theorem 2.4. We stress that the quadratic convergence rate for the  $H^1$ -seminorm in Table 2 is better than the linear rate observed in [2, Table 4] for the finite element scheme considered there. This suggests that the delicate treatment of the boundary nodes in our finite difference scheme (2.15) is crucial to obtain the optimal convergence rate in Theorem 2.4.

In Figure 1 we show a simulation for mean curvature flow of a sphere with an inscribed torus. In particular, the initial surface selfintersects on the equator of the sphere, and has genus 0. For the scheme (2.15) we choose  $J = 1024$  and  $\Delta t = 10^{-4}$ . Under mean curvature flow, the torus attempts to shrink to a circle. For the generating curve, this means that the cusp or swallow tail tries to disappear. Of course, for the approximated partial differential equation this represents a singularity, where the curvatures of the curve, and of the corresponding axisymmetric surface, blow up. However, the discrete scheme (2.15) is blind to the self-intersection and the associated singularity. Hence the finite difference approximation simply integrates across the singularity. The same behavior can be seen, for example, in [8, Figure 4.2] and [3,




FIG. 1. Evolution for a torus inscribed within a sphere. Plots are at times  $t = 0, 0.1, 0.14, 0.2$ .

Figure 6]. Continuing the evolution in Figure 1 would show the curve approaching a shrinking semicircle, that eventually vanishes at the origin.

In the recent article [2], the present authors numerically studied the Angenent torus, see [1, 17], as an example of a self-shrinker for mean curvature flow. Here we recall that the surface  $\mathcal{S}(0)$  is called a self-shrinker, if the self-similar family of surfaces

$$\mathcal{S}(t) = [1 - t]^{\frac{1}{2}} \mathcal{S}(0)$$

is a solution to (1.1). In what follows, we would like to use our approximation (2.15) in order to investigate self-shrinkers of genus-0. It was shown in [15] that the only bounded embedded genus-0 self-shrinker in  $\mathbb{R}^3$  is the sphere of radius 2. Note that the unit sphere has an extinction time of  $T_0 = \frac{1}{4}$ , recall (4.1). On the other hand, in [9] the existence of infinitely many immersed self-shrinkers with rotational symmetry was proved. Hence, inspired by [9, Figure 3], we would like compute such a self-similar evolution for mean curvature flow. To this end, we use the open curve analogue of [2, (5.7), (5.8)] in order to calculate a profile curve of a self-shrinker that has three self-intersections. Using the obtained curve as initial data for the scheme (2.15) yields the self-similar evolution displayed in Figure 2. Here we used the discretization parameters  $J = 512$  and  $\Delta t = 10^{-4}$ . Note that the numerical method appears to confirm the unit extinction time. In fact, continuing the evolution until the method breaks down yields the behaviour of the approximate surface area

$$A^m = 2\pi h \sum_{j=1}^J \vec{X}_j^m \cdot \vec{e}_1 |\delta^- \vec{X}_j^m|$$

as shown in Figure 3, with the expected linear decay and an approximate extinction time of 1.

Finally, we include a numerical experiment to demonstrate that our scheme can also deal with initial data that violate the  $90^\circ$  contact angle condition in (1.6b). To this end, in Figure 4 we start a simulation for a surface that has two cone singularities: an inward cone and an outward cone. The generating curve has a  $45^\circ$  contact angle at the axis of rotation, which induces a discontinuous jump in time for the solution of the partial differential equation. For the simulation we choose  $J = 512$  and  $\Delta t = 10^{-4}$  for the scheme (2.15). It can be observed that the outward cone very quickly smoothens to a rounded tip, while the inward cone also smoothens and rises at the same time. Eventually the curve approaches a shrinking semicircle, that will shrink to a point.




FIG. 2. Self-similar evolution for a surface with three self-intersections. Plots are at times  $t = 0, 0.1, \dots, 0.9$ , and again at times  $t = 0$  and  $t = 0.9$ .




FIG. 3. A plot of the approximate surface area  $A^m$ , for the simulation in Figure 2, over time.

### Appendix A. Properties of the solution.

LEMMA A.1 (Behaviour at the boundary). *Let  $\vec{x} : [0, 1] \times [0, T] \rightarrow \mathbb{R}^2$  satisfy Assumption 2.1. Then we have*

$$(A.1a) \quad \vec{x}_{\rho\rho} \cdot \vec{e}_1 = \vec{x}_{\rho\rho} \cdot \vec{x}_\rho = 0 \quad \text{on } \{0, 1\} \times [0, T],$$

$$(A.1b) \quad \vec{x}_{\rho\rho\rho} \cdot \vec{e}_2 = 0 \quad \text{on } \{0, 1\} \times [0, T],$$

$$(A.1c) \quad \vec{x}_t = 2 \frac{\vec{x}_{\rho\rho} \cdot \vec{e}_2}{|\vec{x}_\rho|^2} \vec{e}_2 \quad \text{on } \{0, 1\} \times [0, T].$$

*Proof.* We have from (1.6b) that

$$(A.2) \quad \vec{x}_\rho^\perp(0, t) = (|\vec{x}_\rho(0, t)| \vec{e}_1)^\perp = -|\vec{x}_\rho(0, t)| \vec{e}_2,$$

and so we obtain with the help of L'Hospital's rule that

$$(A.3) \quad \lim_{\rho \searrow 0} \frac{1}{|\vec{x}_\rho(\rho, t)|^2} \frac{\vec{x}_\rho(\rho, t) \cdot \vec{e}_2}{\vec{x}(\rho, t) \cdot \vec{e}_1} \vec{x}_\rho^\perp(\rho, t) = \frac{1}{|\vec{x}_\rho(0, t)|^2} \frac{\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_2}{\vec{x}_\rho(0, t) \cdot \vec{e}_1} \vec{x}_\rho^\perp(0, t) \\ = -\frac{\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_2}{|\vec{x}_\rho(0, t)|^2} \vec{e}_2.$$

Thus (1.6a) implies that

$$(A.4) \quad \vec{x}_t(0, t) = \frac{\vec{x}_{\rho\rho}(0, t)}{|\vec{x}_\rho(0, t)|^2} + \frac{\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_2}{|\vec{x}_\rho(0, t)|^2} \vec{e}_2.$$

Observing from (1.6b) that  $\vec{x}_t(0, t) \cdot \vec{e}_1 = 0$ , we infer from (A.4) that  $\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_1 = 0$ , which together with (1.6b) proves (A.1a) at  $\rho = 0$ . In particular,  $\vec{x}_{\rho\rho}(0, t) = (\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_2) \vec{e}_2$ .




FIG. 4. Evolution for a surface with two cone singularities. Plots are at times  $t = 0, 0.01, 0.1, 0.4$ .

$\vec{e}_2 \vec{e}_2$ . Combining this with (A.4) yields (A.1c) at  $\rho = 0$ . In order to prove (A.1b), we differentiate (1.6a) with respect to  $\rho$  and obtain

$$(A.5) \quad \vec{x}_{t\rho} = \frac{\vec{x}_{\rho\rho\rho}}{|\vec{x}_\rho|^2} - 2 \frac{\vec{x}_{\rho\rho} \cdot \vec{x}_\rho}{|\vec{x}_\rho|^4} \vec{x}_{\rho\rho} + \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \left( 2 \frac{\vec{x}_{\rho\rho} \cdot \vec{x}_\rho}{|\vec{x}_\rho|^4} \vec{x}_\rho^\perp - \frac{1}{|\vec{x}_\rho|^2} \vec{x}_{\rho\rho} \right) - \left( \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \right)_\rho \frac{1}{|\vec{x}_\rho|^2} \vec{x}_\rho^\perp$$

in  $(0, 1) \times (0, T]$ . A further application of L'Hospital's rule implies that

$$(A.6) \quad \lim_{\rho \searrow 0} \left( \frac{\vec{x}_\rho \cdot \vec{e}_2}{\vec{x} \cdot \vec{e}_1} \right)_\rho (\rho, t) = \lim_{\rho \searrow 0} \frac{(\vec{x}_{\rho\rho}(\rho, t) \cdot \vec{e}_2)(\vec{x}(\rho, t) \cdot \vec{e}_1) - (\vec{x}_\rho(\rho, t) \cdot \vec{e}_2)(\vec{x}_\rho(\rho, t) \cdot \vec{e}_1)}{(\vec{x}(\rho, t) \cdot \vec{e}_1)^2} \\ = \lim_{\rho \searrow 0} \frac{(\vec{x}_{\rho\rho\rho}(\rho, t) \cdot \vec{e}_2)(\vec{x}(\rho, t) \cdot \vec{e}_1) - (\vec{x}_\rho(\rho, t) \cdot \vec{e}_2)(\vec{x}_{\rho\rho}(\rho, t) \cdot \vec{e}_1)}{2(\vec{x}(\rho, t) \cdot \vec{e}_1)(\vec{x}_\rho(\rho, t) \cdot \vec{e}_1)} \\ = \frac{1}{2} \frac{\vec{x}_{\rho\rho\rho}(0, t) \cdot \vec{e}_2}{\vec{x}_\rho(0, t) \cdot \vec{e}_1},$$

since  $\vec{x}_{\rho\rho}(0, t) \cdot \vec{e}_1 = 0$ . Combining (A.5) and (A.6), on noting (A.1a), (1.6b) and (A.2), yields that

$$(A.7) \quad \vec{x}_{t\rho}(0, t) = \frac{\vec{x}_{\rho\rho\rho}(0, t)}{|\vec{x}_\rho(0, t)|^2} + \frac{1}{2} \frac{\vec{x}_{\rho\rho\rho}(0, t) \cdot \vec{e}_2}{|\vec{x}_\rho(0, t)|^2} \vec{e}_2.$$

Since  $\vec{x}_{t\rho}(0, t) \cdot \vec{e}_2 = 0$  in view of (1.6b), we deduce from (A.7) that also (A.1b) holds at the left boundary point. The proof of (A.1) for the other boundary point is analogous.  $\square$

LEMMA A.2. *Let  $\vec{x} : [0, 1] \times [0, T] \rightarrow \mathbb{R}^2$  satisfy Assumption 2.1. Then there exists  $0 < c_2 < \tilde{c}_2$  such that*

$$(A.8) \quad c_2 \rho(1 - \rho) \leq \vec{x}(\rho, t) \cdot \vec{e}_1 \leq \tilde{c}_2 \rho(1 - \rho) \quad \text{for all } (\rho, t) \in [0, 1] \times [0, T].$$

Moreover, there exists  $K > 0$  such that for all  $0 < h \leq \frac{1}{2}\delta$ , with  $\delta$  as in (2.3),

$$(A.9) \quad \frac{1}{\vec{x} \cdot \vec{e}_1} \left| \frac{\partial_t^\ell \vec{x}(\cdot + h, t) - \partial_t^\ell \vec{x}(\cdot - h, t)}{2h} \cdot \vec{e}_2 \right| \leq K \quad \text{in } [h, 1 - h] \times [0, T], \quad \ell = 0, 1.$$

*Proof.* The result (A.8) is an immediate consequence of (2.3).

Let  $t \in [0, T]$  and  $h \leq \rho \leq \frac{1}{2}\delta$ . We infer from (1.6b) and (2.3a) that  $\vec{x}(\rho, t) \cdot \vec{e}_1 \geq \frac{1}{2}c_0\rho$  and hence

$$\begin{aligned} \frac{1}{\vec{x}(\rho, t) \cdot \vec{e}_1} \left| \frac{\vec{x}(\rho + h, t) - \vec{x}(\rho - h, t)}{2h} \cdot \vec{e}_2 \right| &\leq \frac{2}{c_0\rho} \left| \frac{1}{2h} \int_{\rho-h}^{\rho+h} (\vec{x}_\rho(\zeta, t) - \vec{x}_\rho(0, t)) \cdot \vec{e}_2 \, d\zeta \right| \\ &\leq \frac{2}{c_0} \max_{[0, \delta]} |\vec{x}_{\rho\rho}(\cdot, t) \cdot \vec{e}_2| \\ &\leq \frac{2}{c_0} \max_{[0, \delta] \times [0, T]} |\vec{x}_{\rho\rho} \cdot \vec{e}_2|. \end{aligned}$$

We can argue in the same way for  $1 - \frac{1}{2}\delta \leq \rho \leq 1 - h$ , while for  $\frac{1}{2}\delta \leq \rho \leq 1 - \frac{1}{2}\delta$  we have that

$$\begin{aligned} \frac{1}{\vec{x}(\rho, t) \cdot \vec{e}_1} \left| \frac{\vec{x}(\rho + h, t) - \vec{x}(\rho - h, t)}{2h} \cdot \vec{e}_2 \right| &\leq \frac{1}{c_1} \left| \frac{1}{2h} \int_{\rho-h}^{\rho+h} \vec{x}_\rho(\zeta, t) \cdot \vec{e}_2 \, d\zeta \right| \\ &\leq \frac{1}{c_1} \max_{[\frac{\delta}{2}, 1 - \frac{\delta}{2}] \times [0, T]} |\vec{x}_\rho \cdot \vec{e}_2|, \end{aligned}$$

so that (A.9) holds with  $K = \max\{\frac{2}{c_0} \max_{[0, \delta] \times [0, T]} |\vec{x}_{\rho\rho} \cdot \vec{e}_2|, \frac{1}{c_1} \max_{[\frac{\delta}{2}, 1 - \frac{\delta}{2}] \times [0, T]} |\vec{x}_\rho \cdot \vec{e}_2|\}$  in the case  $\ell = 0$ . The case  $\ell = 1$  can be treated in the same way, on noting that  $\vec{x}_{t\rho}(q, t) \cdot \vec{e}_2 = 0$  for  $q \in \{0, 1\}$ .  $\square$

## REFERENCES

- [1] S. B. ANGENENT, *Shrinking doughnuts*, in Nonlinear diffusion equations and their equilibrium states, 3 (Gregynog, 1989), vol. 7 of Progr. Nonlinear Differential Equations Appl., Birkhäuser Boston, Boston, MA, 1992, pp. 21–38.
- [2] J. W. BARRETT, K. DECKELNICK, AND R. NÜRNBERG, *A finite element error analysis for axisymmetric mean curvature flow*, IMA J. Numer. Anal., (2020). (to appear).
- [3] J. W. BARRETT, H. GARCKE, AND R. NÜRNBERG, *On the variational approximation of combined second and fourth order geometric evolution equations*, SIAM J. Sci. Comput., 29 (2007), pp. 1006–1041.
- [4] ———, *On the parametric finite element approximation of evolving hypersurfaces in  $\mathbb{R}^3$* , J. Comput. Phys., 227 (2008), pp. 4281–4307.
- [5] ———, *Variational discretization of axisymmetric curvature flows*, Numer. Math., 141 (2019), pp. 791–837.
- [6] ———, *Parametric finite element approximations of curvature driven interface evolutions*, in Handb. Numer. Anal., A. Bonito and R. H. Nochetto, eds., vol. 21, Elsevier, Amsterdam, 2020, pp. 275–423.
- [7] K. DECKELNICK AND G. DZIUK, *On the approximation of the curve shortening flow*, in Calculus of Variations, Applications and Computations (Pont-à-Mousson, 1994), C. Bandle, J. Bemelmans, M. Chipot, J. S. J. Paulin, and I. Shafrir, eds., vol. 326 of Pitman Res. Notes Math. Ser., Longman Sci. Tech., Harlow, 1995, pp. 100–108.
- [8] K. DECKELNICK, G. DZIUK, AND C. M. ELLIOTT, *Computation of geometric partial differential equations and mean curvature flow*, Acta Numer., 14 (2005), pp. 139–232.
- [9] G. DRUGAN AND S. J. KLEENE, *Immersed self-shrinkers*, Trans. Amer. Math. Soc., 369 (2017), pp. 7213–7250.
- [10] G. DZIUK, *An algorithm for evolutionary surfaces*, Numer. Math., 58 (1991), pp. 603–611.
- [11] ———, *Convergence of a semi-discrete scheme for the curve shortening flow*, Math. Models Methods Appl. Sci., 4 (1994), pp. 589–606.
- [12] K. ECKER, *Regularity Theory for Mean Curvature Flow*, Birkhäuser, Boston, 2004.
- [13] C. M. ELLIOTT AND H. FRITZ, *On approximations of the curve shortening flow and of the mean curvature flow based on the DeTurck trick*, IMA J. Numer. Anal., 37 (2017), pp. 543–603.
- [14] H. GARCKE AND B.-V. MATIOC, *On a degenerate parabolic system describing the mean curvature flow of rotationally symmetric closed surfaces*, J. Evol. Equ., (2020). (to appear).

- [15] S. KLEENE AND N. M. MÖLLER, *Self-shrinkers with a rotational symmetry*, Trans. Amer. Math. Soc., 366 (2014), pp. 3943–3963.
- [16] B. KOVÁCS, B. LI, AND C. LUBICH, *A convergent evolving finite element algorithm for mean curvature flow of closed surfaces*, Numer. Math., 143 (2019), pp. 797–853.
- [17] C. MANTEGAZZA, *Lecture notes on mean curvature flow*, vol. 290 of Progress in Mathematics, Birkhäuser/Springer Basel AG, Basel, 2011.
- [18] A. MIERSWA, *Error estimates for a finite difference approximation of mean curvature flow for surfaces of torus type*, PhD thesis, University Magdeburg, Magdeburg, 2020.