



Minkowski Inequalities via Nonlinear Potential Theory

VIRGINIA AGOSTINIANI, MATTIA FOGAGNOLO &
LORENZO MAZZIERI 

Communicated by A. FIGALLI

Abstract

In this paper, we prove an extended version of the Minkowski Inequality, holding for any smooth bounded set $\Omega \subset \mathbb{R}^n$, $n \geq 3$. Our proof relies on the discovery of *effective monotonicity formulas* holding along the level set flow of the p -capacitary potentials associated with Ω , for every p sufficiently close to 1. These formulas also testify the existence of a link between the monotonicity formulas derived by Colding and Minicozzi for the level set flow of Green's functions and the monotonicity formulas employed by Huisken, Ilmanen and several other authors in studying the geometric implications of the Inverse Mean Curvature Flow. In dimension $n \geq 8$, our conclusions are stronger than the ones obtained so far through the latter mentioned technique.

1. Introduction and Statements of the Main Results

A classical result in the theory of convex hypersurfaces in Euclidean spaces is the so called Minkowski inequality [52], which says that if $\Omega \Subset \mathbb{R}^n$, $n \geq 3$, is a convex domain with smooth boundary and H is the mean curvature of $\partial\Omega$ computed with respect to the outward unit normal, then

$$\left(\frac{|\mathbb{S}^{n-1}|}{|\partial\Omega|} \right)^{1/(n-1)} \leq \int_{\partial\Omega} \frac{H}{n-1} \, d\sigma$$

with equality if and only if Ω is a ball. In other words, the inverse of the surface radius is a sharp lower bound for the averaged total mean curvature of $\partial\Omega$. Observe that the above inequality can be conveniently rephrased as

$$\left(\frac{|\partial\Omega|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \frac{H}{n-1} \, d\sigma, \quad (1.1)$$

so that it can be combined with the standard Isoperimetric Inequality to deduce its volumetric version, also known in the literature as a higher order Isoperimetric Inequality (see [15] and [56])

$$\left(\frac{|\Omega|}{|\mathbb{B}^n|}\right)^{\frac{n-2}{n}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \frac{H}{n-1} \, d\sigma, \quad (1.2)$$

at least when Ω varies in the class of convex domains. It is worth recalling that both the Isoperimetric Inequality and the Minkowski Inequality are part of a family of inequalities involving quermassintegrals that were originally deduced in the context of convex analysis from the classical Aleksandrov–Fenchel mixed volume inequalities [6, 7, 24]. A natural question, raised by several authors (see [14, 15, 36, 61]), is whether the Minkowski Inequality (1.1) as well as its volumetric version (1.2) hold true for larger classes of domains than just for the convex one.

Positive answers to these questions have been proposed so far using the Inverse Mean Curvature Flow (IMCF from now on) and methods based on Optimal Transport. Our main concern in this paper is to propose an alternative technology based on Nonlinear Potential Theory, which is powerful enough to recover, improve and extend all the so far known results on these topics. Surprisingly, this new approach provides simplified arguments, which are also very flexible and likely to be exportable to several interesting frameworks, such as complete manifolds with nonnegative Ricci curvature and asymptotically flat manifolds with nonnegative scalar curvature. In Sect. 2, we will describe in more details the main features of this approach, drawing a systematic comparison with the existing curvature flow techniques. Here, we just anticipate that the cornerstone of our method is the discovery of *effective monotonicity formulas* (see Theorem 4.2 and Theorem 4.4), holding along the level sets of the p -capacitary potential $u_p : \mathbb{R}^n \setminus \overline{\Omega} \rightarrow \mathbb{R}$ associated with Ω . Besides their geometric implications, these formulas have a technical relevance on their own, as they persist through all the possible singularities of the flow. It is worth noticing that, in the present framework, the flow singularities correspond to the critical points of u_p , and these might in principle be arranged in sets of full measure. This means that, albeit the level set flow is possibly subject to jumps, our monotonicity formulas are strong enough to survive them. Finally, from a theoretical point of view, these formulas can be seen as the crucial step towards the completion of a program initiated in the series of works [1, 3, 4, 26] and intended to link the monotonicity formulas employed by Huisken, Ilmanen and other authors in studying the geometric implications of the IMCF (see e.g., [8–10, 21, 27, 29–31, 34, 36, 37, 45, 51, 64]) to the monotonicity formulas discovered by Colding and Minicozzi in [16–18] for the level set flow of the Green’s functions on complete manifolds with nonnegative Ricci curvature. In fact, as explained in Sect. 2.2, the first ones can be recovered from ours in the limit as $p \rightarrow 1^+$, whereas the latter can be reconstructed setting $p = 2$ and letting Ω shrink to a single point (see the Appendix of [1]).

We pass now to describing the main geometric inequalities obtained in this paper. The first one is an extension of the Minkowski Inequality, holding for every bounded and smooth subset of \mathbb{R}^n , in which the total mean curvature of the boundary

is replaced by the L^1 -norm of the mean curvature, whereas the perimeter of the set Ω is replaced by the one of its *strictly outward minimising hull* Ω^* , which is defined in Definition 5.2 below in accordance to [37, pp. 371–372]. For the reader's convenience we briefly recall that a set is called *outward minimising* if it minimises the perimeter among all the sets containing it; moreover, an outward minimising set is called *strictly outward minimising* if it coincides almost everywhere with any outward minimising set containing it and having the same perimeter. Loosely speaking, Ω^* is – up to negligible components – the smallest strictly outward minimising set that contains Ω (see Definitions 5.1 and 5.2 in Sect. 5 for more details). With these concepts at hand, our first main result reads as follows:

Theorem 1.1. (Extended Minkowski Inequality) *If $\Omega \subset \mathbb{R}^n$ is a bounded open set with smooth boundary, then*

$$\left(\frac{|\partial\Omega^*|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathbf{H}}{n-1} \right| d\sigma, \quad (1.3)$$

where Ω^* is the strictly outward minimising hull of Ω defined as in Definition 5.2. Moreover, the dimensional constants appearing here are optimal, in the sense that

$$\min \left\{ |\partial\Omega^*|^{-\frac{n-2}{n-1}} \int_{\partial\Omega} |\mathbf{H}| d\sigma \mid \Omega \Subset \mathbb{R}^n, \text{ with } \partial\Omega \text{ smooth} \right\} = (n-1) |\mathbb{S}^{n-1}|^{\frac{1}{n-1}},$$

and the minimum is achieved on spheres.

As a matter of fact, the Extended Minkowski Inequality (1.3) is deduced as the limit, for $p \rightarrow 1^+$, of the following geometric p -capacitary inequality, which we believe of independent interest.

Theorem 1.2. (L^p -Minkowski Inequality) *Let $\Omega \subset \mathbb{R}^n$ be an open bounded set with smooth boundary. Then, for every $1 < p < n$, the following inequality holds:*

$$C_p(\Omega)^{\frac{n-p-1}{n-p}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathbf{H}}{n-1} \right|^p d\sigma. \quad (1.4)$$

Here $C_p(\Omega)$ is the normalised p -capacity of Ω introduced in Definition 3.1. Moreover, equality holds in (1.4) if and only if Ω is a ball.

In order to deduce (1.3) from (1.4), one needs to compute the limit of the p -capacity of a bounded set with smooth boundary as $p \rightarrow 1^+$. Apart from the case of convex domains, treated in [66], we were unable to find in the literature a complete and satisfactory discussion of this very basic issue. For this reason, we have established that

$$\lim_{p \rightarrow 1^+} C_p(\Omega) = \frac{|\partial\Omega^*|}{|\mathbb{S}^{n-1}|}$$

in Theorem 5.6 of Sect. 5.

As an immediate corollary of Theorem 1.1, we recover the Minkowski Inequality for *outward minimising* sets, since for every Ω in this class it holds $|\partial\Omega| = |\partial\Omega^*|$

(see Remark 5.4) and $H \geq 0$, as a standard variational computation readily shows. Such inequality was originally conceived by Huisken in [36], exploiting the theory of weak solutions to the IMCF, previously developed in [37] (see also [27, Theorem 2–(b)] for a published version of the argument in the case of outward minimising sets with strictly mean-convex boundary). The conceptual differences between this method and our technique are described in Sect. 2, where we also discuss some of the technical subtleties arising in the two approaches.

Corollary 1.3. (Minkowski Inequality for Outward Minimising Sets) *If $\Omega \subset \mathbb{R}^n$ is a bounded outward minimising open set with smooth boundary, then*

$$\left(\frac{|\partial\Omega|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \frac{H}{n-1} \, d\sigma. \quad (1.5)$$

Moreover, the dimensional constants appearing here are optimal, in the sense that

$$\begin{aligned} & \min \left\{ |\partial\Omega|^{-\frac{n-2}{n-1}} \int_{\partial\Omega} H \, d\sigma \mid \Omega \Subset \mathbb{R}^n \text{ outward minimising, with } \partial\Omega \text{ smooth} \right\} \\ & = (n-1) |\mathbb{S}^{n-1}|^{\frac{1}{n-1}}, \end{aligned}$$

and the minimum is achieved on spheres. Viceversa, if the equality holds in (1.5) for some bounded strictly outward minimising open set with smooth and strictly mean-convex boundary, then Ω is isometric to a round ball.

It is worth pointing out that for $n \leq 7$, the above corollary can be shown to be equivalent to Theorem 1.1 at the price of a non completely trivial approximation procedure. Indeed, starting from a possibly non outward minimising Ω , one immediately checks that

$$\frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega^*} \frac{H}{n-1} \, d\sigma \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{H}{n-1} \right| \, d\sigma,$$

where Ω^* is the strictly outward minimising hull of Ω . On the other hand, when $n \leq 7$ the main result in [58] guarantees that $\partial\Omega^*$ is of class $\mathcal{C}^{1,1}$. In particular, with the help of [38, Lemma 2.6], it can be approximated in the $\mathcal{C}^1 \cap W^{2,1}$ -topology by a sequence of smooth and strictly mean-convex hypersurfaces. Arguing as in [37, Lemma 5.6] one realises that these hypersurfaces are bounding strictly outward minimising sets $\{\Omega_\varepsilon^*\}_{\varepsilon>0}$, so that inequality (1.5) is satisfied by the whole approximating sequence. Letting $\varepsilon \rightarrow 0$ yields

$$\left(\frac{|\partial\Omega^*|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega^*} \frac{H}{n-1} \, d\sigma,$$

from which the thesis follows at once. Obviously, this argument breaks down in higher dimension, due to the serious regularity issues coming from the possible presence of a non-empty singular set in the area minimising portion of $\partial\Omega^*$.

Remark 1.4. It is very likely that a perturbation argument similar to the ones in [35] might extend the above argument to $n = 8$, because in this case the singularities of the area minimising portion of the strictly outward minimising hull consist of isolated points.

A simple and very nice consequence of inequality (1.5) is a nearly umbilical estimate for *outward minimising* surfaces in \mathbb{R}^3 with *optimal constant*. The relation between the Minkowski Inequality and the nearly umbilical estimates was suggested by Huisken in [36]. Here, for the sake of reference, we included a proof of this fact in Sect. 6 (see Theorem 6.1). The general nearly umbilical estimate for surfaces in \mathbb{R}^2 with an implicit dimensional constant is a very remarkable theorem, proved in [20] by De Lellis and Müller. We refer the reader to the original paper [20] as well as to the Ph.D. thesis [54] and the references therein for a complete account about the geometric features and implications of such a deep result.

Another direct consequence of Theorem 1.1 is the following optimal version of inequality (1.2), holding for bounded open sets with smooth boundary.

Theorem 1.5. (Volumetric Minkowski Inequality) *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary. Then*

$$\left(\frac{|\Omega|}{|\mathbb{B}^n|} \right)^{\frac{n-2}{n}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathbf{H}}{n-1} \right| d\sigma. \quad (1.6)$$

Moreover, equality holds in (1.6) if and only if Ω is a ball.

Inequality (1.6) is achieved applying the Isoperimetric Inequality to the left hand side of (1.3), and taking into account that $\Omega \subseteq \Omega^*$, the latter being a hull. Observe that the rigidity statement in the above theorem follows essentially from the rigidity statement of the Isoperimetric Inequality. To the authors' knowledge, the above inequality was previously known to hold for domains with a strictly mean-convex boundary of positive scalar curvature (for short $\partial\Omega \in \Gamma_2^+$). On this regard, we refer the reader to the paper [15] and the subsequent [56], where the inequality is proved with methods based on Optimal Transport. For $n \leq 7$, we note that inequality (1.6) can also be deduced from Corollary 1.3 through the approximation argument outlined above.

1.1. Summary

In Sect. 2 we describe the main features of our method through a fairly systematic comparison with the previous approaches, based on the IMCF. Approximations schemes *à la Moser* [53] and formal analogies are employed to make some heuristic considerations as well as to introduce the main technical challenges of the present work. After collecting some preparatory material in Sect. 3, we face these challenges in Sect. 4, which constitutes the core of our analysis. There we solve the issues coming from the presence of critical points, proving *effective monotonicity formulas* (see Theorem 4.2 and Theorem 4.4), whose validity persists also beyond possible jumps. Having this tools at hand, we prove the L^p -Minkowski Inequality (1.4) and then, passing to the limit as $p \rightarrow 1^+$, we prove, in Sect. 5, the

Extended Minkowski Inequality (1.3). Here the main difficulty is to characterise in a geometrically meaningful way the limit of the p -capacity of the domain under consideration. We accomplish this task in Theorem 5.6.

2. Inverse Mean Curvature Flow Versus Nonlinear Potential Theory

Having introduced the main results of this paper in terms of geometric inequalities, we now describe the method that will be employed to deduce them. The most appropriate way to accomplish this task is to compare our approach set in nonlinear potential theory with the one based on the IMCF, in both its smooth and weak version.

2.1. Smooth and weak inverse mean curvature flow

Using the smooth IMCF it is possible to provide an extension of the classical Minkowski Inequality (1.1) for convex domains to the family of *starshaped domains with strictly mean-convex boundary*. This approach has been completely developed in [34] and relies essentially on the results in [32] and [63], where it is proven that if Ω is strictly mean-convex and starshaped, then the IMCF $\{\partial\Omega_t\}_{t \geq 0}$ starting at $\partial\Omega$ is defined and smooth for all times. In this case it is possible to carry out a smooth computation, showing that the function

$$t \longmapsto |\partial\Omega_t|^{-\frac{n-2}{n-1}} \int_{\partial\Omega_t} H \, d\sigma_t \tag{2.1}$$

is non increasing. The Minkowski Inequality then follows from the main theorems in [32,63]. These ensure that, as $t \rightarrow +\infty$, the hypersurfaces $\partial\Omega_t$ converge to a round sphere, once they are suitably rescaled, so that

$$|\partial\Omega|^{-\frac{n-2}{n-1}} \int_{\partial\Omega} H \, d\sigma \geq \lim_{t \rightarrow +\infty} |\partial\Omega_t|^{-\frac{n-2}{n-1}} \int_{\partial\Omega_t} H \, d\sigma_t = (n-1) |\mathbb{S}^{n-1}|^{\frac{1}{n-1}}. \tag{2.2}$$

This approach, which is extremely clean and quite flexible, has found remarkable applications also in non Euclidean contexts (see [10,21,29–31,49]). However, it is suitable only for those hypersurfaces that do not change topology along their evolution. For example, if the ambient manifold is (asymptotically) flat, it applies only to hypersurfaces with spherical topology.

These topological restrictions can be overtaken considering weak solutions of the IMCF starting at the boundary of *outward minimising sets*, as described in Huisken–Ilmanen’s theory [37,38]. In fact, weak solutions are engineered in order to allow for jumps, at which the topological changes take place, preserving at the same time the monotonicity of the quantity (2.1). On this regard, it is worth pointing out explicitly that *the class of outward minimising sets includes the one of starshaped domains with strictly mean-convex boundary*, as it can be easily deduced combining the long time existence established in [32,63] with a simple integration by parts argument.

Needless to say that the larger generality obtained through weak solutions comes at the cost of a much more sophisticated and delicate theory, whose extension to different contexts is not straightforward at all. For example, one of the major difficulties is to show that the monotonicity formulas survive the jumps. In [37] this is achieved by means of an elliptic regularisation procedure in which the weak solution of the IMCF is approximated by a family of smooth functions whose level sets obey a slightly modified version of the desired monotonicity. To the best of our knowledge, such a spectacular though technically demanding construction has never been replied beyond the original context of asymptotically flat Riemannian manifolds [27,36,37,51,64], with the only exception of [43, Theorem 3.2], where the authors have checked that Huisken–Ilmanen theory applies to the case under consideration. Hence, the expected extensions of the results in [10,21,29–31] to the case of outward minimising hypersurfaces are missing so far.

Another crucial point in the weak IMCF theory is the characterisation of the limits of the relevant monotone quantities. To be more specific, one would like to extend the validity of (2.2) to the class of weak solutions of the IMCF starting at an outward minimising domain. A possible way to accomplish this task is to invoke [38, Theorem 2.7-(b)] and then again [32, Theorem 0.1]. Indeed, the former affirms that the weak IMCF eventually becomes strictly starshaped at some large enough time, so that the latter applies yielding the desired asymptotic behaviour. The proof of [38, Theorem 2.7-(b)] relies in turn on a blow-down analysis that can be carried out following the proof of [37, Blowdown Lemma 7.1]. This argument represents a quite delicate point, to treat which a remarkable amount of theory needs to be employed. In the present paper, as explained in the next subsection, both the problem of dealing with possible jumps and the problem of characterising the limit of the relevant monotone quantities will be faced in a completely different way, with the help of nonlinear potential theory.

2.2. Level sets of p -capacitary potentials

The key point in our approach is to replace the weak IMCF technique with a novel analysis of a very natural family of functions, namely the p -capacitary potentials of Ω . These are the weak solutions to the problems

$$\begin{cases} \Delta_p u = 0 & \text{in } \mathbb{R}^n \setminus \overline{\Omega}, \\ u = 1 & \text{on } \partial\Omega, \\ u(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty, \end{cases}$$

with $1 < p < n$.

For heuristic considerations, it is useful to recall that these functions are playing an ancillary role in the weak IMCF theory, by virtue of a beautiful approximation result, originally due to Moser [53] and subsequently extended by Kotschwar and Ni [41] (see also the very recent [50]). This result says that if u_p is a weak solution to the above problem, then the functions $w_p = -(p - 1) \log u_p$ converge locally uniformly in $\mathbb{R}^n \setminus \Omega$ to a proper weak solution of the IMCF, as $p \rightarrow 1^+$. As a formal justification of this fact, it can be noted that the functions w_p satisfy the identities

$$\Delta_p w_p = |Dw_p|^p,$$

which in the limit as $p \rightarrow 1^+$ becomes

$$\operatorname{div} \left(\frac{Dw}{|Dw|} \right) = |Dw|. \quad (2.3)$$

As it is well known, the latter equation corresponds to the level set formulation of the IMCF, since the left hand side equals the mean curvature of the level sets of w .

As a concrete application of this elegant approximation scheme one immediately earns an alternative and flexible approach to the *existence* theory for the weak IMCF. Unfortunately, such a scheme is not as much effective for drawing geometric conclusions. For example, one would hopefully like to use it to deduce the validity of (2.2) from the nowadays classical asymptotic expansions of the u_p 's (see Lemma 3.3). However, the convergence is not strong enough to take the limit of the integral quantities, moreover it only holds on compact subsets of $\mathbb{R}^n \setminus \Omega$, preventing this strategy from being successfully implemented.

In view of these considerations and motivated by the recent discoveries in the field of harmonic functions [1, 2, 4], we then let the p -harmonic functions play the leading role, providing their already rich and well established theory with a so far missing ingredient, namely the existence of *monotonicity formulas* – or a relaxed version of them – holding along their level set flow and in presence of critical points. Once this will be done, we would get a self consistent theory, completely independent of the IMCF and powerful enough to recover and extend all the main geometric conclusions.

To clarify these concepts, let us first discuss the toy-problem case, where the p -capacitary potential has no critical points. In this case, which is treated in [26] assuming Ω convex, one finds that for every $1 < p < n$ the function

$$(0, 1] \ni \tau \mapsto U_p(\tau) = \tau^{-\frac{n-1}{n-p}} \int_{\{u_p=\tau\}} |Du_p|^p \, d\sigma \quad (2.4)$$

is nondecreasing. The monotonicity readily implies (1.4). In fact, computing the limit of U_p as $\tau \rightarrow 0^+$, with the help of Lemma 3.3 (see [26, Lemma 2.6] with $q = p/(p-1)$ for details), gives

$$\left(\frac{n-p}{p-1} \right)^p |\mathbb{S}^{n-1}| C_p(\Omega)^{\frac{n-p-1}{n-p}} = \lim_{\tau \rightarrow 0^+} U_p(\tau) \leq U_p(1) = \int_{\partial\Omega} |Du_p|^p \, d\sigma. \quad (2.5)$$

Next, one computes

$$0 \leq U_p'(1) = \frac{1}{(p-1)} \int_{\partial\Omega} |Du_p|^{p-1} H \, d\sigma - \left(\frac{n-1}{n-p} \right) \int_{\partial\Omega} |Du_p|^p \, d\sigma, \quad (2.6)$$

and thus, by Hölder's inequality, one gets

$$\int_{\partial\Omega} |Du_p|^p \, d\sigma \leq \left(\frac{n-p}{p-1} \right)^p \int_{\partial\Omega} \left| \frac{H}{n-1} \right|^p \, d\sigma.$$

Combining the latter inequality with (2.5) yields (1.4). The Extended Minkowski Inequality (1.3) can thus be obtained in the limit of (1.4), as $p \rightarrow 1^+$, by using the analysis of Sect. 5.

A question arises whether the monotonicity of the U_p 's and the one of the quantity defined in (2.1) are related or not. It turns out that the only way to answer this question is to proceed formally. Setting as above $w_p = -(p - 1) \log u_p$ and $t = -(p - 1) \log \tau$, for every $1 < p < n$, the monotonicity of the function U_p is equivalent to the statement that the function

$$[0, +\infty) \ni t \mapsto e^{-\frac{n-p-1}{n-p}t} \int_{\{w_p=t\}} |Dw_p|^p \, d\sigma$$

is nonincreasing. Taking the formal limit as $p \rightarrow 1^+$, one would get the same monotonicity statement for the function

$$[0, +\infty) \ni t \mapsto e^{-\frac{n-2}{n-1}t} \int_{\{w=t\}} |Dw| \, d\sigma ,$$

where w solves (2.3), and thus $|Dw|(x)$ coincides with the mean curvature of the level set passing through x . Recalling that $|\{w = t\}| = |\partial\Omega_t| = |\partial\Omega| e^t$ along the IMCF, it is easy to realise that the latter monotonicity is equivalent to the one in (2.1). However, it is important to emphasize once more that the above argument is just formal and does not prove anything, since w_p is converging to w only locally uniformly on compact sets and w itself is nothing more than a weak solution to the IMCF.

In presence of critical points for the p -capacitary potentials, the above formal derivation could appear in principle all the more *naïve*, since – unlike in the linear case treated in [4] – the monotonicity of (2.4) is not even *a priori* guaranteed. This phenomenon is typical of the nonlinear setting and is basically due to the loss of analyticity of the solution and the consequent loss of control on the behaviour of the critical points and of the critical values. In particular, in the case of the p -capacitary potential, one cannot exclude *a priori* the presence of clusters of critical points and critical values with full measure. Due to these difficulties, it is impossible to re-adapt the strategy employed in the linear case [4] to earn the *full monotonicity* of the U_p 's. However, we will be able to prove in the next sections that the inequalities

$$0 \leq U_p'(1) \quad \text{and} \quad \lim_{\tau \rightarrow 0^+} U_p(\tau) \leq U_p(1) \tag{2.7}$$

hold true. These inequalities – together with their conformal counterparts defined below (3.10) – will be referred to as *effective inequalities* and will be deduced in Sect. 4 as consequences of some *effective monotonicity* properties of the functions U_p (see Theorems 4.2 and 4.4 combined with formulæ (3.11)). Here, the locution “effective monotonicity” should be understood in contrast with the “full monotonicity” which is instead enjoyed by the U_p 's either in the case where Ω is convex (see [26]) or in the case $p = 2$ (see [4]).

As explained through (2.5) and (2.6), the two conditions in (2.7) are sufficient to deduce the L^p -Minkowski Inequality (1.4) and in turn the Extended Minkowski Inequality (1.3). It must be noticed that the proof of the inequalities (2.7) requires

both a technical and a conceptual enhancement of the previously existing techniques ([3, 4, 26]). This is particularly evident in the analysis leading to the second inequality, which is based on the discovery of a further family of monotonic functions (see Theorem 4.4), whose existence was far beyond the horizon both of the linear case [4] and of the nonlinear convex case [26].

2.3. Further directions

In virtue of the previous observations it is quite clear that methods based on linear and nonlinear potential theory may provide an efficient alternative to the employment of the IMCF techniques in many contexts. To be concrete, let us just mention a couple of projects that represent a natural continuation of the present work.

The first one is the potential revisitation of the IMCF proof of the Riemannian Penrose Inequality due to Huisken and Ilmanen [37]. Indeed, it is not too hard to provide a formal guess of the monotonic quantities which are expected to play the same role as the Hawking Mass in the Geroch's monotonicity scheme. The hard part, as usual, is the treatment of the critical points. There are good reasons to believe that the method and the ideas presented in this work also apply to that situation.

The second one is the extension of the Minkowski Inequalities to the case of complete manifolds with nonnegative Ricci curvature. Proceeding in parallel with the linear theory (compare [4, Theorem 1.1] with [1, Theorem 1.3]), one can prove that if the manifold (M, g) is Asymptotically Locally Euclidean and Ω is an outward minimising domain, then it holds

$$\left(\text{AVR}(g) \frac{|\mathbb{S}^{n-1}|}{|\partial\Omega|} \right)^{1/(n-1)} \leq \int_{\partial\Omega} \frac{H}{n-1} \, d\sigma,$$

where $\text{AVR}(g)$ stands for the Asymptotic Volume Ratio of (M, g) . A detailed proof of this result will appear in a forthcoming manuscript.

Other challenges include the study of natural geometric inequalities in Cartan-Hadamard manifolds as well as in Asymptotically Hyperbolic manifolds.

3. Preparatory Material

3.1. Preliminaries on p -capacitary potentials

We recall the well known notion of p -capacity, introducing at the same time a normalised version of it that is suitable for our applications.

Definition 3.1. (*p -capacity & normalised p -capacity*) Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary.

- The p -capacity of Ω is defined as

$$\text{Cap}_p(\Omega) = \inf \left\{ \int_{\mathbb{R}^n} |\text{D}v|^p \, d\mu \mid v \in \mathcal{C}_c^\infty(\mathbb{R}^n), v \geq 1 \text{ on } \Omega \right\}.$$

- The normalised p -capacity of Ω is defined as

$$C_p(\Omega) = \inf \left\{ \left(\frac{p-1}{n-p} \right)^{p-1} \frac{1}{|\mathbb{S}^{n-1}|} \int_{\mathbb{R}^n} |\text{D}v|^p \, d\mu \mid v \in \mathcal{C}_c^\infty(\mathbb{R}^n), v \geq 1 \text{ on } \Omega \right\}. \quad (3.1)$$

The variational structure of the above definition leads naturally to the formulation of the following problem

$$\begin{cases} \Delta_p u = 0 & \text{in } \mathbb{R}^n \setminus \overline{\Omega}, \\ u = 1 & \text{on } \partial\Omega, \\ u(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty. \end{cases} \quad (3.2)$$

It is well known that, for every bounded open set Ω with smooth boundary and every $1 < p < n$, problem (3.2) admits a unique weak solution. Such a solution is called the p -capacitary potential associated with Ω . For the reader's convenience, we recall that a function v is a weak solution of $\Delta_p v = 0$ in an open set V if $v \in W_{loc}^{1,p}(V)$ and

$$\int_V \left\langle |\text{D}v|^{p-2} \text{D}v \mid \text{D}\psi \right\rangle \, d\mu = 0$$

for any test function $\psi \in \mathcal{C}_c^\infty(V)$. By the important contributions [22,23,44] and [62], we know that weakly p -harmonic functions are $\mathcal{C}_{loc}^{1,\alpha}$ (we are not aware of an explicit formula relating α to p ; we note, however, that such relation cannot be uniform in p). On the other hand, the classical regularity theory for quasilinear nondegenerate elliptic equations (see e.g. [42]) ensures that they are analytic around the points where the gradient does not vanish. We also recall from [46] that the $\mathcal{C}^{1,\alpha}$ -regularity can be extended up to the boundary.

Note that the uniqueness of the solution to problem (3.2) can be easily proved by suitably applying the Comparison Theorem for weakly p -harmonic functions [47, Theorem 2.15] on large balls of radius R , and letting then $R \rightarrow +\infty$. With the same argument one can also show that the solution u to problem (3.2) is such that $0 < u(x) < 1$ for every $x \in \mathbb{R}^n \setminus \Omega$. Finally, we recall that such a solution realises the infimum in (3.1). This can be proved using a standard exhaustion scheme (for example the one proposed in [19]) and invoking the $\mathcal{C}_{loc}^{1,\alpha}$ regularity to guarantee the convergence of the scheme itself. These facts are summarised in the following theorem:

Theorem 3.2. (Existence and regularity of p -capacitary potentials) *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary, and let $1 < p < n$. Then, the following statements hold true:*

- (i) *There exists a unique weak solution $u \in \mathcal{C}_{loc}^{1,\alpha}(\mathbb{R}^n \setminus \overline{\Omega}) \cap \mathcal{C}(\mathbb{R}^n \setminus \Omega)$ to problem (3.2).*
- (ii) *The solution u is analytic at points where $Du \neq 0$.*
- (iii) *The solution u fulfills*

$$C_p(\Omega) = \left(\frac{p-1}{n-p}\right)^{p-1} \frac{1}{|\mathbb{S}^{n-1}|} \int_{\mathbb{R}^n \setminus \overline{\Omega}} |Du|^p \, d\mu, \quad (3.3)$$

where $C_p(\Omega)$ is the normalised p -capacity of Ω defined in (3.1).

Note that since $\partial\Omega$ is assumed to be smooth, by the Hopf Lemma for p -harmonic functions (see [60, Proposition 3.2.1]), we have that $|Du| \neq 0$ in a neighbourhood of this hypersurface. Coupled with this fact, the asymptotic expansions below imply that $\text{Crit}(u) = \{x \in \mathbb{R}^n \setminus \overline{\Omega} \mid Du(x) = 0\}$ is a compact subset of $\mathbb{R}^n \setminus \overline{\Omega}$ (generically depending on p). Finally, it is worth recalling that for $p \neq 2$, the set $\text{Crit}(u)$ is *a priori* allowed to have full measure.

Lemma 3.3. (Asymptotic expansions of u and $|Du|$) *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary, and let $1 < p < n$. Then, the solution u to (3.2) satisfies*

- (i) $\lim_{|x| \rightarrow +\infty} u(x) |x|^{\frac{n-p}{p-1}} = C_p(\Omega)^{\frac{1}{p-1}},$
- (ii) $\lim_{|x| \rightarrow +\infty} |Du(x)| |x|^{\frac{n-1}{p-1}} = \left(\frac{n-p}{p-1}\right) C_p(\Omega)^{\frac{1}{p-1}},$

where $C_p(\Omega)$ is the normalised p -capacity of Ω defined in (3.1). In particular, $\text{Crit}(u)$ is a compact subset of $\mathbb{R}^n \setminus \overline{\Omega}$, possibly with full measure.

For the proof of this lemma we refer the reader to [40] (see also the more recent [55, Lemma 2.3 and (2.2)] for a precise statement). It is also worth mentioning [28], where similar expansions are employed to infer rotational symmetry of starshaped domains supporting a solution to problem (3.2) with constant normal derivative on the boundary. From the point of view of the present paper, the main implication of the above lemma is the computation of the limit

$$\lim_{\tau \rightarrow 0^+} U_p(\tau) = \left(\frac{n-p}{p-1}\right)^p |\mathbb{S}^{n-1}| C_p(\Omega)^{\frac{n-p-1}{n-p}}, \quad (3.4)$$

where $\tau \mapsto U_p(\tau)$ is the function defined in (2.4) (see [26, Lemma 2.6]). The following characterisation of the p -capacity of Ω is widely used in the literature and it is also very useful for our purposes. Hence, we provide a proof.

Lemma 3.4. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary, and let $1 < p < n$. Then, the solution u to (3.2) satisfies*

$$C_p(\Omega) = \left(\frac{p-1}{n-p}\right)^{p-1} \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} |Du|^{p-1} \, d\sigma,$$

where $C_p(\Omega)$ is the normalised p -capacity of Ω defined in (3.1).

Proof. For $\varepsilon > 0$, let V_ε be the ε -tubular neighbourhood of $\text{Crit}(u)$, namely

$$V_\varepsilon = \left\{ x \in \mathbb{R}^n \setminus \overline{\Omega} \mid \text{dist}(x, \text{Crit}(u)) < \varepsilon \right\},$$

where $\text{dist}(x, \text{Crit}(u))$ is the Euclidean distance of x from $\text{Crit}(u)$. By the compactness of $\text{Crit}(u)$ in $\mathbb{R}^n \setminus \overline{\Omega}$, we have that $V_\varepsilon \subset \{u \geq t\}$, for every $\varepsilon > 0$ and $t > 0$ small enough. Since $|\text{Du}| = 0$ on $\text{Crit}(u)$ by definition, we have from identity (3.3) and by Monotone Convergence Theorem that

$$\left(\frac{n-p}{p-1}\right)^{p-1} |\mathbb{S}^{n-1}| C_p(\Omega) = \lim_{\varepsilon \rightarrow 0^+} \lim_{t \rightarrow 0^+} \int_{\{u \geq t\} \setminus V_\varepsilon} |\text{Du}|^p \, d\mu.$$

By the discussions above, u is analytic – and in turn p -harmonic in the classical sense – in the set $\{u \geq t\} \setminus V_\varepsilon$. Therefore, for ε and t small enough, the Divergence Theorem yields

$$\begin{aligned} \int_{\{u \geq t\} \setminus V_\varepsilon} |\text{Du}|^p \, d\mu &= \int_{\{u \geq t\} \setminus V_\varepsilon} \text{div} \left(u |\text{Du}|^{p-2} \text{Du} \right) \, d\mu = \\ &= -t \int_{\{u=t\}} |\text{Du}|^{p-1} \, d\sigma + \int_{\partial V_\varepsilon} |\text{Du}|^{p-1} \, d\sigma \\ &\quad + \int_{\partial V_\varepsilon} u |\text{Du}|^{p-2} \langle \text{Du} \mid \nu \rangle \, d\sigma, \end{aligned}$$

where ν is the inward unit normal to V_ε . Observe that ν is well defined almost everywhere on ∂V_ε and for almost every $\varepsilon > 0$, in view of the Sard-type property for Lipschitz functions proved in [5]. Now observe that for every small enough t the level set $\{u = t\}$ is regular and $\int_{\{u=t\}} |\text{Du}|^{p-1} \, d\sigma$ is constant, by the Divergence Theorem. It follows that the first term in the rightmost hand side of the above formula vanishes as $t \rightarrow 0^+$. Also, when $\varepsilon \rightarrow 0^+$ the integral on ∂V_ε tends to 0, since $|\text{Du}|^{p-1}(x)$ vanishes as x approaches $\text{Crit}(u)$ and u remains bounded. \square

In the following subsection as well as in the remaining part of the paper we will always assume that $1 < p < n$, unless otherwise stated.

3.2. The conformal setting

As shown in [1, 3, 4] and [26], it is very convenient to work in the conformally related Riemannian manifold $(\mathbb{R}^n \setminus \Omega, g)$, where g is given by

$$g = u^{2\left(\frac{p-1}{n-p}\right)} g_{\mathbb{R}^n}. \quad (3.5)$$

In this setting it is also convenient to consider the new variable

$$\varphi = -\frac{(p-1)(n-2)}{(n-p)} \log u. \quad (3.6)$$

By the same formal computations as in [26], the boundary value problem (3.2) translates in terms of g and φ as

$$\left\{ \begin{array}{ll} \Delta_p^g \varphi = 0 & \text{in } \mathbb{R}^n \setminus \overline{\Omega}, \\ \text{Ric}_g - \nabla \nabla \varphi + \frac{d\varphi \otimes d\varphi}{n-2} = \left(\frac{|\nabla \varphi|_g^2}{n-2} - \frac{(p-2)}{(n-2)} \frac{\nabla \nabla \varphi (\nabla \varphi, \nabla \varphi)}{|\nabla \varphi|_g^2} \right) g & \text{in } (\mathbb{R}^n \setminus \overline{\Omega}) \setminus \text{Crit}(\varphi), \\ \varphi = 0 & \text{on } \partial \Omega, \\ \varphi(x) \rightarrow +\infty & \text{as } x \rightarrow \infty. \end{array} \right.$$

Here, ∇ is the Levi-Civita connection of g , $\nabla \nabla$ the Hessian operator, and Δ_p^g is the p -Laplace operator computed with respect to the metric g , explicitly defined as

$$\Delta_p^g \varphi = \text{div}_g (|\nabla \varphi|^{p-2} \nabla \varphi),$$

where div_g is the divergence computed with respect to g . A very useful tool in the study of p -harmonic functions is the Kato-type identity, introduced in [26, Proposition 4.4]. For the reader's convenience, we recall its precise statement in the following proposition:

Proposition 3.5. (Kato-type identity & orthogonal decomposition) *Let (M, g) be a Riemannian manifold, and let φ be a p -harmonic function on M . Then, at any point where $|\nabla \varphi| \neq 0$, the following identity holds true*

$$\begin{aligned} |\nabla \nabla \varphi|^2 - \left(1 + \frac{(p-1)^2}{n-1} \right) |\nabla |\nabla \varphi||^2 &= |\nabla \varphi|^2 \left| \mathfrak{h} - \frac{\mathbf{H}}{n-1} g^\top \right|^2 \\ &+ \left(1 - \frac{(p-1)^2}{n-1} \right) |\nabla^\top |\nabla \varphi||^2, \end{aligned} \quad (3.7)$$

where \mathfrak{h} and \mathbf{H} are respectively the second fundamental form and the mean curvature of the level sets of φ with respect to the unit normal $\nabla \varphi / |\nabla \varphi|$, and g^\top is the metric induced by g on the level sets of φ . Finally, for a given differentiable function f , we agree that $\nabla^\top f$ indicates the tangential part of the gradient, according to the orthogonal decomposition

$$\nabla^\perp f = \left\langle \nabla f \left| \frac{\nabla \varphi}{|\nabla \varphi|} \right\rangle \frac{\nabla \varphi}{|\nabla \varphi|} \quad \text{and} \quad \nabla^\top f = \nabla f - \nabla^\perp f.$$

In particular, the following formula holds true:

$$|\nabla |\nabla \varphi||^2 = |\nabla^\top |\nabla \varphi||^2 + |\nabla^\perp |\nabla \varphi||^2. \quad (3.8)$$

Since the proof of the L^p -Minkowski Inequality outlined in Sect. 2.2 will be carried out in the conformal setting described above, the fundamental conditions (2.7) need to be rephrased accordingly. It is then worth introducing the following definition.

Definition 3.6. (The function Φ_p) For any $1 < p < n$, let g and φ be defined by (3.5) and (3.6). We define the function $\Phi_p : [0, +\infty) \rightarrow \mathbb{R}$ by

$$\Phi_p(s) = \int_{\{\varphi=s\}} |\nabla\varphi|_g^p \, d\sigma_g, \quad (3.9)$$

where $d\sigma_g$ is the area element induced by the ambient measure $d\mu_g$ on the given level set. We agree that

$$\int_{\{\varphi=s\}} |\nabla\varphi|_g^p \, d\sigma_g = \int_{\{\varphi=s\} \setminus \text{Crit}(\varphi)} |\nabla\varphi|_g^p \, d\sigma_g,$$

whenever a critical value is involved.

We conclude this subsection, recalling some of the relevant properties of the function Φ_p just introduced. Their proofs are basically immediate – as they follows from the analogous properties of the corresponding function U_p , defined in (2.4) – and are left to the reader.

- The function Φ_p is bounded at infinity. Moreover, it follows from (3.4) that

$$\lim_{s \rightarrow +\infty} \Phi_p(s) = (n-2)^p |\mathbb{S}^{n-1}| C_p(\Omega)^{\frac{n-p-1}{n-p}}.$$

- The function Φ_p is differentiable at the regular values of φ .
- In terms of Φ_p , the effective inequalities (2.7) correspond to

$$\Phi'_p(0) \leq 0 \quad \text{and} \quad \lim_{s \rightarrow +\infty} \Phi_p(s) \leq \Phi_p(0). \quad (3.10)$$

In fact it is easily seen that

$$\begin{aligned} \Phi_p(s) &= \left[\frac{(n-2)(p-1)}{(n-p)} \right]^p U_p \left(e^{-\frac{(n-p)}{(n-2)(p-1)}s} \right), \\ \Phi'_p(s) &= - \left[\frac{(n-2)(p-1)}{(n-p)} \right]^{p-1} e^{-\frac{(n-p)}{(n-2)(p-1)}s} U'_p \left(e^{-\frac{(n-p)}{(n-2)(p-1)}s} \right), \end{aligned} \quad (3.11)$$

whenever these objects are well defined.

The inequalities (3.10) are at the core of our analysis and will be deduced in Sect. 4, as consequences of our *effective monotonicity formulas* (see Theorem 4.2 and Theorem 4.4).

4. Proof of the L^p -Minkowski Inequality

The aim of this section is to give a complete proof of Theorem 1.2, namely the L^p -Minkowski Inequality

$$C_p(\Omega)^{\frac{n-p-1}{n-p}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathbf{H}}{n-1} \right|^p \, d\sigma.$$

In force of the discussion in Sect. 2.2 (see also Sect. 4.3 below for a fully detailed proof), it is sufficient to establish the validity of the inequalities (2.7) in their conformal version (3.10)

$$\Phi'_p(0) \leq 0 \quad \text{and} \quad \Phi_p(+\infty) = \lim_{s \rightarrow +\infty} \Phi_p(s) \leq \Phi_p(0).$$

Since all the computations of this section will be performed in the conformally related setting, the subscript g will be dropped from the notations.

4.1. First effective inequality: $\Phi'_p(0) \leq 0$.

For a given $1 < p < n$, let us consider the vector field

$$X = e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p-2} \left(\nabla |\nabla\varphi| + (p-2)\nabla^\perp |\nabla\varphi| \right). \quad (4.1)$$

As it can be readily checked, at a regular value of φ one has that

$$e^{-\frac{(n-p)}{(n-2)(p-1)}s} \Phi'_p(s) = \frac{1}{p-1} \int_{\{\varphi=s\}} \left\langle X \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle d\sigma. \quad (4.2)$$

In the next lemma, we compute the divergence of X .

Lemma 4.1. (Divergence of X) *For any $1 < p < n$, let g and φ be defined by (3.5) and (3.6), and let X be the vector field defined in (4.1). Then, the following identity holds at any point $x \in \mathbb{R}^n \setminus \overline{\Omega}$ such that $|\nabla\varphi|(x) \neq 0$.*

$$\operatorname{div} X = e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} Q \geq 0, \quad (4.3)$$

where

$$Q = |\nabla\varphi|^{p-3} \left\{ |\nabla\varphi|^2 \left| \mathbf{h} - \frac{\mathbf{H}}{n-1} g^\top \right|^2 + (p-1) \left| \nabla^\top |\nabla\varphi| \right|^2 + \frac{(p-1)^2}{n-1} \left| \nabla^\perp |\nabla\varphi| \right|^2 \right\}, \quad (4.4)$$

where \mathbf{h} and \mathbf{H} are respectively the second fundamental form and the mean curvature of the level sets of φ with respect to the unit normal $\nabla\varphi/|\nabla\varphi|$.

Proof. For the sake of clearness, we write

$$X = e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} (W + Z),$$

where

$$W = |\nabla\varphi|^{p-2} \nabla |\nabla\varphi| \quad \text{and} \quad Z = (p-2) |\nabla\varphi|^{p-2} \nabla^\perp |\nabla\varphi|.$$

Using the same computation as in [26, Proposition 4.3] with $q = p/(p-1)$ (in the notation of that paper), one finds that the divergence of W is given by

$$\begin{aligned} \operatorname{div} W &= \left(\frac{n-p}{n-2} \right) \langle W \mid \nabla \varphi \rangle + \\ &+ |\nabla \varphi|^{p-3} \left\{ |\nabla \nabla \varphi|^2 - \left| \nabla |\nabla \varphi| \right|^2 \right. \\ &\left. + (p-2) \left[\left| \nabla^\perp |\nabla \varphi| \right|^2 - \frac{\nabla \nabla |\nabla \varphi| (\nabla \varphi, \nabla \varphi)}{|\nabla \varphi|} \right] \right\}. \end{aligned} \quad (4.5)$$

Plugging the Kato-type identity (3.7) in (4.5) and using the standard decomposition (3.8), we immediately get

$$\begin{aligned} \operatorname{div} W &= \left(\frac{n-p}{n-2} \right) \langle W \mid \nabla \varphi \rangle + \\ &+ |\nabla \varphi|^{p-3} \left\{ |\nabla \varphi|^2 \left| \mathfrak{h} - \frac{\mathbf{H}}{n-1} g^\top \right|^2 - (p-2) \frac{\nabla \nabla |\nabla \varphi| (\nabla \varphi, \nabla \varphi)}{|\nabla \varphi|} + \right. \\ &\left. + \left| \nabla^\top |\nabla \varphi| \right|^2 + \left[\frac{(p-1)^2}{n-1} + (p-2) \right] \left| \nabla^\perp |\nabla \varphi| \right|^2 \right\}. \end{aligned} \quad (4.6)$$

Let us now compute the divergence of Z . Clearly, by the p -harmonicity of φ , we have

$$\operatorname{div} Z = (p-2) |\nabla \varphi|^{p-2} \left\langle \nabla \left(\left\langle \nabla |\nabla \varphi| \mid \frac{\nabla \varphi}{|\nabla \varphi|^2} \right\rangle \right) \mid \nabla \varphi \right\rangle.$$

Expanding the right hand side and using the identity

$$\frac{\nabla \nabla \varphi (\nabla |\nabla \varphi|, \nabla \varphi)}{|\nabla \varphi|} = \left| \nabla |\nabla \varphi| \right|^2$$

yield

$$\operatorname{div} Z = (p-2) |\nabla \varphi|^{p-3} \left\{ \frac{\nabla \nabla |\nabla \varphi| (\nabla \varphi, \nabla \varphi)}{|\nabla \varphi|} + \left| \nabla^\top |\nabla \varphi| \right|^2 - \left| \nabla^\perp |\nabla \varphi| \right|^2 \right\}. \quad (4.7)$$

Finally, combining (4.6) and (4.7), and observing that

$$\langle X \mid \nabla \varphi \rangle = e^{-\frac{(n-p)}{(n-2)(p-1)} \varphi} (p-1) \langle W \mid \nabla \varphi \rangle,$$

we arrive at

$$\operatorname{div} X = e^{-\frac{(n-p)}{(n-2)(p-1)} \varphi} \left(\operatorname{div} W + \operatorname{div} Z - \left(\frac{n-p}{n-2} \right) \langle W \mid \nabla \varphi \rangle \right) = e^{-\frac{(n-p)}{(n-2)(p-1)} \varphi} Q.$$

This completes the proof of the lemma. \square

In absence of critical points, the Divergence Theorem applied to the vector field X on the open region $\{s < \varphi < S\}$, with $0 < s < S$, easily yields the inequality

$$\int_{\{\varphi=s\}} \left\langle X \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma \leq \int_{\{\varphi=S\}} \left\langle X \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma,$$

and in turns, thanks to (4.2), the inequality (4.8) below. In presence of a possibly wild critical set, this direct argument is no longer working. Fortunately, some of the new ideas introduced in [4] to treat the same issues in the case of harmonic functions are exportable to the case of p -harmonic functions, where one does not know *a priori* that the critical set is $(n - 1)$ -negligible. As a consequence, we are still able to provide an effective version of the considered monotonicity, showing that (4.8) is actually in force, provided s is small enough and S is large enough. The desired effective inequality $\Phi'_p(0) \leq 0$, will follow at once.

Theorem 4.2. (Effective Monotonicity Formula – I) *For any $1 < p < n$, let g and φ be defined by (3.5) and (3.6) and let $0 < s_p < S_p < +\infty$ be such that $\text{Crit}(\varphi) \subset \{s_p < \varphi < S_p\}$. Then, for every $0 \leq s \leq s_p < S_p \leq S$, the inequality*

$$\frac{\Phi'_p(s)}{e^{\frac{(n-p)}{(n-2)(p-1)}s}} \leq \frac{\Phi'_p(S)}{e^{\frac{(n-p)}{(n-2)(p-1)}S}} \quad (4.8)$$

holds true, where Φ_p is the function defined in (3.9). In particular, one has that $\Phi'_p(0) \leq 0$.

Proof. For a given $\varepsilon > 0$, we consider a smooth nonnegative cut-off-function $\chi : [0, +\infty) \rightarrow \mathbb{R}$, such that

$$\begin{cases} \chi(t) = 0 & \text{in } t < \frac{1}{2}\varepsilon, \\ \dot{\chi}(t) \geq 0 & \text{in } \frac{1}{2}\varepsilon \leq t \leq \frac{3}{2}\varepsilon, \\ \chi(t) = 1 & \text{in } t > \frac{3}{2}\varepsilon. \end{cases} \quad (4.9)$$

Since $\chi(|\nabla\varphi|) = 0$ on $\text{Crit}(\varphi)$, we can apply the Divergence Theorem to the smooth vector field

$$\tilde{X} = \chi(|\nabla\varphi|) X$$

in the domain $\{s < \varphi < S\}$. Observe that, choosing ε small enough, we can make sure that $\chi(|\nabla\varphi|) = 1$ on $\{\varphi = s\}$ and $\{\varphi = S\}$, since $\text{Crit}(\varphi) \subset \{s_p < \varphi < S_p\}$. Having this in mind, we compute

$$\begin{aligned} \int_{\{\varphi=S\}} \left\langle X \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma - \int_{\{\varphi=s\}} \left\langle X \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma &= \int_{\{s < \varphi < S\}} \text{div} \tilde{X} d\mu \\ &= \int_{\{s < \varphi < S\} \setminus U_{\varepsilon/2}} \chi(|\nabla\varphi|) \text{div} X d\mu + \int_{U_{3\varepsilon/2} \setminus U_{\varepsilon/2}} \dot{\chi}(|\nabla\varphi|) \langle X | \nabla|\nabla\varphi| \rangle d\mu, \end{aligned} \quad (4.10)$$

where in the last identity we have used the tubular neighbourhood of $\text{Crit}(\varphi)$ defined for every $\delta > 0$ as $U_\delta = \{|\nabla\varphi| \leq \delta\}$. In view of (4.2), (4.3), (4.9) and (4.10), the

inequality (4.8) is proved if we show that $\langle X | \nabla |\nabla \varphi| \rangle \geq 0$ on $U_{3\varepsilon/2} \setminus U_{\varepsilon/2}$. On the other hand, a direct computation gives

$$\begin{aligned} \langle X | \nabla |\nabla \varphi| \rangle &= e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla \varphi|^{p-2} \left[|\nabla |\nabla \varphi||^2 + (p-2) |\nabla^\perp |\nabla \varphi||^2 \right] = \\ &= e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla \varphi|^{p-2} \left[|\nabla^\top |\nabla \varphi||^2 + (p-1) |\nabla^\perp |\nabla \varphi||^2 \right] \geq 0. \end{aligned}$$

This completes the proof of the first part of the statement. It remains to show that $\Phi'_p(0) \leq 0$. From (4.8) it follows at once that, for every $S \geq S_p$, it holds

$$e^{\frac{(n-p)}{(n-2)(p-1)}S} \Phi'_p(0) \leq \Phi'_p(S).$$

Integrating both sides of the above inequality on an interval of the form (S_p, S) , with $S_p < S$, we obtain

$$\begin{aligned} \frac{(n-2)(p-1)}{(n-p)} e^{\frac{(n-p)}{(n-2)(p-1)}S} \Phi'_p(0) + \Phi_p(S_p) \\ - \frac{(n-2)(p-1)}{(n-p)} e^{\frac{(n-p)}{(n-2)(p-1)}S_p} \Phi'_p(0) \leq \Phi_p(S). \end{aligned}$$

If by contradiction, $\Phi'_p(0) > 0$, then, letting $S \rightarrow +\infty$ in the above inequality, we would deduce that $\Phi_p(S) \rightarrow +\infty$, against the boundedness of Φ_p discussed at the end of Sect. 3.2. \square

4.2. Second effective inequality: $\Phi_p(+\infty) \leq \Phi_p(0)$.

As already observed several times, the presence of critical points and critical values possibly arranged in sets with full measure makes the full monotonicity not expectable in general. In fact, the lack of a sufficiently strong Sard-type property for the p -capacitary potentials prevents any kind of straightforward adaptation of the arguments presented in [4] (it is worth mentioning though [12, 13], where a generic non-fattening property is proved for the level sets of p -harmonic functions). In other words, there is no hope for deducing the global inequality $\Phi_p(+\infty) \leq \Phi_p(0)$ from the pointwise inequality $\Phi'_p(s) \leq 0$ through integration, since the latter inequality may fail to be true – or even well defined – for too many values of $s \in [0, +\infty)$. To face the main difficulty of our program, we craft a new family of effective monotonicity formulas. For a given $1 < p < n$ and a given $0 < \lambda < 1$, we consider the vector field

$$Y_\lambda = \left(e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda \right) X - \left(\frac{n-p}{n-2} \right) |\nabla \varphi|^{p-1} \nabla \varphi, \quad (4.11)$$

where X has been defined in (4.1). It is convenient to observe that at a regular value of φ it holds

$$\begin{aligned} & \left(\frac{e^{\frac{(n-p)}{(n-2)(p-1)}s} - \lambda}{e^{\frac{(n-p)}{(n-2)(p-1)}s}} \right) \Phi'_p(s) - \frac{(n-p)}{(n-2)(p-1)} \Phi_p(s) \\ &= \frac{1}{(p-1)} \int_{\{\varphi=s\}} \left\langle Y_\lambda \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle d\sigma. \end{aligned} \quad (4.12)$$

In the next lemma, we compute the divergence of Y_λ .

Lemma 4.3. (Divergence of Y_λ) *For any $1 < p < n$ and any $0 < \lambda < 1$, let g and φ be defined by (3.5) and (3.6) and let Y_λ be the vector field defined in (4.11). Then, the following identity holds at any point $x \in \mathbb{R}^n \setminus \bar{\Omega}$ such that $|\nabla\varphi|(x) \neq 0$*

$$\operatorname{div} Y_\lambda = \left(\frac{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda}{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi}} \right) Q \geq 0,$$

where Q is the nonnegative quantity defined in (4.4).

Proof. By the very definition of Y_λ , we have that

$$\begin{aligned} \operatorname{div} Y_\lambda &= \left(e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda \right) \operatorname{div} X + \frac{(n-p)}{(n-2)(p-1)} e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} \langle X | \nabla\varphi \rangle \\ &\quad - \left(\frac{n-p}{n-2} \right) \operatorname{div} (|\nabla\varphi|^{p-1} \nabla\varphi). \end{aligned}$$

Using the definition (4.1) of the vector field X , we compute

$$\langle X | \nabla\varphi \rangle = (p-1) e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p-2} \langle \nabla | \nabla\varphi | | \nabla\varphi \rangle.$$

Exploiting the p -harmonicity of φ , we get

$$\operatorname{div} (|\nabla\varphi|^{p-1} \nabla\varphi) = |\nabla\varphi|^{p-2} \langle \nabla | \nabla\varphi | | \nabla\varphi \rangle.$$

We conclude that

$$\operatorname{div} Y_\lambda = \left(e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda \right) \operatorname{div} X = \left(\frac{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda}{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi}} \right) Q,$$

where in the last equality we made use of the identity (4.3). \square

Again, in absence of critical points, the Divergence Theorem applied to the vector field Y_λ on the open region $\{s < \varphi < S\}$ easily yields the inequality

$$\int_{\{\varphi=s\}} \left\langle Y_\lambda \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle d\sigma \leq \int_{\{\varphi=S\}} \left\langle Y_\lambda \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle d\sigma,$$

and in turn, thanks to (4.12), the inequality (4.13) below. As usual, the difficult part is the treatment of the critical points. However, a quite surprising computation in the spirit of Theorem 4.2 shows that it is always possible to deduce the second effective inequality.

Theorem 4.4. (Effective Monotonicity Formula – II) *For any $1 < p < n$, let g and φ be defined by (3.5) and (3.6) and let $0 < s_p < S_p < +\infty$ be such that $\text{Crit}(\varphi) \subset \{s_p < \varphi < S_p\}$. Then, for every $0 < \lambda < 1$ and every $0 \leq s \leq s_p < S_p \leq S$, the inequality*

$$\begin{aligned} & \left(\frac{e^{\frac{(n-p)s}{(n-2)(p-1)}} - \lambda}{e^{\frac{(n-p)s}{(n-2)(p-1)}}} \right) \Phi'_p(s) - \frac{(n-p)\Phi_p(s)}{(n-2)(p-1)} \\ & \leq \left(\frac{e^{\frac{(n-p)S}{(n-2)(p-1)}} - \lambda}{e^{\frac{(n-p)S}{(n-2)(p-1)}}} \right) \Phi'_p(S) - \frac{(n-p)\Phi_p(S)}{(n-2)(p-1)} \end{aligned} \quad (4.13)$$

holds true, where Φ_p is the function defined in (3.9). In particular, one has that $\Phi_p(+\infty) \leq \Phi_p(0)$.

Proof. Let $\chi : [0, +\infty) \rightarrow \mathbb{R}$ be the same smooth nonnegative cut-off function as in the proof of Theorem 4.2, so that the properties (4.9) are in force. To simplify the notation, let us also set

$$\eta_\lambda(\varphi) = \frac{1}{e^{\frac{(n-p)\varphi}{(n-2)(p-1)}} - \lambda}.$$

Finally, let us consider the smooth vector field

$$\tilde{Y}_\lambda = \chi(\eta_\lambda(\varphi) |\nabla\varphi|) Y_\lambda,$$

where Y_λ has been defined in (4.11). Again, choosing ε small enough, we can suppose $\tilde{Y}_\lambda = Y_\lambda$ on $\{\varphi = s\}$ and $\{\varphi = S\}$, with s and S as in the statement. Hence, applying the Divergence Theorem to the smooth vector field \tilde{Y}_λ on the region $\{s < \varphi < S\}$ gives

$$\begin{aligned} & \int_{\{\varphi=S\}} \left\langle Y_\lambda \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma - \int_{\{\varphi=s\}} \left\langle Y_\lambda \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right\rangle d\sigma = \int_{\{s < \varphi < S\}} \text{div} \tilde{Y}_\lambda d\mu \\ & = \int_{\{s < \varphi < S\} \setminus U_{\varepsilon/2}} \chi(\eta_\lambda(\varphi) |\nabla\varphi|) \text{div} Y_\lambda d\mu \\ & \quad + \int_{U_{3\varepsilon/2} \setminus U_{\varepsilon/2}} \dot{\chi}(\eta_\lambda(\varphi) |\nabla\varphi|) \left\langle Y_\lambda \mid \nabla(\eta_\lambda(\varphi) |\nabla\varphi|) \right\rangle d\mu, \end{aligned}$$

where this time the tubular neighbourhoods of $\text{Crit}(\varphi)$ are defined, for every $\delta > 0$, as $U_\delta = \{\eta_\lambda(\varphi) |\nabla\varphi| \leq \delta\}$. Since, as observed in Lemma 4.3, the divergence of Y_λ is nonnegative on $\{s \leq \varphi \leq S\} \setminus U_{\varepsilon/2}$, where clearly $|\nabla\varphi| \neq 0$, the inequality (4.13) is proved if we can show that

$$\left\langle Y_\lambda \mid \nabla(\eta_\lambda(\varphi) |\nabla\varphi|) \right\rangle \geq 0$$

on $U_{3\varepsilon/2} \setminus U_{\varepsilon/2}$. A direct – though not immediately evident – computation, combined with the definition (4.11) of Y_λ yields

$$\begin{aligned}
 & \left\langle Y_\lambda \left| \nabla \left(\frac{|\nabla\varphi|}{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} - \lambda} \right) \right. \right\rangle \\
 &= e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p-2} \left[|\nabla^\top |\nabla\varphi||^2 + (p-1) |\nabla^\perp |\nabla\varphi||^2 \right] \\
 &\quad - 2 \left(\frac{n-p}{n-2} \right) \eta_\lambda(\varphi) |\nabla\varphi|^p \left\langle \nabla |\nabla\varphi| \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle + \\
 &\quad + \left(\frac{n-p}{n-2} \right)^2 \eta_\lambda^2(\varphi) \frac{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p+2}}{p-1} \\
 &= e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p-2} |\nabla^\top |\nabla\varphi||^2 \\
 &\quad + \left[\left(\frac{n-p}{n-2} \right) \eta_\lambda(\varphi) \left(\frac{e^{\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p+2}}{(p-1)} \right)^{1/2} \right. \\
 &\quad \left. - \left\langle \nabla |\nabla\varphi| \left| \frac{\nabla\varphi}{|\nabla\varphi|} \right. \right\rangle \left((p-1) e^{-\frac{(n-p)}{(n-2)(p-1)}\varphi} |\nabla\varphi|^{p-2} \right)^{1/2} \right]^2
 \end{aligned}$$

This completes the proof of the first part of the statement, since the rightmost hand side is manifestly nonnegative. It remains to show that $\Phi_p(+\infty) \leq \Phi_p(0)$. Applying the inequality (4.13) with $0 < \lambda < 1$, $s = 0$ and $S_p \leq S$, we get

$$\begin{aligned}
 \frac{(n-p)}{(n-2)(p-1)} \left(\Phi_p(S) - \Phi_p(0) \right) &\leq -(1-\lambda) \Phi'_p(0) \\
 &\quad + \left(\frac{e^{\frac{(n-p)}{(n-2)(p-1)}S} - \lambda}{e^{\frac{(n-p)}{(n-2)(p-1)}S}} \right) \Phi'_p(S).
 \end{aligned}$$

Observe now that (4.8) holds also for $S_p < s < S$ (the cut-off argument is not even necessary in this case). Then, the very same reasoning employed to deduce that $\Phi'_p(0) \leq 0$ gives also $\Phi'_p(s) \leq 0$ for any $s > S_p$. In particular, Φ_p is a definitely bounded monotone function, and this implies $\liminf_{S \rightarrow +\infty} \Phi'_p(S) \leq 0$. Hence, passing to the (inferior) limit as $S \rightarrow +\infty$ in the above inequality yields

$$\frac{(n-p)}{(n-2)(p-1)} \left(\lim_{S \rightarrow +\infty} \Phi_p(S) - \Phi_p(0) \right) \leq -(1-\lambda) \Phi'_p(0).$$

Letting $\lambda \rightarrow 1^-$ on the right hand side leads to the second effective inequality $\Phi_p(+\infty) \leq \Phi_p(0)$. \square

4.3. Completion of the proof of Theorem 1.2

We are finally in the position to complete the proof of the L^p -Minkowski inequality, together with the related rigidity statement.

Proof of Theorem 1.2. To obtain inequality (1.4), it is sufficient to detail the proof sketched in Sect. 2.2. As already observed in (3.10), the effective inequalities obtained in Theorems 4.2 and 4.4 correspond to $U'_p(1) \geq 0$ and $U_p(0^+) \leq U_p(1)$, respectively. The first effective inequality implies that

$$\int_{\partial\Omega} \left(\frac{p-1}{n-p} \right) |\mathrm{D}u|^p \, \mathrm{d}\sigma \leq \int_{\partial\Omega} |\mathrm{D}u|^{p-1} \frac{\mathrm{H}}{n-1} \, \mathrm{d}\sigma,$$

since a direct computation shows that

$$U'_p(\tau) = \frac{1}{p-1} \tau^{-\frac{n-1}{n-p}} \int_{\{u=\tau\}} |\mathrm{D}u|^{p-1} \left[\mathrm{H} - \frac{(p-1)(n-1)}{(n-p)} |\mathrm{D} \log u| \right] \, \mathrm{d}\sigma.$$

Applying the Hölder inequality to the above right hand side, with conjugate exponents $a = p/(p-1)$ and $b = p$, one is left with

$$\int_{\partial\Omega} |\mathrm{D}u|^p \, \mathrm{d}\sigma \leq \left(\frac{n-p}{p-1} \right)^p \int_{\partial\Omega} \left| \frac{\mathrm{H}}{n-1} \right|^p \, \mathrm{d}\sigma. \quad (4.14)$$

Using the second effective inequality $U_p(0^+) \leq U_p(1)$ in combination with (3.4) we get

$$\left(\frac{n-p}{p-1} \right)^p |\mathbb{S}^{n-1}| C_p(\Omega)^{\frac{n-p-1}{n-p}} = \lim_{\tau \rightarrow 0^+} U_p(\tau) \leq U_p(1) = \int_{\partial\Omega} |\mathrm{D}u|^p \, \mathrm{d}\sigma,$$

that combined with (4.14) gives the desired

$$C_p(\Omega)^{\frac{n-p-1}{n-p}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathrm{H}}{n-1} \right|^p \, \mathrm{d}\sigma.$$

Assume now that equality holds in (1.2). Then, equality holds in (4.14), and consequently $U'_p(1) = \Phi'_p(0) = 0$. Let $s^* \in (0, +\infty]$ be the first critical value of φ . A straightforward perusal of the proof of Theorem 4.2 shows that $\operatorname{div}_g X = 0$ in $\{\varphi \leq s\}$ for any $s < s^*$. By (4.3) and (3.7) we deduce that $|\nabla \nabla \varphi|_g = 0$ in this region. Then, a very standard argument (see e.g. the proof of [2, Theorem 4.1 (i)]) shows that $(\{\varphi \leq s\}, g)$ is isometric to the cylinder $([0, s] \times \{\varphi = 0\}, dt \otimes dt + g_{\{\varphi=0\}})$, and that $|\nabla \varphi|_g$ equals a (positive) constant in this region. The existence of a critical value $s^* < +\infty$ would thus contradict the continuity of $|\nabla \varphi|_g$, that follows from the \mathcal{C}^1 -regularity of p -harmonic functions. Then, $|\nabla \nabla \varphi|_g = 0$ on the whole $\mathbb{R}^n \setminus \overline{\Omega}$, and then [2, Theorem 4.1 (ii)] implies that $\partial\Omega$ is a sphere. \square

5. Proof of the Extended Minkowski Inequality

In this section we derive the Extended Minkowski Inequality (1.3)

$$\left(\frac{|\partial\Omega^*|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\mathrm{H}}{n-1} \right| \, \mathrm{d}\sigma,$$

by letting $p \rightarrow 1^+$ in the L^p -Minkowski Inequality (1.4). The main task here (see Theorem 5.6) is to compute – and characterise geometrically – the limit of the

variational p -capacity of a bounded set with smooth boundary. As we are going to see, this limit turns out to be related to the *strictly outward minimising hull* of Ω , a notion that plays a central role in the formulation of the weak Inverse Mean Curvature Flow introduced in [37].

5.1. (Strictly) outward minimising sets and the strictly outward minimising hull

The notion of *outward minimising sets* and *strictly outward minimising sets* are given in the context of sets with finite perimeter. We refer the reader to [48] for a comprehensive treatment of the basic notions that we are going to recall.

Definition 5.1. (*Outward minimising and strictly outward minimising sets*) Let $E \subset \mathbb{R}^n$ be a bounded measurable set with finite perimeter. We say that E is *outward minimising* if for any $F \subset \mathbb{R}^n$ with $E \subseteq F$ we have $|\partial^* E| \leq |\partial^* F|$, where by $\partial^* F$ we denote the reduced boundary of a set F . We say that E is *strictly outward minimising* if it is outward minimising and any time $|\partial^* E| = |\partial^* F|$ for some $F \subset \mathbb{R}^n$ with $E \subseteq F$ we have $|F \setminus E| = 0$.

It is easily seen that a bounded open set with finite perimeter is (strictly) outward minimising if and only if any measure zero modification of it is (strictly) outward minimising. In order to define appropriate representatives for these sets, we recall the definition of the *measure theoretic interior* of a set E with $|E| < +\infty$ as the points of density 1 for E , namely

$$\text{Int}(E) = \left\{ x \in \mathbb{R}^n \mid \lim_{r \rightarrow 0^+} \frac{|E \cap B(x, r)|}{|B(x, r)|} = 1 \right\}.$$

It follows from Lebesgue Differentiation Theorem (see [48, Theorem 5.16]) that

$$|E \Delta \text{Int}(E)| = 0.$$

Importantly, a set with finite perimeter E satisfies,

$$\partial \text{Int}(E) = \overline{\partial^* E}, \tag{5.1}$$

that is, the topological boundary of the measure theoretic interior of a set with finite perimeter coincides with the closure of its reduced boundary. We address the reader to [11, Theorem 10] for a proof of this nice property. We are now ready to define the *strictly outward minimising hull* of a set. As it can be checked, this concept essentially coincides with the one outlined in [37, p. 371].

Definition 5.2. (*Strictly outward minimising hull*) Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary. We define the *strictly outward minimising hull* of Ω as the measure theoretic interior of a set E solving the minimisation problem

$$\inf_{E \in \text{SOMBE}(\Omega)} |E|,$$

where

$$\text{SOMBE}(\Omega) = \{ E \mid \Omega \subseteq E \text{ and } E \text{ is bounded and strictly outward minimising} \}.$$

According to [25, Theorem 1.1], every bounded subset $\Omega \subset \mathbb{R}^n$ with smooth boundary admits a bounded strictly outward minimising hull Ω^* , which is characterised as a maximal volume solution to the least area problem with obstacle Ω . In particular, it satisfies

$$|\partial^* \Omega^*| = \inf \{ |\partial^* F| \mid \Omega \subseteq F \}. \quad (5.2)$$

We recall that the main result in [58] (see also the comprehensive [37, Theorem 1.3]) provides us with a regularity result for any solution E to (5.2) such that $\partial E = \overline{\partial^* E}$. Note that Ω^* fulfils this requirement (in view of (5.1) and the fact that $\text{Int}(\Omega^*) = \Omega^*$, by the very definition of Ω^*). The topological boundary $\partial \Omega^*$ is thus equipped with the following regularity.

Theorem 5.3. (Regularity of the strictly outward minimising hull) *Let $\Omega \subset \mathbb{R}^n$ be a bounded set with smooth boundary. Then*

- (i) $\partial \Omega^*$ is a $\mathcal{C}^{1,1}$ hypersurface in a neighbourhood of any point in $\partial \Omega^* \cap \partial \Omega$.
- (ii) $\partial \Omega^*$ is area minimising in $\partial \Omega^* \setminus \partial \Omega$. In particular there exists a singular set $\text{Sing} \subset \partial \Omega^* \setminus \partial \Omega$, with Hausdorff dimension at most $n - 8$, such that $\partial \Omega^* \setminus \partial \Omega$ is a real analytic hypersurface in a neighbourhood of any point in $(\partial \Omega^* \setminus \partial \Omega) \setminus \text{Sing}$.

An obvious consequence of this theorem is the fact that $|\partial^* \Omega^*| = |\partial \Omega^*|$.

Remark 5.4. Since, as already pointed out, the boundary of Ω^* has least area among sets enclosing Ω , we have that a bounded open set $\Omega \subset \mathbb{R}^n$ with smooth boundary is outward minimising if and only if

$$|\partial \Omega| = |\partial \Omega^*|.$$

Observe that the inequality $|\partial \Omega^*| \leq |\partial \Omega|$ is automatically satisfied, due to (5.2).

Since $|\partial^* \Omega^*| = |\partial \Omega^*|$ we can apply the nice interior approximation result [57, Theorem 1.1] to $B(x, R) \setminus \overline{\Omega^*}$, with $\Omega^* \Subset B(x, R)$, to obtain the following exterior approximation result.

Lemma 5.5. (Smooth exterior approximation of the strictly outward minimising hull) *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary. Then, there exists a sequence of bounded sets $\{\Omega_k\}_{k \in \mathbb{N}}$ with smooth boundary such that*

$$\Omega^* \subset \Omega_k, \quad |\partial \Omega_k| \rightarrow |\partial \Omega^*|.$$

5.2. Minimising hulls and p -capacities

Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary. Recall from Definition 3.1 that for $1 < p < n$ one has

$$\text{Cap}_p(\Omega) = \inf \left\{ \int_{\mathbb{R}^n} |Df|^p \, d\mu \mid f \geq \chi_\Omega, \, f \in \mathcal{C}_c^\infty(\mathbb{R}^n) \right\}. \quad (5.3)$$

We can define, analogously, the 1-capacity of a bounded open set with smooth boundary Ω as

$$\text{Cap}_1(\Omega) = \inf \left\{ \int_{\mathbb{R}^n} |Df| \, d\mu \mid f \geq \chi_\Omega, f \in \mathcal{C}_c^\infty(\mathbb{R}^n) \right\}. \quad (5.4)$$

The following result says that $\text{Cap}_1(\Omega)$ can indeed be recovered as the limit of $\text{Cap}_p(\Omega)$, as $p \rightarrow 1^+$, and that these quantities are also related with the strictly outward minimising hull of Ω .

Theorem 5.6. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with smooth boundary. Then,*

$$\lim_{p \rightarrow 1^+} \text{Cap}_p(\Omega) = |\partial\Omega^*|,$$

where Ω^* is the strictly outward minimising hull of Ω .

Proof. Let us first observe that for any $f \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ with $f \geq \chi_\Omega$ we have, by co-area formula,

$$\int_{\mathbb{R}^n} |Df| \, d\mu \geq \int_0^1 |\{f = t\}| \, dt \geq \inf \left\{ |\partial E| \mid \Omega \subset E, \partial E \text{ smooth} \right\} \geq |\partial\Omega^*|,$$

where the last inequality is due to (5.2). In particular, taking the infimum over any such f , we get

$$|\partial\Omega^*| \leq \text{Cap}_1(\Omega). \quad (5.5)$$

We now prove that

$$\text{Cap}_1(\Omega) \leq \liminf_{p \rightarrow 1^+} \text{Cap}_p(\Omega). \quad (5.6)$$

This will be done by passing to the limit as $p \rightarrow 1^+$ in the inequality appearing in the proof of [67, Theorem 3.2], keeping track of the appearing constants (which results in inequality (5.10) below). Namely, for every $f \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ with $f \geq \chi_\Omega$ and any positive exponent q , the function f^q is an admissible competitor in (5.3) and (5.4). Then, by definition of the 1-capacity and by Hölder inequality we have

$$\begin{aligned} \text{Cap}_1(\Omega) &\leq \int_{\mathbb{R}^n} |Df^q| \, d\mu = q \int_{\mathbb{R}^n} f^{q-1} |Df| \, d\mu \\ &\leq q \left(\int_{\mathbb{R}^n} f^{(q-1)\frac{p}{p-1}} \, d\mu \right)^{(p-1)/p} \left(\int_{\mathbb{R}^n} |Df|^p \, d\mu \right)^{1/p}. \end{aligned} \quad (5.7)$$

Let now q satisfy $(q-1)p/(p-1) = p^*$, where $p^* = pn/(n-p)$ is the Sobolev conjugate exponent of p , that is

$$q = 1 + p^* \frac{(p-1)}{p}. \quad (5.8)$$

Observe that with this choice $q > 1$. Then, we obtain, applying the Sobolev inequality to the first integral in the right hand side of (5.7),

$$\begin{aligned} \text{Cap}_1(\Omega) &\leq q T_{n,p}^{p^*(p-1)/p} \left(\int_{\mathbb{R}^n} |Df|^p \, d\mu \right)^{p^*(p-1)/p^2+1/p} \\ &= q T_{n,p}^{q-1} \left(\int_{\mathbb{R}^n} |Df|^p \, d\mu \right)^{(n-1)/(n-p)}, \end{aligned} \quad (5.9)$$

where $T_{n,p}$ is Talenti's best constant in the Sobolev inequality, obtained in [59]. We recall that the precise value of such constant is

$$T_{n,p} = \frac{1}{\pi^{1/2} n^{1/p}} \left(\frac{p-1}{n-p} \right)^{(p-1)/p} \left[\frac{\Gamma(1+n/2)\Gamma(n)}{\Gamma(n/p)\Gamma(1+n-n/p)} \right]^{1/n},$$

where Γ is Euler's Gamma function. Taking the infimum over any $f \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ with $f \geq \chi_\Omega$ in (5.9), we obtain

$$\text{Cap}_1(\Omega) \leq q T_{n,p}^{q-1} \text{Cap}_p(\Omega)^{(n-1)/(n-p)}. \quad (5.10)$$

As $p \rightarrow 1^+$, one can check that $T_{n,p}$ converges to a positive constant, precisely (compare with [59, p. 355])

$$\lim_{p \rightarrow 1^+} T_{n,p} = \left(\frac{1}{|\mathbb{S}^{n-1}|} \right)^{1/n},$$

and this implies

$$\lim_{p \rightarrow 1^+} T_{n,p}^{q-1} = 1.$$

Note that $q \rightarrow 1$, as $p \rightarrow 1^+$, in view of (5.8). In turn, passing to the limit in (5.10), we get (5.6).

We are left to prove the inequality

$$\limsup_{p \rightarrow 1^+} \text{Cap}_p(\Omega) \leq |\partial\Omega^*|. \quad (5.11)$$

Let E be any open and bounded set in \mathbb{R}^n with smooth boundary such that $\Omega \subset E$. Define, for $x \in \mathbb{R}^n$, the function $d_E(x) = \text{dist}(x, E)$. Moreover, let us introduce a smooth cut-off function χ_ε fulfilling

$$\begin{cases} \chi_\varepsilon(t) = 1 & \text{in } t < \varepsilon, \\ \dot{\chi}_\varepsilon(t) \leq 0 & \text{in } \varepsilon \leq t \leq 2\varepsilon \\ \chi_\varepsilon(t) = 0 & \text{in } t > 2\varepsilon, \end{cases}$$

and let us set $\eta_\varepsilon(x) = \chi_\varepsilon(d_E(x))$. Choosing ε small enough, it is easily seen, by the regularity of d_E in a neighbourhood of E (see [33, Lemma 14.16]), that the function η_ε is an admissible competitor in (5.3) and (5.4). Then,

$$\text{Cap}_p(\Omega) \leq \int_{\mathbb{R}^n} |D\eta_\varepsilon|^p \, d\mu$$

for any $p \geq 1$. Letting $p \rightarrow 1^+$, we get

$$\begin{aligned} \limsup_{p \rightarrow 1^+} \text{Cap}_p(\Omega) &\leq \int_{\mathbb{R}^n} |\text{D}\eta_\varepsilon| \, \text{d}\mu \leq - \sup_{s \in (\varepsilon, 2\varepsilon)} |\{d_E = s\}| \times \int_\varepsilon^{2\varepsilon} \dot{\chi}_\varepsilon(t) \, \text{d}t \\ &= \sup_{s \in (\varepsilon, 2\varepsilon)} |\{d_E = s\}|, \end{aligned}$$

where in the second inequality we have applied the coarea formula. Since, as $\varepsilon \rightarrow 0^+$, we clearly have

$$\sup_{s \in (\varepsilon, 2\varepsilon)} |\{d_E = s\}| \rightarrow |\partial E|,$$

we conclude that

$$\limsup_{p \rightarrow 1^+} \text{Cap}_p(\Omega) \leq |\partial E|$$

for any bounded open set E with smooth boundary containing Ω . In particular, considering a sequence of bounded sets $\{\Omega_k\}_{k \in \mathbb{N}}$ with smooth boundary containing Ω^* and with $|\partial\Omega_k| \rightarrow |\partial\Omega^*|$ as $k \rightarrow \infty$, provided in Lemma 5.5, we get (5.11). The inequalities (5.5), (5.6) and (5.11) combine as

$$|\partial\Omega^*| \leq \text{Cap}_1(\Omega) \leq \liminf_{p \rightarrow 1^+} \text{Cap}_p(\Omega) \leq \limsup_{p \rightarrow 1^+} \text{Cap}_p(\Omega) \leq |\partial\Omega^*|,$$

completing the proof. □

5.3. Proof of Theorem 1.1 and Corollary 1.3

We are now in the position to prove Theorem 1.1 together with its Corollary 1.3.

Proof of Theorem 1.1. It suffices to pass to the limit as $p \rightarrow 1^+$ in (1.4), that is

$$\text{C}_p(\Omega)^{\frac{n-p-1}{n-p}} \leq \frac{1}{|\mathbb{S}^{n-1}|} \int_{\partial\Omega} \left| \frac{\text{H}}{n-1} \right|^p \, \text{d}\sigma. \quad (5.12)$$

Indeed, recalling the relation between p -capacity and normalised p -capacity given in Definition 3.1, Theorem 5.6 shows that the left hand side of the above inequality behaves as

$$\lim_{p \rightarrow 1^+} \text{C}_p(\Omega)^{\frac{n-p-1}{n-p}} = \left(\frac{|\partial\Omega^*|}{|\mathbb{S}^{n-1}|} \right)^{\frac{n-2}{n-1}},$$

while the right-hand side of (5.12) is immediately seen to converge to the right hand side of (1.3). Spheres show the optimality of the estimate since their mean curvature is given by $(n-1)/R$, where R is the radius of the ball they enclose. □

Proof Corollary 1.3. Inequality (1.5) immediately follows from the fact that outward minimising sets with smooth boundary satisfy $|\partial\Omega^*| = |\partial\Omega|$ (see Remark 5.4) and the mean curvature of their boundaries (computed with respect to the outward unit normal) is nonnegative, for otherwise one might easily provide compactly supported outward perturbations of $\partial\Omega$ that decrease the area. To see this, it is sufficient to recall that the first variation of $|\partial\Omega|$ with respect to the variation $\psi\nu$, where $\psi \in C_c^\infty(\partial\Omega)$ is nonnegative and ν is the outward unit normal, is given by

$$\int_{\partial\Omega} H\psi \, d\sigma .$$

To complete the proof of Corollary 1.3, we are left to consider the equality case in (1.5) for some strictly outward minimising set with smooth and strictly mean-convex boundary. To this aim, let $\{\partial\Omega_t\}_{t \in [0, T]}$ be the evolution of $\partial\Omega$ under smooth IMCF, up to some $T > 0$. By [37, Lemma 2.4], the weak IMCF starting at $\partial\Omega$ coincides with the smooth flow $\{\Omega_t\}_{t \in [0, T^*)}$, for some $0 < T^* \leq T$. In particular, by [37, Lemma 1.4], Ω_t is strictly outward minimising and strictly mean-convex for every $t \in [0, T^*)$, and then (1.5) holds for every $\partial\Omega_t$ with $t \in [0, T^*)$. We can then define, for $t \in [0, T^*)$, the monotonic quantity already discussed in Sect. 2.1, namely

$$\mathcal{Q}(t) = |\partial\Omega_t|^{-\frac{n-2}{n-1}} \int_{\partial\Omega_t} H \, d\sigma .$$

Observe that inequality (1.5) is equivalent to $\mathcal{Q}(0) \geq (n-1)|\mathbb{S}^{n-1}|^{1/(n-1)}$, and assuming equality in (1.5) is equivalent to $\mathcal{Q}(0) = (n-1)|\mathbb{S}^{n-1}|^{1/(n-1)}$. By the smoothness of the flow, the function $\mathcal{Q}(t)$ is differentiable for $t \in [0, T)$, and then a straightforward computation involving the standard evolution equations provided e.g. in [39, Theorem 3.2] shows that

$$\mathcal{Q}'(0) = -|\partial\Omega|^{-\frac{n-2}{n-1}} \int_{\partial\Omega} \frac{|\mathring{h}|^2}{H} \, d\sigma \leq 0 . \tag{5.13}$$

However, since we assumed $\mathcal{Q}(0) = (n-1)|\mathbb{S}^{n-1}|^{1/(n-1)}$, the strict inequality $\mathcal{Q}'(0) < 0$ would imply $\mathcal{Q}(t) < (n-1)|\mathbb{S}^{n-1}|^{1/(n-1)}$ for some $t \in (0, T^*)$, which is equivalent to contradict (1.5) for some outward minimising Ω_t with strictly mean-convex boundary. Then $\mathcal{Q}'(0) = 0$ and in turn, by formula (5.13), $\partial\Omega$ is totally umbilical. Therefore, $\partial\Omega$ must be a sphere. \square

6. Optimal Nearly Umbilical Estimates for Outward Minimising Sets

This section is devoted to the proof of an optimal version of the well celebrated De Lellis–Müller nearly umbilical estimates for outward minimising domains.

Theorem 6.1. (Optimal Nearly Umbilical Estimate) *If $\Omega \subset \mathbb{R}^3$ is a bounded outward minimising open domain with smooth boundary, then*

$$\int_{\partial\Omega} \left| h - \frac{\bar{H}}{2} g_{\partial\Omega} \right|^2 d\sigma \leq 2 \int_{\partial\Omega} |\mathring{h}|^2 d\sigma, \quad (6.1)$$

where $g_{\partial\Omega}$ is the metric induced on $\partial\Omega$ by the Euclidean metric of \mathbb{R}^3 , and

$$\bar{H} = \int_{\partial\Omega} H d\sigma, \quad \mathring{h} = h - \frac{H}{2} g_{\partial\Omega}.$$

Moreover, the equality is achieved in (6.1) by some strictly mean-convex and strictly outward minimising domain Ω if and only if Ω is isometric to a round ball.

A first main tool we are going to use in order to deduce Theorem 6.1 from Theorem 1.1 is the classical Gauss' equation for surfaces in \mathbb{R}^3 , yielding

$$R_{\partial\Omega} = H^2 - |h^2|, \quad (6.2)$$

where $R_{\partial\Omega}$ is the scalar curvature of $\partial\Omega$ computed with respect to the metric $g_{\partial\Omega}$ induced on it by the Euclidean metric of \mathbb{R}^3 . A second main tool we need to recall is the famous Gauss–Bonnet formula, stating that

$$\int_{\partial\Omega} R_{\partial\Omega} d\sigma = 4\pi \chi(\partial\Omega), \quad (6.3)$$

where $\chi(\partial\Omega)$ is the Euler characteristic of the surface $\partial\Omega$.

We are finally going to show how the Minkowski inequality (1.1) combined with these basic identities in differential geometry gives the optimal nearly umbilical estimate (6.1).

Proof. (Proof of Theorem 6.1) Expanding the squares, it is straightforwardly seen that (6.1) is equivalent to

$$\int_{\partial\Omega} (|H|^2 - |h|^2) d\sigma \leq \frac{\bar{H}^2}{2} |\partial\Omega|.$$

Invoking Gauss' equation (6.2) and Gauss–Bonnet formula (6.3), we then obtain that (6.1) is equivalent to

$$8\pi \chi(\partial\Omega) = 2 \int_{\partial\Omega} R_{\partial\Omega} d\sigma = 2 \int_{\partial\Omega} (|H|^2 - |h|^2) d\sigma \leq \bar{H}^2 |\partial\Omega|,$$

that is, to

$$\sqrt{2\pi \chi(\partial\Omega) |\partial\Omega|} \leq \int_{\partial\Omega} \frac{H}{2} d\sigma. \quad (6.4)$$

Since obviously $\chi(\partial\Omega) \leq 2$, the inequality (6.4) follows from the Minkowski inequality (1.5).

Assume now that equality holds for some outward minimising set Ω with smooth and strictly mean-convex boundary. Let $\{\partial\Omega_t\}_{t \in [0, T]}$ be evolving by IMCF with initial datum $\partial\Omega$. By [37, Lemma 2.4], the weak IMCF $\{E_t\}_{t \in [0, \infty)}$ starting at $\partial\Omega$ coincides with Ω_t for $t \in [0, T^*)$, for some T^* possibly smaller than T . In particular, Ω_t is outward minimising and strictly mean-convex for any $t \in [0, T^*)$, for some $T^* > 0$, and then (6.1) holds for $\partial\Omega_t$ for any $t \in [0, T^*)$. We can then define, for $T \in [0, T^*)$, the quantity

$$\mathcal{P}(t) = \int_{\partial\Omega_t} |\mathring{h}|^2 \, d\sigma - \frac{1}{2} \int_{\partial\Omega_t} \left(H - \frac{1}{|\partial\Omega_t|} \int_{\partial\Omega_t} H \right)^2 \, d\sigma,$$

introduced in [54, Chapter 3]. Observe that inequality (6.1) is equivalent to $\mathcal{P}(0) \geq 0$, assuming equality in (6.1) is equivalent to $\mathcal{P}(0) = 0$. By the smoothness of the flow, the function $\mathcal{P}(t)$ is differentiable for $t \in [0, T)$, and then [54, Lemma 3.4] yields

$$\mathcal{P}'(0) = -\bar{H} \int_{\partial\Omega} \frac{|\mathring{h}|^2}{H} \, d\sigma \leq 0. \tag{6.5}$$

However, since we assumed $\mathcal{P}(0) = 0$, $\mathcal{P}'(0) < 0$ would imply $\mathcal{P}(t) < 0$ for some $t \in (0, T^*)$ that is equivalent to falsify (6.1) for some outward minimising Ω_t with strictly mean-convex boundary. Then $\mathcal{P}'(0) = 0$, and by formula (6.5) this means that $\partial\Omega$ is totally umbilical, thus a sphere. \square

Inequality (6.4) in the above proof, that we just showed to be equivalent to the nearly umbilical estimate (6.1), actually coincides with the Minkowski inequality for mean-convex hypersurfaces (1.5) if $\chi(\partial\Omega) = 2$, that is, if $\partial\Omega$ is diffeomorphic to a sphere. We want to show, with the following easy proposition, that such a diffeomorphism exists each time the right-hand side of (6.1) is smaller than 16π .

Proposition 6.2. *Let $\Omega \subset \mathbb{R}^3$ be a bounded open set with smooth and mean-convex boundary. If*

$$\int_{\partial\Omega} \left| \mathring{h} - \frac{H}{2} g_{\partial\Omega} \right|^2 \, d\sigma \leq 8\pi \tag{6.6}$$

then $\partial\Omega$ is diffeomorphic to a sphere. In particular, if (6.6) holds, then the Minkowski inequality is equivalent to the optimal nearly umbilical estimate.

Proof. If (6.6) holds, we have, by the classical Willmore inequality [65],

$$\int_{\partial\Omega} \left| \mathring{h} - \frac{H}{2} g_{\partial\Omega} \right|^2 \, d\sigma \leq 8\pi \leq \frac{1}{2} \int_{\partial\Omega} H^2 \, d\sigma,$$

which implies

$$\int_{\partial\Omega} (|H|^2 - |\mathring{h}|^2) \, d\sigma \geq 0. \tag{6.7}$$

Moreover, since equality is attained in the Willmore inequality if and only if $\partial\Omega$ is isometric to a sphere with the round metric, the same rigidity statement holds if

equality is attained in (6.7). On the other hand, applying the Gauss' equation (6.2) and the Gauss–Bonnet formula (6.3), (6.7) is equivalent to

$$\chi(\partial\Omega) \geq 0.$$

If $\chi(\partial\Omega) = 0$ then by the characterisation of the equality case in the Willmore inequality $\partial\Omega$ would be even isometric to a sphere, and this is a contradiction. Then $\chi(\partial\Omega) = 2$, as claimed. \square

Acknowledgements. The authors are grateful to G. De Philippis, N. Fusco, G. Huisken, A. Malchiodi, C. Mantegazza, L. Mari, J. Scheuer and F. Schulze for their interest in the present work, as well as for useful comments and discussions during the preparation of the manuscript. A special thank goes to L. Benatti and F. Oronzio for the careful reading of this paper. The authors are members of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA), which is part of the Istituto Nazionale di Alta Matematica (INdAM), and they are partially funded by the GNAMPA project "Aspetti geometrici in teoria del potenziale lineare e nonlineare".

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. AGOSTINIANI, V., FOGAGNOLO, M., MAZZIERI, L.: Sharp geometric inequalities for closed hypersurfaces in manifolds with nonnegative Ricci curvature. *Invent. Math.* **222**, 1033–1101, 2020
2. AGOSTINIANI, V., MAZZIERI, L.: Riemannian aspects of potential theory. *J. Math. Pures Appl.* **104**(3), 561–586, 2015
3. AGOSTINIANI, V., MAZZIERI, L.: On the geometry of the level sets of bounded static potentials. *Commun. Math. Phys.* **355**, 261–301, 2017
4. AGOSTINIANI, V., MAZZIERI, L.: Monotonicity formulas in potential theory. *Calc. Var. Partial Differ. Equ.* **59**(1), 6, 2019
5. ALBERTI, G., BIANCHINI, S., CRIPPA, G.: Structure of level sets and Sard-type properties of Lipschitz maps. *Annali della Scuola Normale Superiore di Pisa. Classe di Scienze. Serie V*, 4, 2011
6. ALEKSANDROV, A.D.: Zur Theorie der gemischten Volumina von konvexen Körpern, II. Neue Ungleichungen zwischen den gemischten Volumina und ihre Anwendungen. *Mat. Sbornik N.S.* **2**, 1205–1238, 1937 (in Russian)
7. ALEKSANDROV, A.D.: Zur Theorie der gemischten Volumina von konvexen Körpern, III. Die Erweiterung zweier Lehrsätze Minkowskis über die konvexen Polyeder auf beliebige konvexe Flächen. *Mat. Sbornik N.S.* **3**, 27–46, 1938 (in Russian)

8. BRAY, H.L., MIAO, P.: On the capacity of surfaces in manifolds with nonnegative scalar curvature. *Invent. Math.* **172**(3), 459–475, 2008
9. BRAY, H.L., NEVES, A.: Classification of prime 3-manifolds with σ -invariant greater than $\mathbb{R}P^3$. *Ann. Math.* **159**(1), 407–424, 2004
10. BRENDLE, S., HUNG, P.-K., WANG, M.-T.: A Minkowski inequality for hypersurfaces in the Anti-de Sitter–Schwarzschild manifold. *Commun. Pure Appl. Math.* **69**(1), 124–144, 2016
11. CARABALLO, D.G.: Local simplicity, topology, and sets of finite perimeter. *Interfaces Free Bound.* **13**(2), 171–189, 2011
12. CHAMBOLLE, A., MORINI, M., NOVAGA, M., PONSIGLIONE, M.: Existence and uniqueness for anisotropic and crystalline mean curvature flows. *J. Am. Math. Soc.* **32**(3), 779–824, 2019
13. CHAMBOLLE, A., MORINI, M., PONSIGLIONE, M.: Nonlocal curvature flows. *Arch. Ration. Mech. Anal.* **218**(3), 1263–1329, 2015
14. CHANG, S.-Y.A., WANG, Y.: On Aleksandrov–Fenchel Inequalities for k -Convex Domains. *Milan J. Math.* **79**(1), 13, 2011
15. CHANG, S.-Y.A., WANG, Y.: Some higher order isoperimetric inequalities via the method of optimal transport. *Int. Math. Res. Not.* **2014**, 05, 2013
16. COLDING, T.H.: New monotonicity formulas for Ricci curvature and applications. I. *Acta Math.* **209**(2), 229–263, 2012
17. COLDING, T.H., MINICOZZI, W.P.: Monotonicity and its analytic and geometric implications. *Proc. Natl. Acad. Sci.* **110**(48), 19233–19236, 2013
18. COLDING, T.H., MINICOZZI, W.P.: Ricci curvature and monotonicity for harmonic functions. *Calc. Var. Partial Differ. Equ.* **49**(3), 1045–1059, 2014
19. COLESANTI, A., SALANI, P.: The Brunn–Minkowski inequality for p -capacity of convex bodies. *Math. Ann.* **327**(3), 459–479, 2003
20. DE LELLIS, C., MÜLLER, S.: Optimal rigidity estimates for nearly umbilical surfaces. *J. Differ. Geom.* **69**(1), 75–110, 2005
21. DE LIMA, L.L., GIRÃO, F.: An Alexandrov–Fenchel-type Inequality in Hyperbolic Space with an application to a Penrose Inequality. *Ann. Henri Poincaré* **17**(4), 979–1002, 2016
22. DiBENEDETTO, E.: $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. *Nonlinear Anal.* **7**(8), 827–850, 1983
23. EVANS, L.C.: A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic p.d.e. *J. Differ. Equ.* **45**(3), 356–373, 1982
24. FENCHEL, W.: Über Krümmung und Windung geschlossener Raumkurven. *Math. Ann.* **101**, 238–252, 1929
25. FOGAGNOLO, M., MAZZIERI, L.: Minimising hulls, p -capacity and isoperimetric inequality on complete Riemannian manifolds. [arXiv:2012.09490](https://arxiv.org/abs/2012.09490)
26. FOGAGNOLO, M., MAZZIERI, L., PINAMONTI, A.: Geometric aspects of p -capacitary potentials. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **36**(4), 1151–1179, 2019
27. FREIRE, A., SCHWARTZ, F.: Mass-capacity inequalities for conformally flat manifolds with boundary. *Commun. Partial Differ. Equ.* **39**(1), 98–119, 2014
28. GAROFALO, N., SARTORI, E.: Symmetry in exterior boundary value problems for quasilinear elliptic equations via blow-up and a priori estimates. *Adv. Differ. Equ.* **4**(2), 137–161, 1999
29. GE, Y., WANG, G., WU, J.: Hyperbolic Alexandrov–Fenchel quermassintegral inequalities i. *J. Differ. Geom.* **98**, 03, 2013
30. GE, Y., WANG, G., WU, J.: Hyperbolic Alexandrov–Fenchel quermassintegral inequalities ii. *J. Differ. Geom.* **98**(2), 237–260, 2014
31. GE, Y., WANG, G., WU, J., XIA, C.: A Penrose inequality for graphs over Kottler space. *Calc. Var. Partial Differ. Equ.* **52**(3), 755–782, 2015
32. GERHARDT, C.: Flow of nonconvex hypersurfaces into spheres. *J. Differ. Geom.* **32**(1), 299–314, 1990
33. GILBARG, D., TRUDINGER, N. S.: *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition

34. GUAN, P., LI, J.: The quermassintegral inequalities for k -convex starshaped domains. *Adv. Math.* **221**(5), 1725–1732, 2009
35. HARDT, R., SIMON, L.: Area minimizing hypersurfaces with isolated singularities. *Journal für die Reine und Angewandte Mathematik [Crelle's Journal]* **362**, 102–129, 1985
36. HUISKEN, G.: An isoperimetric concept for the mass in general relativity. Video, available at <https://www.youtube.com/watch?v=Qt09iMPUCYY>
37. HUISKEN, G., ILMANEN, T.: The inverse mean curvature flow and the Riemannian Penrose inequality. *J. Differ. Geom.* **59**(3), 353–437, 2001
38. HUISKEN, G., ILMANEN, T.: Higher regularity of the inverse mean curvature flow. *J. Differ. Geom.* **80**(3), 433–451, 2008
39. HUISKEN, G., POLDEN, A.: Geometric evolution equations for hypersurfaces. In *Calculus of variations and geometric evolution problems (Cetraro, 1996)*, volume 1713 of *Lecture Notes in Math.*, pages 45–84. Springer, Berlin, 1999
40. KICHENASSAMY, S., VÉRON, L.: Singular solutions of the p -Laplace equation. *Math. Ann.* **275**(4), 599–615, 1986
41. KOTSCHWAR, B., NI, L.: Local gradient estimates of p -harmonic functions $1/H$ -flow and an entropy formula. *Ann. Sci. Éc. Norm. Supér. (4)* **42**(1), 1–36, 2009
42. LADYZHENSKAYA, O.A., URAL'TSEVA, N.N.: *Linear and quasilinear elliptic equations*. Translated from the Russian by Scripta Technica, Inc. Translation editor: Leon Ehrenpreis. Academic Press, New York, 1968
43. LEE, A.D., NEVES, A.: The Penrose inequality for asymptotically locally hyperbolic spaces with nonpositive mass. *Commun. Math. Phys.* **327**, 339–352, 2015
44. LEWIS, J.L.: Regularity of the derivatives of solutions to certain degenerate elliptic equations. *Indiana Univ. Math. J.* **32**(6), 849–858, 1983
45. LI, H., WEI, Y.: On inverse mean curvature flow in Schwarzschild space and Kottler space. *Calc. Var. Partial Differ. Equ.* **56**(3), 62, 2017
46. LIEBERMAN, G.M.: Boundary regularity for solutions of degenerate elliptic equations. *Nonlinear Anal.* **12**(11), 1203–1219, 1988
47. LINDQVIST, P.: Notes on the p -Laplace equation, 2nd edn. University of Jyväskylä, Jyväskylä (2017)
48. MAGGI, F.: *Sets of Finite Perimeter and Geometric Variational Problems: An Introduction to Geometric Measure Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge (2012)
49. MAKOWSKI, M., SCHEUER, J.: Rigidity results, inverse curvature flows and Alexandrov–Fenchel type inequalities in the sphere. *Asian J. Math.* **20**, 869–892, 2016
50. MARI, L., RIGOLI, L., SETTI, A.G.: On the $1/h$ -flow by p -Laplace approximation: new estimates via fake distances under Ricci lower bounds. [arXiv:1905.00216](https://arxiv.org/abs/1905.00216)
51. MCCORMICK, S.: On a Minkowski-like inequality for asymptotically flat static manifolds. *Proc. Am. Math. Soc.* **146**, 09, 2017
52. MINKOWSKI, H.: Volumen und Oberfläche. *Math. Ann.* **57**(4), 447–495, 1903
53. MOSER, R.: The inverse mean curvature flow and p -harmonic functions. *J. Eur. Math. Soc.* **9**(1), 77–83, 2007
54. PEREZ, D.R.: *On nearly umbilical hypersurfaces*. Ph.D. thesis, University of Zurich, 2011
55. POGGESI, G.: Radial symmetry for p -harmonic functions in exterior and punctured domains. *Appl. Anal.*, 1–14, 2018
56. QIU, G.: A family of higher-order isoperimetric inequalities. *Commun. Contemp. Math.* **17**(03), 1450015, 2015
57. SCHMIDT, T.: Strict interior approximation of sets of finite perimeter and functions of bounded variation. *Proc. Am. Math. Soc.* **143**(5), 2069–2084, 2015
58. STERNBERG, P., ZIEMER, W.P., WILLIAMS, G.: $C^{1,1}$ -regularity of constrained area minimizing hypersurfaces. *J. Differ. Equ.* **94**(1), 83–94, 1991
59. TALENTI, G.: Best constant in Sobolev inequality. *Ann. Mat. Pura Appl.* **4**(110), 353–372, 1976

60. TOLKSDORF, P.: On the Dirichlet problem for quasilinear equations in domains with conical boundary points. *Commun. Partial Differ. Equ.* **8**(7), 773–817, 1983
61. TRUDINGER, N.S.: Isoperimetric inequalities for quermassintegrals. *Annales de l'I.H.P. Analyse non linéaire* **11**(4), 411–425, 1994
62. URAL'TSEVA, N.N.: Degenerate quasilinear elliptic systems. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **7**, 184–222, 1968
63. URBAS, J.I.E.: On the expansion of starshaped hypersurfaces by symmetric functions of their principal curvatures. *Math. Z.* **205**(3), 355–372, 1990
64. WEI, Y.: On the Minkowski-type inequality for outward minimizing hypersurfaces in Schwarzschild space. *Calc. Var. Partial Differ. Equ.* **57**(2), 46, 2018
65. WILLMORE, T.J.: Mean curvature of immersed surfaces. *An. Şti. Univ. "Al.I. Cuza" Iaşi Secţ. I a Mat. (N.S.)* **14**, 99–103, 1968
66. XIAO, J.: P-capacity vs surface-area. *Adv. Math.* **308**, 1318–1336, 2017
67. XU, X.: Some results on functional capacity and their applications to p-Laplacian problems involving measure data. *Nonlinear Anal. Theory Methods Appl.* **27**(1), 17–36, 1996

V. AGOSTINIANI · L. MAZZIERI

Università degli Studi di Trento,

Via Sommarive 14,

38123 Povo

TN

Italy.

e-mail: virginia.agostiniani@unitn.it

e-mail: lorenzo.mazzieri@unitn.it

and

M. FOGAGNOLO

Centro di Ricerca Matematica Ennio De Giorgi,

Scuola Normale Superiore,

Piazza dei Cavalieri 3,

56126 Pisa

PI

Italy.

e-mail: mattia.fogagnolo@sns.it

(Received January 15, 2021 / Accepted January 18, 2022)

© The Author(s) (2022)