



Element Formation in Radiation-hydrodynamics Simulations of Kilonovae

Fabio Magistrelli¹ , Sebastiano Bernuzzi¹ , Albino Perego^{2,3} , and David Radice^{4,5,6,7} ¹ Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, 07743 Jena, Germany; fabio.magistrelli@uni-jena.de, sebastiano.bernuzzi@uni-jena.de² Dipartimento di Fisica, Università di Trento, Via Sommarive 14, 38123 Trento, Italy; albino.perego@unitn.it³ INFN-TIFPA, Trento Institute for Fundamental Physics and Applications, via Sommarive 14, I-38123 Trento, Italy⁴ Institute for Gravitation & the Cosmos, The Pennsylvania State University, University Park, PA 16802, USA; david.radice@psu.edu⁵ Department of Physics, The Pennsylvania State University, University Park, PA 16802, USA⁶ Department of Astronomy & Astrophysics, The Pennsylvania State University, University Park, PA 16802, USA

Received 2024 May 17; revised 2024 August 27; accepted 2024 August 27; published 2024 October 1

Abstract

Understanding the details of r -process nucleosynthesis in binary neutron star merger (BNSM) ejecta is key to interpreting kilonova observations and identifying the role of BNSMs in the origin of heavy elements. We present a self-consistent, two-dimensional, ray-by-ray radiation-hydrodynamic evolution of BNSM ejecta with an online nuclear network (NN) up to a timescale of days. For the first time, an initial numerical relativity ejecta profile composed of the dynamical component and spiral-wave and disk winds is evolved including detailed r -process reactions and nuclear heating effects. A simple model for the jet energy deposition is also included. Our simulation highlights that the common approach of relating in postprocessing the final nucleosynthesis yields to the initial thermodynamic profile of the ejecta can lead to inaccurate predictions. Moreover, we find that neglecting the details of the radiation-hydrodynamic evolution of the ejecta in nuclear calculations can introduce deviations of up to 1 order of magnitude in the final abundances of several elements, including very light and second r -process peak elements. The presence of a jet affects element production only in the innermost part of the polar ejecta, and it does not alter the global nucleosynthesis results. Overall, our analysis shows that employing an online NN improves the reliability of nucleosynthesis and kilonova light-curve predictions.

Unified Astronomy Thesaurus concepts: [Hydrodynamics \(1963\)](#); [Nuclear astrophysics \(1129\)](#); [Ejecta \(453\)](#); [R-process \(1324\)](#); [Relativistic binary stars \(1386\)](#); [Novae \(1127\)](#)

1. Introduction

Mass ejecta from binary neutron star mergers (BNSMs) are primary sites for rapid neutron capture (r -process) nucleosynthesis (D. Eichler et al. 1989; see, e.g., J. J. Cowan et al. 2021; A. Perego et al. 2021; A. Arcones & F.-K. Thielemann 2023). The heavy, neutron-rich elements produced in these environments undergo radioactive decays, powering an electromagnetic (EM) transient known as a kilonova. Moreover, some of these mergers can produce short gamma-ray bursts (sGRBs), offering further insights into their astrophysical properties. The unambiguous detection of gravitational waves (GW170817) and EM counterparts (kilonova AT 2017gfo and GRB 170817A) from a BNSM in 2017 August has confirmed theoretical predictions and triggered intense work on the subject (B. P. Abbott et al. 2017a, 2017b, 2017c; D. A. Coulter et al. 2017; V. Savchenko et al. 2017; N. R. Tanvir et al. 2017).

Numerical simulations are the main tool to explore the physics of BNSMs and identify the mechanisms underlying the associated gravitational and EM emissions (S. Rosswog et al. 2014; L. F. Roberts et al. 2017; V. Nedora et al. 2021; D. Radice et al. 2022; D. Radice & S. Bernuzzi 2023; L. Combi & D. M. Siegel 2023a, K. Kiuchi et al. 2024; F. Schianchi et al. 2024; F. Zappa et al. 2023; 2023b; C. Musolino et al. 2024). Simulations are used to carry out extensive investigations on r -

process nucleosynthesis and its relationship with kilonovae. Lanthanide and actinide production is consistently found in low electron fraction ejected material ($Y_e \lesssim 0.2$; O. Korobkin et al. 2012; J. Lippuner & L. F. Roberts 2015; A. Perego et al. 2021). Low values of Y_e can be found in the equatorial component of dynamical ejecta (S. Wanajo et al. 2014; D. Radice et al. 2016; K. Kiuchi et al. 2023) and in the late disk wind (e.g., A. M. Beloborodov 2003; D. M. Siegel & B. D. Metzger 2017; K. Kiuchi et al. 2024; T. M. Sprouse et al. 2024). The high opacity associated with these elements commonly links the outer regions of the (low-latitude) ejecta with the red component of the kilonova light curves (B. D. Metzger & R. Fernandez 2014; A. Perego et al. 2017). However, small expansion timescales can inhibit neutron capture and thus produce a blue-UV precursor powered by the decay of free neutrons (B. D. Metzger et al. 2015; D. Radice et al. 2018a; L. Combi & D. M. Siegel 2023a). Higher values of $Y_e \gtrsim 0.3$ associated with most of the disk and the majority of polar ejecta prevent strong r -processes, thus leading to low-opacity matter contributing to the blue part of the kilonova spectra (B. D. Metzger & R. Fernandez 2014; D. Kasen et al. 2017; M. Tanaka et al. 2017; S. Curtis et al. 2024). In the polar regions, this is enhanced by interactions with neutrinos emitted by the central remnant until its possible collapse to a black hole (BH; D. Radice & S. Bernuzzi 2024). The passage of a relativistic jet can in principle affect the nucleosynthesis by injecting extra energy in the system and thus reigniting suppressed reactions (A. Janiuk 2014). Understanding the details of the nuclear evolution of the ejecta and how this affects its dynamics is crucial for accurately interpreting observational data and assessing the role of BNSMs in explaining the origin of r -process elements in the Universe.

⁷ Alfred P. Sloan fellow.

One of the main open challenges toward the accurate prediction of r -process nucleosynthesis in BNSM ejecta is capturing the dependence of the nuclear composition outcomes on the initial thermodynamic profile and detailed hydrodynamic evolution of the ejecta. Traditional approaches assume homologously expanding ejecta and neglect radiation transport, hydrodynamics, and ejecta self-interaction (e.g., O. Korobkin et al. 2012; D. Radice et al. 2016; S. Rosswog et al. 2017; A. Perego et al. 2022; S. Curtis et al. 2024), although some recent efforts have considered detailed radiation transport (e.g., C. E. Collins et al. 2023; L. J. Shingles et al. 2023). Alternatively, nuclear networks (NNs) are employed in a postprocessing step on hydrodynamics profiles to obtain yields and heating rates (D. Martin et al. 2015; O. Just et al. 2015; L. F. Roberts et al. 2017). The latter are sometimes fitted as time-domain functions and reused in hydrodynamical simulations to improve ejecta evolution (e.g., S. Rosswog et al. 2014; Z. Wu et al. 2022; G. Ricigliano et al. 2024).

In this work, we present, for the first time, a self-consistent simulation of r -process nucleosynthesis for a fiducial BNSM ejecta using a two-dimensional, ray-by-ray radiation-hydrodynamics code coupled with an online NN. In Section 2, we present our model and discuss how the NN is coupled with the radiation-hydrodynamic equations to update the ejecta composition and calculate the associated nuclear power in each hydrodynamic step. We also describe our implementation of thermalization processes and an extra energy source modeling a short gamma-ray burst (sGRB). Section 3 presents the nucleosynthesis and kilonova light-curve predictions for the ejecta profiles we extract from a fiducial BNSM numerical relativity simulation. In Section 4, we conclude by discussing the implications of our findings and emphasizing the importance of implementing an online NN for self-consistent predictions of r -process nucleosynthesis patterns.

2. Method

2.1. Numerical Relativity Profiles

Our simulation is initialized with initial ejecta profiles extracted from ab initio numerical relativity simulations that include microphysics, M0 neutrino transport,⁸ and magnetic-field-induced turbulence (D. Radice et al. 2018b; A. Perego et al. 2019; S. Bernuzzi 2020; V. Nedora et al. 2021). The considered binary has a mass ratio $q = 1.43$ and produces a short-lived remnant that collapses to a BH at about 10 ms postmerger. The simulation is run with the LS220 equation of state (EoS) and a typical resolution of 185 m up to 28.5 ms after merger (V. Nedora et al. 2021). Throughout the simulation, the unbound material is identified via the Bernoulli criterion and collected at an extraction radius of $R_{\text{ext}} \simeq 295$ km. The ejecta consists of the dynamical component of mass $\sim 1.24 \times 10^{-2} M_{\odot}$ and a short spiral-wave wind ($\sim 0.80 \times 10^{-2} M_{\odot}$). In order to prepare the subsequent long-term evolution, the dependence of the ejecta properties on the azimuthal coordinate is integrated out, while the polar dependence is accounted for by discretizing the polar angle in 51 angular sections. The time-dependent ejecta profile is mapped to a Lagrangian profile by positioning the latest shell at R_{ext} and progressively layering previously ejected shells on top as prescribed by the constraint $m(r) = 4\pi \int_{R_{\text{ext}}}^r \rho(r) r^2 dr$, where ρ is the mass density and r is

the position of the radius including the mass m (Z. Wu et al. 2022). An analytical disk wind profile of mass $\sim 2.75 \times 10^{-2} M_{\odot}$ is constructed from a similar simulation but performed on an equal-mass binary with the M1 neutrino transport (D. Radice et al. 2022), which better describes the optically thick regime (F. Zappa et al. 2023). The simulation of the $q = 1$ binary is run with the SFHo EoS for ~ 270 ms postmerger.⁹ The disk profile is constructed by interpolating the spherically averaged density, temperature, and electron fraction of the unbound matter over time and using second-order Padé approximants. We assume a constant velocity and entropy and rescale the density evolution to get a total ejection of 40% of the total disk mass (of the LS220 simulation) assuming a $\propto \sin^2 \theta$ angular distribution (A. Perego et al. 2017; R. Fernandez et al. 2019). The complete initial profiles of two selected angular sections are shown in the two bottom panels of Figure 1.

2.2. Ray-by-ray Radiation Hydrodynamics

The ejecta profile is evolved with the system of Lagrangian radiation-hydrodynamic equations as implemented in V. Morozova et al. (2015) and Z. Wu et al. (2022). In particular, the energy equation in spherical symmetry reads

$$\frac{\partial \epsilon}{\partial t} = \frac{P}{\rho} \frac{\partial \ln \rho}{\partial t} - 4\pi r^2 Q \frac{\partial v}{\partial m} - \frac{\partial L}{\partial m} + \dot{\epsilon}_{\text{nucl}}, \quad (1)$$

where m is the mass coordinate, ϵ is the specific internal energy, t is time, P is pressure, v (radial) is velocity, L is luminosity, and Q is the von Neumann–Richtmyer artificial viscosity (J. Von Neumann & R. D. Richtmyer 1950). The specific energy deposition $\dot{\epsilon}_{\text{nucl}}$ accounts for the local production and partial thermalization of the energy released by all possible nuclear reactions occurring in the expanding material. To calculate the luminosity, we employ the same analytic, time-independent opacity introduced in Z. Wu et al. (2022).

The set of hydrodynamic equations is closed by the EoS. In the high-temperature regime, we implement the tabulated Helmholtz EoS introduced in F. X. Timmes & F. D. Swesty (2000) and also used in J. Lippuner & L. F. Roberts (2017). For lower temperatures, where the contribution of positrons is negligible, we switch to the Paczynski EoS introduced in B. Paczynski (1986) and utilized in V. Morozova et al. (2015) and Z. Wu et al. (2022). This analytical EoS describes a mixture of ideal gases, namely, a Boltzmann gas of nondegenerate, nonrelativistic ions; an ideal gas of arbitrarily degenerate and relativistic electrons; and a photon ideal gas. The inclusion of an online NN, as described in Section 2.3, improves the realism of these EoSs, as the mean molecular weight and the electron fraction, which enter the EoS, are self-consistently calculated on the fly.

To account for the angular dependency of the ejecta properties, we employ the spherically symmetric hydrodynamic equations in a ray-by-ray fashion. For each angular section, we first map the profile into an effective one-dimensional problem

⁸ See F. Zappa et al. (2023) for the importance of including neutrino heating effects.

⁹ We take the results of this simulation, where the BH collapse occurs around $t \simeq 10$ ms postmerger, as a representative history of disk ejecta. The general properties of disks and disk ejecta are found to be relatively stable against variations in the properties of the compact binary or numerical prescriptions (e.g., EoS, effective viscosity, neutrino treatment), as long as the BH collapse occurs on a timescale of $\mathcal{O}(10)$ ms (R. Fernández & B. D. Metzger 2013; O. Just et al. 2021, 2023; K. Kiuchi et al. 2023, 2024; A. Camilletti et al. 2024).




Figure 1. Evolution of the ejecta composition for every mass shell of the angular sections $\theta \simeq 12^\circ, 90^\circ$ (first and second column, respectively). The x-axis indicates the Lagrangian mass coordinate, linear in the shell index. The vertical black dotted lines indicate the point where the analytical disk wind is attached to the dynamical ejecta profile. The color maps in panels (a)–(c) and (h)–(l) represent the mass fractions of neutrons (gray); protons (red); first (blue), second (orange), and third (purple) r -process peak elements; and rare earths (brown). Panels (d), (m) and (e), (n) show the evolution of the parameter $h(t)$ defined in Equation (4) (blue to red) and of the electron fraction (green to pink), respectively. In (e) and (n), we also plot in blue the amount of mass δm included in each shell, rescaled by the total mass of the angular section $M_{\text{ej}}(\theta)$, and in black the position of the photosphere. Subplots (f), (g) and (o), (p) show the initial values of $Y_{e,0}$, s , T , and τ as a function of the included mass. The horizontal dotted lines indicate the limit values of $Y_{e,0} = 0.22$, $s_0 = 100 \text{ k}_B \text{ baryon}^{-1}$, $T_0 = 7 \text{ GK}$, and $\tau_b = 5 \text{ ms}$.

by multiplying the total mass by the scaling factor $\lambda_\theta = 4\pi/\Delta\Omega$, where $\Delta\Omega \simeq 2\pi \sin \theta d\theta$ represents the solid angle included in the angular section. This ensures that all the intensive quantities (including density) remain fixed. Then, we independently evolve the different angular sections by discretizing the one-dimensional hydrodynamic equations over n_{sh} spherical fluid elements (mass shells). We ran tests for some selected sections with grid resolutions $n_{\text{sh}} = 300, 600$, and 1000 . We found $n_{\text{sh}} = 600$ sufficiently high to resolve the details of the nucleosynthesis and decided to use this resolution for the complete simulation. Note that nonradial flows of matter and radiation are neglected, as well as higher-dimensional fluid instabilities (e.g., Kelvin–Helmholtz), convection, and the possibility of shells surpassing each other. This could in principle lead to over- or underestimating the pressure work exchanged between the radial shells and introduce artificial

shocks in the system. Finally, we map the problem back to the axisymmetric scenario by keeping the intensive quantities unchanged, rescaling the extensive ones by the scaling factor $1/\lambda_\theta$, and combining the results. In particular, we calculate the global mass fractions and abundances with a mass-weighted average over all the mass shells and angular sections. Kilonova light curves are recombined, accounting for the angle of view as in D. Martin et al. (2015) and A. Perego et al. (2017).

2.3. NN Coupling

We calculate the nuclear composition and specific energy deposition, ϵ_{nuc} , in a self-consistent way with an online implementation of the NN *SkyNet* (J. Lippuner & L. F. Roberts 2017). The NN includes 7836 isotopes up to ^{337}Cn and uses the JINA REACLIB (R. H. Cyburt et al. 2010) and the same setup as in J. Lippuner & L. F. Roberts (2015) and

A. Perego et al. (2022). The simulation is started by initializing matter composition from the initial temperature $T_0(m)$ and density $\rho_0(m)$ assuming nuclear statistical equilibrium (NSE). We thus impose, for each isotope considered in the NN, the abundance (J. J. Cowan et al. 2021; A. Perego et al. 2021)

$$Y_{i,0} = \frac{n_{i,0}}{n_{b,0}} = Y_{p,0}^{Z_i} Y_{n,0}^{(A_i-Z_i)} \frac{G_i(T) A_i^{3/2}}{2^{A_i}} \left(\frac{\rho}{m_b} \right)^{A_i-1} \times \left(\frac{2\pi\hbar^2}{m_b k_B T} \right)^{3(A_i-1)/2} e^{BE_i/k_B T}, \quad (2)$$

where m_b and n_b are the baryon mass and number density, A_i and Z_i are the mass and atomic numbers of the i th isotope, and n_i and BE_i are its number density and total binding energy.¹⁰ Here, $G_i(T)$ is the internal partition function of the nucleus i at temperature T . The two conditions $\sum_i A_i Y_i = 1$ and $\sum_i Z_i Y_i = Y_e$, where Y_e is the electron fraction, constrain the NSE composition, which thus depends on ρ , T , and Y_e . See J. Lippuner & L. F. Roberts (2017) for details about SkyNet’s implementation of Equation (2). The isotopic mass fractions can be calculated from the element abundances via $X_i = A_i Y_i$.

At the initial time t_0 of each hydrodynamic step Δt , an independent instance of SkyNet is called in each mass shell. The NN evolves the composition with its own time substeps until $t_0 + \Delta t$. The (nonthermalized) power released by nuclear reactions is averaged over Δt to get the total nuclear power that will then be thermalized into $\dot{\epsilon}_{\text{nucl}}$ as described below. During the substeps, the temperature is constant (no self-heating) and fixed by $T(t_0)$, and it will only be updated by the hydrodynamic step. The density evolution is prescribed with a log-interpolation between $\rho(t_0)$ and $\rho(t_0 + \Delta t)$. The latter is calculated from the density and radial velocity profiles at $t = t_0$. Note that in our setup, mass shells cannot mix.

To calculate the specific energy deposition in Equation (1), we independently thermalize the different contributions to $\dot{\epsilon}_{\text{nucl}} = \dot{\epsilon}_{\text{nucl}}^\gamma + \dot{\epsilon}_{\text{nucl}}^\alpha + \dot{\epsilon}_{\text{nucl}}^\beta + \dot{\epsilon}_{\text{nucl}}^{\text{oth}}$ coming from γ -rays, α particles, electrons, and other nuclear reaction products. We compute the fraction of energy thermalized by the emission of γ -rays as

$$\dot{\epsilon}_{\text{nucl}}^\gamma(t) = \sum_j f_j^\gamma(t) \frac{\langle E_j^\gamma \rangle Y_j(t)}{\tau_j m_p}, \quad (3)$$

where j is the isotope index and m_p is the proton mass. The average lifetimes τ_j and the mean energy $\langle E_j^\gamma \rangle$ released by each of these nuclei in γ -rays via radioactive decay are taken from the ENDF/B-VIII.0 database¹¹ (D. A. Brown et al. 2018). The thermalization factor $f_j^\gamma(t)$ is calculated, as in K. Hotokezaka & E. Nakar (2020) and L. Combi & D. M. Siegel (2023a), starting from the detailed composition of the ejecta and the same effective opacity tables from the NIST-XCOM catalog¹²

¹⁰ Several mass shells have $T < 5$ GK at the end of the numerical relativity simulation (see Figure 1). In the case of low-entropy tidal ejecta, their initial composition should be close to the one set by cold NSE conditions inside isolated neutron stars. In all the other cases, we expect matter to reach hot NSE conditions inside our extraction radius, before matter expansion causes an NSE freeze-out. We plan to attest to the impact of the early out-of-NSE evolution on our results by developing and comparing alternative initialization procedures for cold fluid elements in a future work.

¹¹ <https://www-nds.iaea.org/exfor/endl>

¹² <https://www.nist.gov/pml/xcom-photon-cross-sections-database>

(M. Berger et al. 2010) used in J. Barnes et al. (2016). For the energy injected in electrons and α particles and their thermalization factors, we use the analytic expressions of D. Kasen & J. Barnes (2019). Neutrinos from β -decays do not deposit energy into the system, while fission fragments and daughter nuclei tend to thermalize very efficiently (J. Barnes et al. 2016). We approximate this behavior by assuming that half of the remaining (nonthermalized) heating rate from SkyNet is deposited into the ejecta.

2.4. Jet Energy Deposition

We consider a jet model following the “thermal bomb” prescription of V. Morozova et al. (2015). Essentially, an extra energy term is added to the right-hand side of Equation (1) for the innermost shells of the ejecta during a chosen time interval. The thermal bomb parameters are fixed assuming that the released isotropic energy from the sGRB is proportional to the kinetic energy of the structured jet described by G. Ghirlanda et al. (2019), $E_{\text{iso}}(\theta) = E_0/[1 + (\theta/\theta_j)^{5.5}]$, where θ_j is the jet’s opening angle. The total energy E_j released by the jet is related to E_{iso} by $E_{\text{iso}} = (4\pi/\Delta\Omega_j)E_j$, where $\Delta\Omega_j = 2\pi(1 - \cos\theta_j)$ is the solid angle covered by the jet. We fix $\theta_j = 15^\circ$, within the range also explored in H. Hamidani et al. (2020) for the sGRB associated with GW170817. We simulate a jet with the same duration as the observed sGRB from B. P. Abbott et al. (2017c), $\Delta t_j = 100$ ms. We launch the jet at $t = 200$ ms and fix the parameter $E_0 = 10^{51}$ erg to be compatible with the isotropic luminosity range given by H. Hamidani et al. (2020). Their Figure 9 shows that, in the case of GRB 170817A, choosing the isotropic luminosity to be $L_{\text{iso}} \sim 10^{52}$ erg s^{−1} constrains the jet to be launched no more than 300 ms after merger.

3. Results

3.1. Fluid Element Composition Evolution

Our simulation allows us to self-consistently monitor the nuclear composition of the matter during its expansion. Figure 1 shows the time evolution of the mass fraction of protons, neutrons, rare earths, and r -process peak elements (panels (a)–(c) and (h)–(l)), the parameter $h(t)$ (see below; panels (d) and (m)), and the electron fraction (panels (e) and (n)), together with the initial thermodynamic conditions of the ejecta (panels (f), (g) and (o), (p)), as a function of the Lagrangian mass coordinate. The left and right panels refer to two different angular sections, $\theta = 12^\circ, 90^\circ$. Mass shells with, respectively, $m \lesssim 50 \times 10^{-6} M_\odot$ and $m \lesssim 1.4 \times 10^{-3} M_\odot$ correspond to the wind, while larger values correspond to the dynamical ejecta. In the latter, the entropy slightly increases toward the outer shells, where a high-entropy, low-mass tail is observed at all angles. Material with low initial electron fractions, $Y_{e,0} < 0.22$, is found in the outer regions of the ejecta and, for $m \sim 2.5 \times 10^{-3} M_\odot$, around the equatorial plane. During the evolution, the electron fraction rises to $Y_e > 0.22$ for almost every mass shell at around $t \sim 1$ s due to the β -decays following the neutron captures and neutron freeze-out.

The color maps in panels (d) and (m) of Figure 1 show the evolution of the parameter:

$$h(t) \equiv \log_{10} \left(\frac{\rho(t)}{\rho_{\text{SkyNet}}(t | \rho_0, \tau_b)} \right). \quad (4)$$

The density evolution for the analytical expansion ρ_{skyNet} is parameterized by the initial density ρ_0 and expansion timescale τ_b as defined in Equation (1) of J. Lippuner & L. F. Roberts (2015). Here, τ_b is calculated from the expansion timescale measured at the extraction radius as prescribed in D. Radice et al. (2016). The same homologous expansion is used by A. Perego et al. (2022) and Z. Wu et al. (2022) for the time-domain fitting of the heating rates. Positive (negative) values of h indicate regions of the t - M plane where the mass shells are characterized by higher (lower) densities than the one predicted by the simple analytical model. Shells where h is constant in time are expanding homologously. This happens for most of the shells in our simulation for $t \gtrsim 100$ s, in agreement with the results of S. Rosswog et al. (2014). If $h \neq 0$, the evolution happens on a curve, $\rho_{\text{skyNet}}(t | \rho'_0, \tau'_b)$, different from the one associated with the values of ρ_0, τ_b of the considered mass shell. We identify a deviation of approximately 2–3 orders of magnitude between our numerical solution and the specific analytical expansion $\rho_{\text{skyNet}}(t | \rho_0, \tau_b)$ in all the significant timescales. As further explained below, such a different expansion in the early evolution ($t \lesssim 1$ s, when the production of heavy nuclei is still ongoing) can lead to inconsistent nucleosynthesis predictions. In Appendix A, we discuss an example of how a full evolution of a fluid element can differ from the one obtained by an independent NN assuming $\rho_{\text{skyNet}}(t | \rho_0, \tau_b)$. A comparison with the results from a run using the fitted heating rates of Z. Wu et al. (2022) instead of the data from the coupled NN reveals that the disagreement between the predicted density evolution and the simulated one builds up mostly because of hydrodynamic effects. The relative differences in the nuclear heating are not big enough to impact the hydrodynamic evolution while competing with the pressure work in Equation (1) (see Appendix B).

At $t \sim 100$ ms, a sudden increase of h is observed in the disk ejecta. The analysis of the released nuclear energy reveals a correspondent absorption of energy from the fluid to the NN ($\dot{\epsilon}_{\text{nucl}} < 0$). In the same region, the abrupt change in the h parameter reveals ongoing shocks. A series of shock-induced photodissociation processes could explain the negative heating rates, in analogy with supernova simulations (see, e.g., H.-T. Janka 2012; A. Burrows 2013). Such an effect is genuinely due to the coupling between the NN and hydrodynamics, and it cannot be reproduced by offline analyses. In the polar regions, a further compression at $t \sim 200$ ms, followed by a more rapid expansion, is caused by the extra energy deposited by the jet.

Panels (a)–(c) and (h)–(l) of Figure 1 show the evolution of the cumulative mass fractions of groups of selected isotopes. Consistently throughout the ejecta, shells with $Y_{e,0} \lesssim 0.22$ produce first-peak elements (embedded in a sea of free neutrons) within $t \lesssim 10^{-4}$ s. On a timescale of 1 s, these nuclei are converted to second- and third-peak elements and a small fraction of rare earths. At $T_0 \gtrsim 7$ GK, most of the ejecta is initially composed of free neutrons and protons (their ratio depending on the initial electron fraction) and light elements, which will act as seeds for r -process nucleosynthesis. In regions where $T_0 \lesssim 7$ GK, the initial temperature is low enough in certain mass shells (see the arrows in panel (i)) to make the binding energy term in Equation (2) strongly affect the NSE composition. Hence, a significant fraction of first-peak elements is already formed at $t = 0$ and can be directly used

as r -process seeds. This can ease the heavy-element production and, in some cases, partially induce strong r -processes even for $Y_e \gtrsim 0.22$ (see, for example, the high- $Y_{e,0}$, low- s_0 third-peak production in the $\theta = 90^\circ$ case).

High enough values of the initial entropy ($s_0 \gtrsim 100 k_B$ baryon $^{-1}$) can lead to the production of rare earths, lanthanides, and actinides even for intermediate values of the initial electron fraction, $0.22 \lesssim Y_e \lesssim 0.35$. This outcome is compatible with an α -rich freeze-out, characterized by a large neutron-to-seed ratio (O. Just et al. 2015; J. J. Cowan et al. 2021). However, not all shells with high initial entropy are sites of strong r -process nucleosynthesis. This is clearly shown, for example, by the $\theta = 12^\circ$ angular section in the region $74.07 \lesssim m/(10^{-6} M_\odot) \lesssim 74.60$; see panels (a) and (b). When strong r -processes are activated, free neutrons are efficiently consumed, reducing the final abundance of free protons. Starting from $m \sim 74.07 \times 10^{-6} M_\odot$ and moving outward in the ejecta, while the initial T_0, s_0 , and τ_b profiles are almost constant (panels (f) and (g)), strong r -processes get progressively suppressed. The suppression is smooth even given the jump in the initial conditions around $m \sim 74.60 \times 10^{-6} M_\odot$, which also does not change the second-peak production in panel (c). This discussion gives an example of how only the detailed evolution of the local thermo- and hydrodynamics conditions determine with accuracy the subsequent nucleosynthesis yields.

For low entropies, $s_0 \lesssim 100 k_B$ baryon $^{-1}$, and intermediate values of the initial electron fractions, $0.22 \lesssim Y_{e,0} \lesssim 0.35$, the fewer free neutrons are rapidly captured by seed nuclei, and only second-peak elements are produced at the end of the nucleosynthesis. Note again the exception represented by the shells at around $m \gtrsim 2.704 \times 10^{-3} M_\odot$ in the equatorial section. At the high $Y_{e,0} \gtrsim 0.35$ reached in the polar section, weak r -processes occur at low entropy, and only first-peak elements can be produced.

Extremely high values of the initial entropy, $s_0 \gtrsim 200 k_B$ baryon $^{-1}$, can be reached in the outermost shells of the ejecta. Despite the large neutron-to-seed ratio, neutron capture is not effective in the fast ejecta tail ($v \gtrsim 0.6$). Nucleosynthesis is therefore hindered. In this rarefied environment, most of the neutrons remain free after the r -process freeze-out and start β -decaying on a timescale of $t \sim 10$ minutes while crossing the photosphere, whose position is tracked in panels (e) and (n). This could power a UV/blue kilonova precursor on a timescale of hours, as discussed in Section 3.3. After 10^4 s, no heavy element is produced, and free protons dominate the final matter composition. This is reflected in the high final electron fractions of these shells. If the strong rise in entropy is combined with an initial electron fraction $Y_{e,0} \gtrsim 0.35$, all neutrons are initially bound in stable first-peak elements. In these outer regions, the composition remains essentially frozen. We can thus predict a negligible nuclear contribution to the kilonova light curves from these shells throughout the ejecta evolution, regardless of the photosphere position.

The qualitative changes in the nucleosynthesis patterns caused by the jet are negligible. The only quantitative effect observed on the mass fractions depicted in Figure 1 is a very slight delay in the production of second r -process peak elements in the innermost shells of the polar sections. No significant effects arise in the regions near the equatorial plane, as the jet energy is negligible at low latitudes, and we do not account for the coupling between different angular sections.




Figure 2. Global r -process nucleosynthesis yields at $t \simeq 5 \times 10^4$ s. Top panel: in blue and black, the final global abundances $\langle Y_{A,f} \rangle$ as a function of the mass number A , obtained, respectively, from our SNEC simulation and with a postprocess analysis similar to A. Perego et al. (2022). In orange and green are the results obtained from our SNEC simulation excluding the disk ejecta or including only the matter emitted at $\theta = 12^\circ$ (at all times). Red dots represent the solar r -abundances (N. Prantzos et al. 2020) scaled to match the average value of the second peak. The histogram shows, for the complete simulation, the global cumulative mass fractions $\langle X(A) \rangle$ of selected groups of elements, in particular, first (blue), second (orange), and third (purple) r -process peaks and rare earths (brown). We report the initial value of the average electron fraction of the ejecta in the top left corner of the plot. Bottom panel: in red, the relative difference against the model with prescribed homologous expansion (see Equation (5)), and in black and purple, analogous comparisons between our global results and the postprocess analysis and between the results of the complete simulation with and without the jet. The differences are only shown for mass numbers $\hat{A}$ such that $\langle Y_{\hat{A},f} \rangle > 10^{-10}$.

In summary, our results indicate that it is not possible to predict the final nucleosynthesis yields based solely on the initial thermodynamic conditions of the ejecta.

3.2. Global Nucleosynthesis Yields

We calculate the global abundances $\langle Y_i \rangle$ as mass-weighted averages over all the mass shells of all the angular sections. In Figure 2, we show the matter composition at $t \simeq 5 \times 10^4$ s. Except for small modifications due to long-lived α -decaying isotopes with $A \gtrsim 220$, this is a good representation of the final r -process nucleosynthesis yields. All the r -process peaks and (in part) rare earths are produced. This is expected, as matter is mostly ejected along the equatorial plane, where the neutron-rich dynamical ejecta keeps the global initial average electron fraction at $\langle Y_{e,0} \rangle \simeq 0.24$. The production of heavy elements is partially hindered in the polar sections, consistent with their higher Y_e and the picture outlined in Figure 1. By comparing the results of our complete model with the ones obtained by excluding the late disk wind, we establish that the relative production of heavy elements is already saturated in the dynamical ejecta. Most of the lighter elements with $A \lesssim 130$ (in particular elements between the first and second peaks and isotopes of the iron group) are more strongly produced by the disk wind. These results are in agreement with D. Martin et al. (2015), J. J. Cowan et al. (2021), S. Curtis et al. (2023), and L. Chiesa et al. (2024).

In the bottom panel of Figure 2, we show the final value of the relative difference,

$$\Delta \langle Y_A^{\text{ch}} \rangle = \frac{|\langle Y_A \rangle - \langle Y_A^{\text{ch}} \rangle|}{\min(\langle Y_A \rangle, \langle Y_A^{\text{ch}} \rangle)}, \quad (5)$$

between our results and the predictions $\langle Y_A^{\text{ch}} \rangle$ obtained with a SNEC simulation of the ejecta with coupled NN and a prescribed homologous expansion. We evolve the system with the full hydrodynamic equations up to $t_0 = 10$ ms and then force the density to evolve as $\rho(t, m) = \rho(t_0, m)(t/t_0)^{-3}$. Neglecting the details of the hydrodynamic evolution introduces an error of $\gtrsim 10\%$ for almost all values of mass numbers. The biggest discrepancies are observed for some light and second-peak elements. We also compare our results against a postprocess analysis similar to the method described in A. Perego et al. (2022) but adapted to our initialization method. The agreement is overall worse in the new case, and the position of the third r -process peak is shifted. The extra source of error is introduced here by effectively coarsening the resolution on the initial thermodynamic conditions (see Appendix C for details). Figure 3 shows the evolution of the abundances for free protons and neutrons and a few selected isotopes. The r -process nucleosynthesis takes place between $t \sim 10^{-2}$ s and $t \sim 1$ s, when the first drop in $\langle Y_n \rangle$ occurs and the abundances of lanthanides and actinides saturate at $\langle Y_i \rangle \simeq 10^{-4}$. The plateau at $Y_n \simeq 10^{-4}$ in panel (a) indicates the presence of regions of the ejecta where the material undergoes incomplete neutron burning. The second drop in the abundance of neutrons at $t \sim 10$ minutes, connected with an increase in the proton fraction, is due to free neutron β -decays. Our analysis shows qualitative features similar to those discussed in A. Perego et al. (2022). However, a direct comparison between the two models shows discrepancies of several orders of magnitude in the abundance evolution of various isotopes; see panel (c). In particular, note the 6 orders of magnitude difference in the neutron abundance for the postnucleosynthesis plateau and the overproduction of ^{56}Fe from the postprocessing of about 2 and 1 orders of magnitude at timescales between 10^{-2} – 10^3 s and $t \sim 10^4$ s, respectively. At late times, the postprocessing has underproduced ^4He , ^2H , and ^3H and free protons of about 1, 3, and 6 orders of magnitude.

The initial global abundance of ^4He is of the order of 10^{-2} because of shells undergoing α -rich freeze-out; see panel (a). This isotope is partially consumed at early times to construct heavier elements, e.g., ^{88}Sr . In the innermost shells, it is produced again during the r -process nucleosynthesis by α -decays of the freshly produced, heavy, neutron-rich elements; see panel (b). These regions are crossed by the jet, which produces a rapid increase in the abundances of protons and deuterium at $t \simeq 200$ ms. ^{88}Sr is initially strongly suppressed, but it rises enough at early times to effectively act as a seed for r -process nucleosynthesis; see panel (a). At late times, this element is produced again as part of the first peak by β -decaying neutron-rich isotopes. This is particularly evident in panel (b). We also show the abundance of ^{56}Fe as representative of the most bounded nuclei around the iron peak. Its initial formation is favored by its high binding energy.




Figure 3. Evolution of the abundances of a few selected isotopes and the cumulative abundances of lanthanides and actinides. Panel (a) shows global results obtained with mass-weighted averages all over the ejecta. Panel (b) is restricted to an average over the innermost shell of all the angular sections in order to highlight the jet effects. Panel (c) shows the ratio $\Delta \langle Y_i \rangle = \langle Y_i \rangle / \langle Y_i^{pp} \rangle$ between our results and the prediction obtained with a postprocess procedure similar to the one of A. Perego et al. (2022); see Appendix C for details. Panel (d) shows a similar comparison between the results obtained with and without the jet. The vertical dashed lines in panels (b) and (d) indicate the time when the jet is launched. In panels (c) and (d), the results are only shown for times t such that $\langle Y_i \rangle(t) > 10^{-12}$.




Figure 4. AB apparent magnitudes predicted by our simulation for the Gemini bands *u*, *g*, *i*, and *K_s* for an observation angle of $\theta = 20^\circ$ and a distance of 40 Mpc. Left panel: results of the complete simulation (solid lines) against the predictions obtained by excluding the jet (dashed lines) or the disk component (dashed-dotted lines). Central panel: isotropized $\theta = 12^\circ, 90^\circ$ angular sections (solid and dashed lines, respectively). Right panel: light curves obtained assuming a constant thermalization factor $f_{th} = 0.5$ with (solid lines) or without (with the same heating rate fits of Z. Wu et al. 2022; dashed lines) a coupled NN.

Like strontium, ^{56}Fe acts as a seed nucleus, but it is only partially produced again by later nuclear reactions.

The additional energy released by the jet significantly affects only a fraction of the inner shells at the highest latitudes. The mass involved is too small a fraction of the ejecta to yield visible effects on the global results presented in Figures 2 and 3. In particular, the final nuclear yields are compatible to a few percent between the complete simulations with and without the jet (see the bottom panel of Figure 2). Nevertheless, the examination of the nucleosynthesis within the inner shells reveals some modifications of the final abundances for the selected elements of Figure 3; see panel (d). The sudden acceleration induced by the jet slows neutron captures down, therefore leaving, at the end of the nucleosynthesis, fewer lanthanides, more free β -decaying neutrons, and, consequently, free protons. The faster dynamics also inhibits the burning of

light elements, leading to an increase in the abundance of hydrogen isotopes at the end of the simulation.

3.3. Light Curves

In Figure 4, we show the kilonova light curves predicted by our model for a few selected UV/visible/IR bands for an observation angle of $\theta = 20^\circ$ and a distance of 40 Mpc.

The disk is predicted to contribute to the blue component of the kilonova due to its lower opacity (see, e.g., B. D. Metzger & R. Fernandez 2014; D. M. Siegel 2019; S. Curtis et al. 2024). However, at all angles, the photosphere enters the disk region only at late times ($t \sim 10$ days), when the temperature drop has already made the disk spectra red. Essentially, the dynamical ejecta acts as a curtain, preventing the disk from significantly boosting the blue part of the kilonova. Correspondingly, in the left panel of Figure 4, all the curves show an analogous

decrease in their luminosity when the disk is excluded, only due to the removal of part of the ejected mass. The central panel of Figure 4 shows that, around the equator, the opaque lanthanide curtain associated with the dynamical ejecta produces a red kilonova. The blue components of the spectra become dominant at higher latitudes, suggesting a main contribution to the blue kilonova from the angular sections around the disk edge, where a still significant mass of relatively neutron-poor material is ejected.

To assess the impact of the online NN on the kilonova predictions, we plot in the right panel of Figure 4 the results obtained without evolving the matter composition but taking as input the heating rate fits employed already in Z. Wu et al. (2022).¹³ Around $t \sim 4 \times 10^{-2}$ days, the simulation with a coupled NN shows a bump in the blue light curves, compatible with the UV precursor predicted in B. D. Metzger et al. (2015) and L. Combi & D. M. Siegel (2023a) to be powered by β -decaying free neutrons. This feature is not reproduced by the simulation with no coupled NN. The reason is that the fits of Z. Wu et al. (2022) do not capture the free neutron contribution to the heating rate (see their Figure 2 at $t \sim 10^{-2}$ days). Most importantly, the simulation without a coupled NN fails in predicting the position and brightness of the main peak of the high-frequency filters. It predicts instead a faster dimming at late times of the low-frequency ones. Thus, even if a coupled NN does have a negligible impact on the hydrodynamic evolution, it introduces significant corrections in the predicted light curves. While the heating rate competes with the pdV work and thus has a negligible effect on the hydrodynamics (see also Appendix B), it acts more directly on the luminosity of the optically thin regions above the photosphere. Moreover, relatively small changes in the temperature of the photosphere can lead to an observable shift in the frequency peak of its blackbody radiation.

The jet slightly increases the early ($t \sim 0.1$ –1 days) light curves in all bands. The effect is stronger and can be seen earlier for higher frequencies (see the left panel of Figure 4). This is due to the extra energy input increasing the temperature and thus leading to a bluer and brighter emission. A similar effect is discussed in L. Nativi et al. (2020), where the jet is found to clear a significant fraction of the lanthanide curtain.

4. Conclusions

In this work, we investigated the nucleosynthesis process in a BNSM ejecta by means of a two-dimensional, ray-by-ray, radiation-hydrodynamics simulation incorporating an online NN.

Our results challenge the widely used approach of predicting r -process nucleosynthesis from isolated fluid elements. Figure 1 illustrates that it is not possible to fully predict the detailed distribution of the (final) nucleosynthesis yields solely from the initial thermodynamic conditions of a set of fluid elements. Moreover, the most commonly employed approximations for homologous expansions fail to capture the asymptotic hydrodynamic evolution of the unbound material. Therefore, even estimates for the nuclear heating rate obtained by postprocessing isolated fluid elements (see, e.g., S. Rosswog

et al. 2014; S. Wanajo 2018; Z. Wu et al. 2022; C. E. Collins et al. 2023; L. J. Shingles et al. 2023; G. Ricigliano et al. 2024; S. Rosswog & O. Korobkin 2024) may lead to inconsistent results. To get a consistent picture of the ongoing nuclear processes, the effects related to radiation transfer and ejecta self-interaction must be taken into account. On the other hand, the corrections introduced by an online NN on the nuclear energy supplied to or removed from the expanding material do not introduce significant changes in the hydrodynamic evolution of the ejecta. However, nuclear reactions have a more direct impact on their EM emission (see Figure 4). Providing a radiation-hydrodynamic simulation of the ejecta from compact binary mergers with an online NN thus also appears crucial for improving the realism of the predicted kilonova light curves.

Some of the outer layers of the ejecta primarily consist of β -decaying free neutrons (see Figure 1), in agreement with J. de Jesús Mendoza-Temis et al. (2015) and D. Radice et al. (2018a). In our simulations, having a very rapidly ($\tau \lesssim 5$ ms) expanding ejecta is not sufficient to prevent the production of third-peak elements. Very high entropies, $s \gtrsim 100 k_B \text{ baryon}^{-1}$, are also required to leave a strong abundance of free neutrons after the r -process freeze-out.

From a qualitative point of view, our averaged results for nucleosynthesis yields in Figure 2 align with other predictions in the literature (e.g., A. Perego et al. 2021; J. J. Cowan et al. 2021). Our analysis reveals a connection between the production of light (below second peak) and third-peak elements with late disk and dynamical ejecta, respectively, in agreement with D. Martin et al. (2015) and J. J. Cowan et al. (2021). Figure 4 shows that the outer lanthanide curtain shields the radiation produced by the inner disk ejecta, thus preventing them from giving the significant contribution to the blue component of the kilonova predicted by B. D. Metzger & R. Fernandez (2014), D. M. Siegel (2019), and S. Curtis et al. (2024). The polar part of our ejecta evolves from higher Y_e and produces few heavy r -process elements; this is consistent with the link discussed in D. Kasen et al. (2017), M. Tanaka et al. (2017), M. Nicholl et al. (2017), and A. Perego et al. (2017) between the blue kilonova component and these sections of the unbound material.

We compared the final results for the nucleosynthesis yields with a SNEC run where a homologous expansion is prescribed from $t = 10$ ms. Neglecting the details of the radiation-hydrodynamics evolution of the ejecta introduces deviations $\gtrsim 10\%$ for all the mass numbers (see the bottom panel of Figure 2). In particular, the final abundances of some light ($A \lesssim 50$) and second r -process peak elements differ by almost 1 order of magnitude. We also compared the time evolution of the nuclear abundances with a postprocessing method analogous to a previous work that utilized the same NN (A. Perego et al. 2022). We found significant quantitative differences for some of the analyzed elements (a qualitative agreement is found only for some specific shells in our simulation). In particular, we found discrepancies of several orders of magnitude in the evolution of ^4He , ^{56}Fe , deuterium, tritium, and free protons; see Figure 3. The differences are due both to the coupling of the NN with the radiation-hydrodynamic evolution and to the grid introduced by the postprocessing method on the initial thermodynamic conditions.

The inclusion of an extra energy term mimicking a jet only affects the dynamic evolution of the polar regions of the ejecta,

¹³ Without the composition information given by the NN, we cannot calculate the thermalization factors as described in Section 2.3. We therefore follow Z. Wu et al. (2022) by imposing a constant thermalization factor $f_{\text{th}} = 0.5$ and compare the results against a complete simulation with a coupled NN but with the same simple thermalization.

without altering the global qualitative predictions of the r -process nucleosynthesis.

Our work will be extended and improved in several directions. We aim at implementing an improved thermalization of charged particles and a more realistic opacity treatment based on the complete, tracked information about matter composition. A following paper will report a systematic investigation of kilonova light curves using numerical relativity profiles that are hundreds of milliseconds long and the improved thermalization and opacity models. Finally, the methods developed here will be ported to three-dimensional simulations.

Acknowledgments

F.M. acknowledges support from the Deutsche Forschungsgemeinschaft (DFG) under grant No. 406116891 within the Research Training Group RTG 2522/1. F.M. also acknowledges ECT* for organizing the MICRA2023 workshop and TIFPA for their invitation to Trento in 2023 December. Both occasions were sites of useful discussions. S.B. acknowledges funding from the EU Horizon under ERC Consolidator grant No. InspiReM-101043372 and from the Deutsche Forschungsgemeinschaft (DFG) project MEMI No. BE 6301/2-1. The work of A.P. is partially funded by the European Union under NextGenerationEU, PRIN 2022 Prot. No. 2022KX2Z3B. D.R. acknowledges funding from the U.S. Department of Energy, Office of Science, Division of Nuclear Physics under award Nos. DE-SC0021177 and DE-SC0024388 and from the National Science Foundation under grant Nos. PHY-2011725, PHY-2020275, PHY-2116686, and AST-2108467.

Simulations were performed on the ARA and DRACO clusters at Friedrich Schiller University Jena, the supercomputer SuperMUC-NG at the Leibniz-Rechenzentrum (LRZ; www.lrz.de) Munich, and the national HPE Apollo Hawk at the High Performance Computing Center Stuttgart (HLRS). The ARA cluster is funded in part by DFG grants INST 275/334-1 FUGG and INST 275/363-1 FUGG and ERC Starting Grant, grant agreement No. BinGraSp-714626. The authors acknowledge the Gauss Centre for Supercomputing e. V. (www.Gauss-center.eu) for funding this project by providing computing time on the GCS Supercomputer SuperMUC-NG at LRZ (allocations pn36ge and pn36jo). The authors acknowledge HLRS for funding this project by providing access to the supercomputer HPE Apollo Hawk under the grant No. INTRHYGUE/44215.

Appendix A Hydrodynamic Effects on NN

To get an idea of how the prescribed thermodynamic trajectory and the hydrodynamical one differ, we compare in Figure 5 the density ρ , temperature T , and electron fraction Y_e evolutions for the innermost shell of the 12° , 90° angular sections of our SNEC profile. We find orders-of-magnitude discrepancies in ρ and T , while Y_e is generally more compatible between the two evolutions. Note, in the left column, how the jet affects the more polar trajectory, reducing the density and causing a discontinuity in the derivatives of $T(t)$ and $Y_e(t)$. Note that Figure 5 is only meant to work as an example and is not representative of the general behavior of the ejecta. For instance, note in Figure 1 how h settles at different final values (i.e., the expansion becomes homologous) at different times for




Figure 5. Evolution of the density (black), temperature (orange), and electron fraction (green) in the innermost shell of the 12° (left column) and 90° (right column) angular sections of the ejecta as calculated from the complete simulation (solid lines) and an independent *SkyNet* run (dashed lines). The bottom panels show the relative differences calculated with an expression analogous to Equation (5). The evolution of the temperature from *SkyNet* is plotted until it is actively updated by the NN and used in the nuclear calculations. The vertical dashed line at $t = 200$ ms indicates the launch of the jet.

different fluid elements. Better or worse agreement between the compared evolutions can be found looking at other mass shells and angular sections.

Appendix B NN Effects on Hydrodynamics

We discuss here why the introduction of an online NN in a hydrodynamic simulation of the ejecta can affect the kilonova predictions while having negligible effects on the hydrodynamic evolution. To do that, we run a SNEC simulation using the heating rate fits of Z. Wu et al. (2022) instead of the information from a coupled NN. In Figure 6, we compare the total pressure work done on the ejecta and released nuclear energy, as a function of time, against a SNEC run with an online NN. In both simulations, we assume a constant thermalization factor $f_{\text{th}} = 0.5$. The same assumption is made in Z. Wu et al. (2022), as the composition information needed by the detailed thermalization and coming from the NN is missing.

As discussed in Section 3.1, the hydrodynamical evolution of the density profile from our complete run does not agree with the considered common prescription for homologous expansion. The same conclusion holds if the run using the heating rate fits is used. The introduction of a coupled NN introduces global corrections in the heating rates of $\sim 10\%$. If specific fluid elements are considered, these corrections can grow up to a factor 10. Anyway, as shown in Figure 6, the global corrections are typically $\mathcal{O}(10^{-4})$, the total pressure work done by the mass shells during their expansion. As a consequence, the effect of the online NN on their thermodynamic trajectories through Equation (1) is negligible.




Figure 6. Top panel: evolution of total pressure work pdV (blue) and nuclear heating $\epsilon = \dot{\epsilon}_{\text{nuc}} dt$ (orange) averaged all over the ejecta and integrated over time from the SNEC runs with (solid lines) and without (dashed lines) a coupled NN and a constant thermalization factor $f_{\text{th}} = 0.5$. Bottom panel: difference in the nuclear heating between the two simulations normalized over pdV (black line) or ϵ (red line) from the simulation using the heating rate fits.

However, the kilonova light curves are affected by the NN, as the heating rate changes act more directly on the properties of the photosphere and the radiation emitted by the shells outside of it (see Section 3.3).

Appendix C Postprocessing

A. Perego et al. (2022) compute the nucleosynthesis in postprocessing using our same NN and imposing homologically expanding ejecta profiles. A grid on the initial entropy s_0 , expansion timescale τ_0 , and electron fraction $Y_{e,0}$ is introduced to have a series of ready-to-use tabulated results. To adapt their postprocess procedure to our initialization method, we first distribute the mass shells of the initial ejecta profiles (the same as we run with SNEC, averaged over the azimuthal angle) on a grid based on the initial thermodynamic conditions. We take the same grid in s_0 , τ_0 , and $Y_{e,0}$ used in A. Perego et al. (2022) and introduce a fine grid (150 values) on the temperature T_0 to be consistent with our NN initialization procedure. We then run an independent instance of SkyNet on each of the grid points. To be consistent with the homology assumption of D. Radice et al. (2016), we base our grid on the value $\tau_0 = R_E/v_0$, where R_E is the extraction radius and v_0 is the initial velocity; initialize each NN at $t_0 = \tau_0$ and $T = T_0$ imposing NSE; and evolve it with the prescription $\rho(t) = \rho_0((t + \tau_0)/\tau_0)^{-3}$.

The introduction of the grid on s_0 , τ_0 , and $Y_{e,0}$ coarsens the resolution on the initial thermodynamic conditions, thus introducing another layer of approximation in the postprocessing on top of the assumption of homologous expansion. As discussed in Section 3.2, this is also a relevant source of error (note anyway that a step that reduces the number of NNs to run is crucial to define a postprocess procedure able to give results in a reasonable amount of time).

ORCID iDs

Fabio Magistrelli <https://orcid.org/0009-0005-0976-7851>

Sebastiano Bernuzzi <https://orcid.org/0000-0002-2334-0935>

Albino Perego <https://orcid.org/0000-0002-0936-8237>

David Radice <https://orcid.org/0000-0001-6982-1008>

References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017a, *PhRvL*, **119**, 161101
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017b, *ApJL*, **848**, L12
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017c, *ApJL*, **848**, L13
- Arcones, A., & Thielemann, F.-K. 2023, *A&ARv*, **31**, 1
- Barnes, J., Kasen, D., Wu, M.-R., & Martinez-Pinedo, G. 2016, *ApJ*, **829**, 110
- Beloborodov, A. M. 2003, *ApJ*, **588**, 931
- Berger, M., Hubbell, J.H., Seltzer, S.M., et al. 2010, XCOM: Photon Cross Sections Database, NIST Standard Reference Database 8 (XGAM), doi:10.18434/T48G6X, <http://physics.nist.gov/xcom>
- Bernuzzi, S. 2020, *GRGr*, **52**, 108
- Brown, D. A., Chadwick, M. B., Capote, R., et al. 2018, *NDS*, **148**, 1
- Burrows, A. 2013, *RvMP*, **85**, 245
- Camilletti, A., Perego, A., Guercilena, F. M., Bernuzzi, S., & Radice, D. 2024, *PhRvD*, **109**, 063023
- Chiesa, L., Perego, A., & Guercilena, F. M. 2024, *ApJL*, **962**, L24
- Collins, C. E., Bauswein, A., Sim, S. A., et al. 2023, *MNRAS*, **521**, 1858
- Combi, L., & Siegel, D. M. 2023a, *ApJ*, **944**, 28
- Combi, L., & Siegel, D. M. 2023b, *PhRvL*, **131**, 231402
- Coulter, D., Foley, R.J., Kilpatrick, C.D., et al. 2017, *Sci*, **358**, 1556
- Cowan, J. J., Sneden, C., Lawler, J. E., et al. 2021, *RvMP*, **93**, 15002
- Curtis, S., Bosch, P., Mosta, P., et al. 2024, *ApJL*, **961**, L26
- Curtis, S., Miller, J. M., Frohlich, C., et al. 2023, *ApJL*, **945**, L13
- Cybart, R. H., Amthor, A. M., Ferguson, R., et al. 2010, *ApJS*, **189**, 240
- de Jesús Mendoza-Temis, J., Wu, M.-R., Martinez-Pinedo, G., et al. 2015, *PhRvC*, **92**, 055805
- Eichler, D., Livio, M., Piran, T., & Schramm, D. N. 1989, *Natur*, **340**, 126
- Fernández, R., & Metzger, B. D. 2013, *MNRAS*, **435**, 502
- Fernandez, R., Tchekhovskoy, A., Quataert, E., Foucart, F., & Kasen, D. 2019, *MNRAS*, **482**, 3373
- Ghirlanda, G., Salafia, O. S., Paragi, Z., et al. 2019, *Sci*, **363**, 968
- Hamidani, H., Kiuchi, K., & Ioka, K. 2020, *MNRAS*, **491**, 3192
- Hotokezaka, K., & Nakar, E. 2020, *ApJ*, **891**, 152
- Janiuk, A. 2014, *A&A*, **568**, A105
- Janka, H.-T. 2012, *ARNPS*, **62**, 407
- Just, O., Bauswein, A., Pulpillo, R. A., Goriely, S., & Janka, H.-T. 2015, *MNRAS*, **448**, 541
- Just, O., Goriely, S., Janka, H.-T., Nagataki, S., & Bauswein, A. 2021, *MNRAS*, **509**, 1377
- Just, O., Vijayan, V., Xiong, Z., et al. 2023, *ApJL*, **951**, L12
- Kasen, D., & Barnes, J. 2019, *ApJ*, **876**, 128
- Kasen, D., Metzger, B., Barnes, J., Quataert, E., & Ramirez-Ruiz, E. 2017, *Natur*, **551**, 80
- Kiuchi, K., Fujibayashi, S., Hayashi, K., et al. 2023, *PhRvL*, **131**, 011401
- Kiuchi, K., Rebol-Salze, A., Shibata, M., & Sekiguchi, Y. 2024, *NatAs*, **298**, 8
- Korobkin, O., Rosswog, S., Arcones, A., & Winteler, C. 2012, *MNRAS*, **426**, 1940
- Lippuner, J., & Roberts, L. F. 2015, *ApJ*, **815**, 82
- Lippuner, J., & Roberts, L. F. 2017, *ApJS*, **233**, 18
- Martin, D., Perego, A., Arcones, A., et al. 2015, *ApJ*, **813**, 2
- Metzger, B. D., Bauswein, A., Goriely, S., & Kasen, D. 2015, *MNRAS*, **446**, 1115
- Metzger, B. D., & Fernandez, R. 2014, *MNRAS*, **441**, 3444
- Morozova, V., Piro, A. L., Renzo, M., et al. 2015, *ApJ*, **814**, 63
- Musolino, C., Duqué, R., & Rezzolla, L. 2024, *ApJL*, **966**, L31
- Nativi, L., Bulla, M., Rosswog, S., et al. 2020, *MNRAS*, **500**, 1772
- Nedora, V., Bernuzzi, S., Radice, D., et al. 2021, *ApJ*, **906**, 98
- Nicholl, M., Berger, E., Kasen, D., et al. 2017, *ApJL*, **848**, L18
- Paczynski, B. 1986, *ApJL*, **308**, L43
- Perego, A., Bernuzzi, S., & Radice, D. 2019, *EPJA*, **55**, 124
- Perego, A., Radice, D., & Bernuzzi, S. 2017, *ApJL*, **850**, L37
- Perego, A., Thielemann, F. K., & Cescutti, G. 2021, in *Handbook of Gravitational Wave Astronomy*, ed. C. Bambi, S. Katsanevas, & K. D. Kokkotas (Berlin: Springer), 13
- Perego, A., Vescovi, D., Fiore, A., et al. 2022, *ApJ*, **925**, 22
- Prantzos, N., Abia, C., Cristallo, S., Limongi, M., & Chieffi, A. 2020, *MNRAS*, **491**, 1832
- Radice, D., & Bernuzzi, S. 2023, *ApJ*, **959**, 46
- Radice, D., & Bernuzzi, S. 2024, *JPhCS*, **2742**, 012009
- Radice, D., Bernuzzi, S., Perego, A., & Haas, R. 2022, *MNRAS*, **512**, 1499
- Radice, D., Galeazzi, F., Lippuner, J., et al. 2016, *MNRAS*, **460**, 3255
- Radice, D., Perego, A., Hotokezaka, K., et al. 2018a, *ApJL*, **869**, L35
- Radice, D., Perego, A., Hotokezaka, K., et al. 2018b, *ApJ*, **869**, 130

- Ricigliano, G., Perego, A., Borhanian, S., et al. 2024, [MNRAS](#), **529**, 647
- Roberts, L. F., Lippuner, J., Duez, M. D., et al. 2017, [MNRAS](#), **464**, 3907
- Rosswog, S., Feindt, U., Korobkin, O., et al. 2017, [CQGra](#), **34**, 104001
- Rosswog, S., & Korobkin, O. 2024, [AnP](#), **536**, 2200306
- Rosswog, S., Korobkin, O., Arcones, A., Thielemann, F.-K., & Piran, T. 2014, [MNRAS](#), **439**, 744
- Savchenko, V., Ferrigno, C., Kuulkers, E., et al. 2017, [ApJL](#), **848**, L15
- Schianchi, F., Gieg, H., Nedora, V., et al. 2024, [PhRvD](#), **109**, 044012
- Shingles, L. J., Collins, C. E., Vijayan, V., et al. 2023, [ApJL](#), **954**, L41
- Siegel, D. M. 2019, [EPJA](#), **55**, 203
- Siegel, D. M., & Metzger, B. D. 2017, [PhRvL](#), **119**, 231102
- Sprouse, T. M., Lund, K. A., Miller, J. M., McLaughlin, G. C., & Mumpower, M. R. 2024, [ApJ](#), **962**, 79
- Tanaka, M., Utsumi, Y., Mazzali, P.A., et al. 2017, [PASJ](#), **69**, 102
- Tanvir, N. R., Levan, A. J., Gonzalez-Fernandez, C., et al. 2017, [ApJL](#), **848**, L27
- Timmes, F. X., & Swesty, F. D. 2000, [ApJS](#), **126**, 501
- Von Neumann, J., & Richtmyer, R. D. 1950, [JAP](#), **21**, 232
- Wanajo, S. 2018, [ApJ](#), **868**, 65
- Wanajo, S., Sekiguchi, Y., Nishimura, N., et al. 2014, [ApJL](#), **789**, L39
- Wu, Z., Ricigliano, G., Kashyap, R., Perego, A., & Radice, D. 2022, [MNRAS](#), **512**, 328
- Zappa, F., Bernuzzi, S., Radice, D., & Perego, A. 2023, [MNRAS](#), **520**, 1481