

Graviton production in noninflationary cosmology

Ruth Durrer and Massimiliano Rinaldi

Université de Genève, Département de Physique Théorique, 24 quai Ernest Ansermet, CH-1211 Genève 4, Switzerland

(Received 6 January 2009; published 11 March 2009)

We discuss the creation of massless particles in a Universe, which transits from a radiation-dominated era to any other expansion law. We calculate in detail the generation of gravitons during the transition to a matter-dominated era. We show that the resulting gravitons generated in the standard radiation/matter transition are negligible. We use our result to constrain one or more previous matter-dominated era, or any other expansion law, which may have taken place in the early Universe. We also derive a general formula for the modification of a generic initial graviton spectrum by an early matter-dominated era.

DOI: [10.1103/PhysRevD.79.063507](https://doi.org/10.1103/PhysRevD.79.063507)

PACS numbers: 98.80.-k, 04.30.Db, 98.80.Cq

I. INTRODUCTION

One of the most interesting aspect of inflation is that it leads to the generation of a scale-invariant spectrum of scalar perturbations [1] and of gravitational waves [2] (see also [3,4]). The origin of these perturbations is the quantum generation of field correlations in a time-dependent background (scalar field modes for the scalar perturbations and gravitons for the tensor perturbations). Since this generation takes place mainly on superhorizon scales, it is not correct to talk of “particles”. However, long after inflation, when the perturbations reenter the horizon, the particle concept, e.g. for gravitons becomes meaningful and we can calculate, e.g. the energy density of the gravitons which have been generated during inflation.

It is natural to ask whether particle production takes place also in an ordinary expanding but noninflationary Friedmann Universe. The answer is that particle production (or more generically the quantum generation of field correlations) can indeed take place after inflation, but if there is no inflationary phase to start with, the initial vacuum state is in general not known, and the production rate cannot be computed. Examples where particle creation taking place after inflation (or after pre-big bang) modifies the final spectrum are given in Refs. [5–8]. Especially in Ref. [7] it has been studied how inflationary perturbations are modified if the subsequent expansion is not standard radiation but some other expansion law.

In general, the vacuum state, and hence the particle concept, is well defined only if the spacetime is static or very slowly varying [9]. Let us consider a mode of fixed (comoving) frequency k in a Friedmann universe. The above condition then corresponds to $k/\mathcal{H} \gg 1$, where $\mathcal{H} = aH$ is the comoving Hubble scale. In this sense, the scale (wavelength) under consideration must be “inside the horizon”. However, the production of a particle with a given energy k can only take place if the energy scale of expansion is larger or of the order of the energy of the associated mode, i.e. $k \lesssim \mathcal{H}$. Therefore, having a well defined initial vacuum state, and subsequent particle creation, usually requires a decreasing comoving Hubble rate.

This is verified only during inflation or during a collapsing Friedmann Universe, like in the pre-big bang or in bouncing models.

However, one important exception to this general rule exists, and it is the subject of the present paper: in a radiation-dominated Friedmann background, massless perturbations do not couple to the expansion of the Universe, and evolve like in ordinary Minkowski space. This has already been realized and studied to some extent in Ref. [10]. In a radiation-dominated Universe we therefore can provide vacuum initial conditions for all modes of a massless field, including superhorizon modes. Thus, when the expansion law changes, e.g. from radiation- to matter-dominated, the massless modes couple to the expansion of the Universe, and those with $k/\mathcal{H} < 1$ are amplified.

In this paper we study this phenomenon in two situations of interest. In the first, we investigate graviton production during the ordinary radiation/matter transition at redshift $z_{\text{eq}} \approx 3500$. We determine the amplitude and the spectrum of the generated gravity wave background, and we show that the spectrum is flat and the amplitude is negligibly small. In the second case, we investigate the production of gravitons during an arbitrary matter-dominated phase, which could take place in the early Universe, e.g., if a (very weakly interacting) particle becomes massive, and succeeds to dominate the Universe for a period of time before it decays into radiation. We derive a general formula for the gravity wave spectrum generated by any number of such intermediate periods of matter domination. We also determine the gravity waves produced by a transition into an arbitrary other expansion law. Finally, we discuss the modifications of our results which occur when the initial state is not the vacuum but some arbitrary state which may already contain particles. In this work we concentrate on graviton production, but all our results are equally applicable to other massless particles.

The remainder of this paper is organized as follows. In the next section we present the setup and the basic formulas used in our work. In Sec. III, we calculate the gravitational wave production during the standard radiation/matter transition. We also give the results for the transition from

radiation to some generic expansion law. In Sec. IV we consider the effect of one or several additional transitions in the early Universe and we derive results for general nonvacuum initial conditions. In Sec. V we derive the consequences of our results and we draw some conclusions.

Notation: We work in a spatially flat Friedmann Universe, and we denote conformal time by t , so that

$$ds^2 = a^2(t)(-dt^2 + \delta_{ij}dx^i dx^j).$$

An over-dot denotes the derivative with respect to the conformal time. We use natural units $c = \hbar = 1$, except for Newton's constant G , which is related to the reduced Planck mass by $8\pi G = m_p^{-2}$. We normalize the scale factor, so that $a_0 = a(t_0) = 1$ at the present time.

II. GRAVITON CREATION IN COSMOLOGY

We now consider tensor perturbations of the Friedmann metric, namely

$$ds^2 = a^2(t)[-dt^2 + (\delta_{ij} + 2h_{ij})dx^i dx^j],$$

where h_{ij} is a transverse and traceless tensor. In Fourier space we have

$$h_{ij}(\mathbf{k}, t) = h_+(k, t)e_{ij}^{(+)}(\hat{\mathbf{k}}) + h_-(k, t)e_{ij}^{(-)}(\hat{\mathbf{k}}), \quad (1)$$

where $e_{ij}^{(\pm)}(\hat{\mathbf{k}})$ denote the positive and negative helicity polarization tensors, and $k^i e_{ij}^{(\pm)} = 0$. In a perfect fluid background, i.e. if there are no anisotropic stresses, both amplitudes satisfy the same wave equation,

$$\square h = \ddot{h} + 2\frac{\dot{a}}{a}\dot{h} + k^2 h = 0, \quad (2)$$

where $h \equiv h_{\pm}$. This equation of motion is obtained when expanding the gravitational action in a Friedman universe to second order in h ,

$$S + \delta S = -\frac{m_p^2}{2} \int d^4x \sqrt{-(g + \delta g)}(R + \delta R). \quad (3)$$

A brief calculation shows that the lowest (second order) contribution to δS can be written in Minkowski-space canonical form,

$$\delta S^{(2)} = -\frac{1}{2} \int d^4x \left(\partial_\mu \phi \partial^\mu \phi - \frac{\ddot{a}}{a} \phi^2 \right) = \int d^4x \mathcal{L}, \quad (4)$$

if we rescale h as

$$h(\mathbf{x}, t) = \frac{1}{\sqrt{2}m_p a(t)} \phi(\mathbf{x}, t). \quad (5)$$

Equation (4) is the action of a canonical scalar field with time-dependent effective squared mass $m^2(t) = -\ddot{a}/a$. If the expansion of the Universe is slow enough (compared to the frequency of the mode under consideration), then the effective mass is negligible, and the theory describes a

massless scalar field in Minkowski space, which can be quantized according to the usual procedure: we first promote the field to an operator

$$\hat{\phi}(\mathbf{x}, t) = \int \frac{d^3k}{(2\pi)^3} [e^{i\mathbf{k}\cdot\mathbf{x}} \chi_{\mathbf{k}}(t) \hat{b}_{\mathbf{k}} + e^{-i\mathbf{k}\cdot\mathbf{x}} \chi_{\mathbf{k}}^*(t) \hat{b}_{\mathbf{k}}^\dagger], \quad (6)$$

then we impose the commutation rules

$$[b_{\mathbf{k}}, b_{\mathbf{k}'}^\dagger] = (2\pi)^3 \delta^3(\mathbf{k} - \mathbf{k}'),$$

$$[b_{\mathbf{k}}, b_{\mathbf{k}'}] = [b_{\mathbf{k}}^\dagger, b_{\mathbf{k}'}^\dagger] = 0.$$

The field equations derived from the action (4) lead to the mode equation

$$\ddot{\chi}_{\mathbf{k}} + \left[k^2 - \frac{\ddot{a}}{a} \right] \chi_{\mathbf{k}} = 0. \quad (7)$$

Within linearized gravity we can therefore quantize the metric fluctuations, provided the Universe expands adiabatically, by making use of the above rescaling of the amplitude h .

In particular, we now assume that the Universe is initially radiation-dominated, so that $a(t) = t$, and $\ddot{a}/a = 0$, and ϕ represents exactly a massless scalar field in Minkowski space. We consider the vacuum initial conditions for the modes $\chi_{\mathbf{k}}(t)$ as given by

$$\chi_{\mathbf{k}}(t) = \frac{1}{\sqrt{2k}} e^{-ikt}. \quad (8)$$

More general initial conditions will be considered at the end of Sec. IV. The field normalization is determined by the Klein-Gordon norm

$$i\chi^* \overleftrightarrow{\partial}_0 \chi \equiv i(\chi^* \partial_0 \chi - \chi \partial_0 \chi^*) = 1. \quad (9)$$

Then, the field operator $\hat{\phi}$ and its canonically conjugate momentum, $\hat{\Pi} = \partial \mathcal{L} / \partial (\partial_0 \hat{\phi})$ satisfy the canonical commutation relations. The operators $\hat{b}_{\mathbf{k}}$ define the *in* vacuum by $\hat{b}_{\mathbf{k}} |0_{\text{in}}\rangle = 0 \forall \mathbf{k}$. In the following, this is the initial vacuum, void of particles by construction.

Suppose now that at a time $t = t_1$, the Universe changes abruptly from radiation-dominated to another expansion law. Then, the effective squared mass no longer vanishes, and the initial modes, with $kt_1 < 1$ are amplified. Continuity requires that χ and $\dot{\chi}$ match at $t = t_1$, and these conditions determine the Bogoliubov coefficients, which relate the new modes χ_{out} and operators $\hat{b}_{\mathbf{k},\text{out}}$ to the old ones, χ_{in} and $\hat{b}_{\mathbf{k},\text{in}}$ by [9]

$$\chi_{\text{out}}(t) = \alpha \chi_{\text{in}}(t) + \beta \chi_{\text{in}}^*(t), \quad (10)$$

$$\hat{b}_{\text{out}} = \alpha^* \hat{b}_{\text{in}} - \beta^* \hat{b}_{\text{in}}^\dagger. \quad (11)$$

With these relations, we can compute the number density of the particles [11] created at the transition [9]

$$N(k, t_1) = \langle 0_{\text{in}} | \hat{b}_{\mathbf{k},\text{out}}^\dagger \hat{b}_{\mathbf{k},\text{out}} | 0_{\text{in}} \rangle = |\beta|^2. \quad (12)$$

Thus, the energy density $\rho \equiv \langle T^0_0 \rangle$ can be written as

$$\rho_h = \frac{1}{a^4} \int \frac{d^3k}{(2\pi)^3} k N(k, t_1) = \frac{1}{2\pi^2 a^4} \int dk k^3 |\beta|^2, \quad (13)$$

which implies the usual formula [12]

$$\left. \frac{d\rho_h}{d \log k} \right|_{\text{tot}} = \frac{k^4 |\beta|^2}{\pi^2 a^4}, \quad (14)$$

where we have multiplied Eq. (13) by a factor 2 to take into account both polarizations. Note also that k denotes comoving momenta/energy so that we had to divide by a^4 to arrive at the physical energy density. The second quantity of interest is the power spectrum $P_h(k, t)$, defined by

$$4\pi \int \frac{dk}{k} P_h(k, t) = \langle 0_{\text{in}} | h(t, \mathbf{x})^2 | 0_{\text{in}} \rangle. \quad (15)$$

Using Eqs. (5) and (6), we obtain

$$P_h(k, t) = \frac{k^2 |\beta|^2}{(2\pi)^3 m_p^2 a^2}, \quad (16)$$

where, again, we have multiplied by 2 to account for both polarizations.

Note that for all this it is not important that we consider a spin 2 graviton. The exactly same mode equation is obtained for a scalar field and also for a fermion field. In the latter case, the commutation relations have to be replaced by the corresponding anti-commutation relations.

III. FROM RADIATION TO MATTER ERA

Before discussing a transition from the radiation-dominated era to the matter era, let us consider the transition from radiation to some generic power law expansion phase, $a \propto t^q$ with $q \neq 1$ at some time t_1 . In the new era $\ddot{a}/a = q(q-1)/t^2 \neq 0$. Note that only if $q > 1$ (or if $q < 0$ which corresponds to inflation or contraction), so that $m^2(t) = -q(q-1)/t^2$ is negative, we will have significant particle production. Since the expansion law is related to the equation of state parameter $w = P/\rho$ via [4]

$$q = \frac{2}{1+3w}, \quad (17)$$

this requires $w < 1/3$.

Let us start in the vacuum during the radiation era, then χ is given entirely by the negative frequency modes, Eq. (8). This means that we consider the situation where there are no significant gravity waves present from an earlier inflationary epoch. Here, we really want to study the production due solely to the radiation/matter transition. The general solution of the mode Eq. (7) in the new era are the spherical Hankel functions [13] of order $q-1$,

$$\chi_{\mathbf{k}}(t) = \frac{\alpha_1}{\sqrt{2k}} z h_{q-1}^{(2)}(z) + \frac{\beta_1}{\sqrt{2k}} z h_{q-1}^{(1)}(z), \quad (18)$$

where $z = kt$. Note that inside the horizon, i.e. for $z \gg 1$,

$z h_{q-1}^{(2)}(z) \propto \exp(-iz)$ corresponds to the negative frequency modes while $z h_{q-1}^{(1)}(z) \propto \exp(iz)$ corresponds to positive frequency modes. We match χ and $\dot{\chi}$ at $t = t_1$ to the radiation-dominated vacuum solution (8). A brief calculation yields the coefficients ($z_1 \equiv kt_1$)

$$\alpha_1 = -\frac{i}{2} e^{-iz_1} [(iz_1 + q) h_{q-1}^{(1)}(z_1) - z_1 h_q^{(1)}(z_1)], \quad (19)$$

$$\beta_1 = -\frac{i}{2} e^{-iz_1} [(iz_1 + q) h_{q-1}^{(2)}(z_1) - z_1 h_q^{(2)}(z_1)]. \quad (20)$$

This instantaneous matching condition is good enough for frequencies for which the transition is rapid, i.e. $z_1 \ll 1$. In fact, for frequencies with $z_1 > 1$, the transition is adiabatic and no particle creation will take place. This can also be seen when considering the limits of the above result for large z_1 . Then $\alpha_1 \rightarrow 1$ and $\beta_1 \rightarrow 0$, but strictly speaking the above approximations are not valid in this regime where no particle creation takes place. We therefore concentrate on $z_1 \ll 1$.

Let us now study the specific case of the radiation-matter transition, i.e. $q = 2$. Then we have to consider spherical Hankel functions of order 1 and the solution is given by

$$\chi_{\mathbf{k}}(t) = \frac{\alpha_1}{\sqrt{2k}} \frac{z-i}{z} e^{-iz} + \frac{\beta_1}{\sqrt{2k}} \frac{z+i}{z} e^{iz}. \quad (21)$$

The matching at t_1 now yields

$$\alpha_1 = 1 + \frac{i}{z_1} - \frac{1}{2z_1^2}, \quad \beta_1 = -\frac{1}{2z_1^2} e^{-2iz_1}. \quad (22)$$

We want to evaluate the quantum field $\hat{\phi}$ at late time, when $z \gg 1$ and the mode k under consideration is sub-horizon. Then, the solution (21) is again the Minkowski solution,

$$\chi_{\mathbf{k}}(t) \simeq \frac{\alpha_1}{\sqrt{2k}} e^{-iz} + \frac{\beta_1}{\sqrt{2k}} e^{iz}. \quad (23)$$

The number of gravitons generated during the matter era (before $z \gg 1$) is, see Eq. (12)

$$N(k, t) = |\beta_1|^2. \quad (24)$$

The graviton power spectrum is given by Eq. (16), and the energy density by Eq. (13).

Using that $\rho_{\text{rad}} a^4 \equiv \rho_{\text{rad}}(t_1) a_1^4 = \frac{3}{2} m_p^2 \mathcal{H}_1^2 a_1^2$ and $|\beta_1|^2 = z_1^{-4}/4$, we obtain

$$\begin{aligned} \frac{d\Omega_h(k)}{d \log k} &= \frac{2\Omega_{\text{rad}}}{3\pi^2} \frac{k^4}{m_p^2 \mathcal{H}_1^2 a_1^2} |\beta_1|^2 = \frac{\Omega_{\text{rad}}}{6\pi^2} \left(\frac{H_1}{m_p} \right)^2 \\ &= \Omega_{\text{rad}} \frac{g_{\text{eff}}}{36 \times 30} \left(\frac{T_1}{m_p} \right)^4. \end{aligned} \quad (25)$$

For the second equal sign we have used that $\mathcal{H}_1 = 1/t_1 = a_1 H_1$, which is strictly true only in the radiation era, in the matter era we have $\mathcal{H} = 2/t$ and at the transition a value

between 1 and 2 would probably be more accurate. But within our approximation of an instant transition, we do not bother about such factors. For the last equal sign we used $H_1^2 = \rho_{\text{rad}}/(6m_p^2)$ with $\rho_{\text{rad}} = g_{\text{eff}} \frac{\pi^2}{30} T^4$ where $g_{\text{eff}} = N_B + \frac{7}{8} N_F$ is the effective number of degrees of freedom.

For a generic transition we obtain $|\beta_1|^2 = z_1^{-2q}/4$ so that

$$\frac{d\Omega_h(k)}{d \log k} \simeq \Omega_{\text{rad}} \left(\frac{T_1}{m_p} \right)^4 z_1^{4-2q}. \quad (26)$$

This spectrum is blue (i.e. growing with k) if $q < 2$ and red otherwise.

As $z_1 < 1$ in the regime of validity of our formula, we need a red spectrum i.e. $q > 2$ to enhance the gravitational wave energy density with respect to the result from the radiation \rightarrow matter transition. According to Eq. (17), this requires $-1/3 < w < 0$, a slightly negative pressure, but still noninflationary expansion.

In the standard radiation to matter transition when three species of left handed neutrinos and the photon are the only relativistic degrees of freedom, we have $g_{\text{eff}} = 29/4$. For this transition

$$T_1 = T_{\text{eq}} \simeq 0.85 \text{ eV} \simeq 0.35 \times 10^{-27} m_p,$$

hence the result (25) is completely negligible.

IV. MORE THAN ONE RADIATION-MATTER TRANSITION

We now consider an early matter-dominated era. At some high temperature $T_1 \gg T_{\text{eq}}$, corresponding to a comoving time t_1 , a massive particle may start to dominate the Universe and render it matter-dominated. At some later time t_2 , corresponding to temperature T_2 , this massive particle decays and the Universe becomes radiation-dominated again, until the usual radiation-matter transition, which takes place at $T_{\text{eq}} \equiv T_3$. We want to determine the gravitational wave spectrum and the spectral density parameter $d\Omega_h/d \log(k)$ as functions of T_1 and T_2 .

Let us first again start with the vacuum state in the radiation era before t_1 . When, we just obtain the results (22) for the Bogoliubov coefficients α_1 and β_1 after the first transition. To evaluate the matching conditions at the second transition, matter to radiation, we set

$$\chi = \frac{\alpha_2}{\sqrt{2k}} e^{-ikt} + \frac{\beta_2}{\sqrt{2k}} e^{ikt}, \quad t \geq t_2.$$

Matching χ and $\dot{\chi}$ at t_2 we can relate the new coefficients α_2 and β_2 to α_1 and β_1 . A brief calculation gives

$$\begin{aligned} \alpha_2 &= \alpha_1 \left(1 - \frac{i}{z_2} - \frac{1}{2z_2^2} \right) + \frac{\beta_1}{2z_2^2} e^{2iz_2} \\ &= \alpha_1 f(z_2) + \beta_1 g(z_2), \end{aligned} \quad (27)$$

$$\beta_2 = \beta_1 \left(1 + \frac{i}{z_2} - \frac{1}{2z_2^2} \right) + \frac{\alpha_1}{2z_2^2} e^{-2iz_2} \quad (28)$$

$$= \beta_1 \bar{f}(z_2) + \alpha_1 \bar{g}(z_2), \quad (29)$$

or in matrix notation

$$\begin{pmatrix} \alpha_2 \\ \beta_2 \end{pmatrix} = M(z_2) \begin{pmatrix} \alpha_1 \\ \beta_1 \end{pmatrix}, \quad \text{with} \quad (30)$$

$$M(z) = \begin{pmatrix} f(z) & g(z) \\ \bar{g}(z) & \bar{f}(z) \end{pmatrix} \quad (31)$$

$$M^{-1}(z) = \begin{pmatrix} \bar{f}(z) & -g(z) \\ -\bar{g}(z) & f(z) \end{pmatrix}. \quad (32)$$

The fact that $M \in SI(2, \mathbb{C})$, i.e., $|f(z)|^2 - |g(z)|^2 = 1$ ensures that the normalization condition (9) which translates to the condition $|\alpha|^2 - |\beta|^2 = 1$ for the Bogoliubov coefficients of a free field, is maintained at the transition. Finally, the matching at the usual radiation-matter transition yields

$$\alpha_3 = \alpha_2 \left(1 + \frac{i}{z_3} - \frac{1}{2z_3^2} \right) - \frac{\beta_2}{2z_3^2} e^{2iz_3}, \quad (33)$$

$$\beta_3 = \beta_2 \left(1 - \frac{i}{z_3} - \frac{1}{2z_3^2} \right) - \frac{\alpha_2}{2z_3^2} e^{-2iz_3},$$

$$\begin{pmatrix} \alpha_3 \\ \beta_3 \end{pmatrix} = M^{-1}(z_3) \begin{pmatrix} \alpha_2 \\ \beta_2 \end{pmatrix} \quad (34)$$

$$\begin{pmatrix} \alpha_3 \\ \beta_3 \end{pmatrix} = M^{-1}(z_3) M(z_2) M^{-1}(z_1) \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (35)$$

To obtain the power spectrum and energy density in this case, we simply have to replace $|\beta|^2$ in Eqs. (16) and (14) by $|\beta_3|^2$. In Fig. 1 we plot $|\beta_3|^2$ as a function of z_3 for different choices of t_2 . The instantaneous transition approximation breaks down for $z_3 > 1$, hence only the left side of the vertical line is physical. For the right side one would have to solve the mode equation numerically, but since we know that particle production is suppressed for these frequencies, we do not consider them. We concentrate on $z_3 \leq 1$. For these wave numbers, also $z_1 < z_2 < z_3 < 1$.

This allows the following approximations,

$$\alpha_1 \simeq \beta_1 \simeq -\frac{1}{2z_1^2}, \quad z_1 \ll 1 \quad (36)$$

$$\alpha_2 \simeq -\beta_2 \simeq \frac{iz_2}{3z_1^2}, \quad z_2 \ll 1 \quad (37)$$

$$\alpha_3 \simeq \beta_3 \simeq -\frac{2z_2}{3z_1^2 z_3}, \quad z_3 \ll 1. \quad (38)$$

To obtain the results (37) and (38) we have to expand the




FIG. 1 (color online). The Bogoliubov coefficient $|\beta_3(k)|^2$ for $t_1 = 1$, and $t_3 = 2000$, with various values of t_2 . Only the k -values left of the vertical dashed line satisfy $kt_3 < 1$. They show clearly a k^{-4} slope and the amplitude is well approximated by (38).

exact expression (27) to fourth order and (33) to second order, but we consider only the largest term in the result given above, using also $z_1 < z_2 < z_3$. Therefore, in the approximate expression (37), where we have neglected a term proportional to z_1/z_2^2 , one no longer sees that $\alpha_2 \rightarrow 1$ and $\beta_2 \rightarrow 0$ when $t_2 \rightarrow t_1$ and hence $z_2 \rightarrow z_1$. In this case there is no intermediate matter-dominated era and therefore no particle creation, hence $\beta_2 = 0$. This can be seen from the exact expression given in Eq. (27).

Within these approximations, Eqs. (16) and (14) lead to

$$P_h(k) = \frac{1}{(2\pi)^3} \left(\frac{t_2}{m_p a t_1^2 t_3 k} \right)^2, \quad kt_3 < 1 \quad (39)$$

$$\begin{aligned} \frac{d\Omega_h}{d \log k} &= \frac{\Omega_{\text{rad}}}{\pi^2} \frac{k^4 |\beta_3|^2}{a_1^4 \rho_{\text{rad}}(t_1)} \\ &= \Omega_{\text{rad}} \frac{g_{\text{eff}}(T_1)}{18 \times 45} \left(\frac{T_1}{m_p} \right)^4 \left(\frac{T_{\text{eq}}}{T_2} \right)^2, \quad kt_{\text{eq}} < 1. \end{aligned} \quad (40)$$

This result can be generalized to several, say N , intermediate radiation \rightarrow matter transitions at times t_{2n-1} and back to radiation at time t_{2n} , $1 \leq n \leq N$, with the result

$$|\beta_{2N+1}|^2 \simeq \frac{1}{(kt_1)^4} \left(\frac{T_3 \cdots T_{2N-1} T_{\text{eq}}}{T_2 \cdots T_{2N}} \right)^2. \quad (41)$$

Hence, each return to the radiation-dominated era at some intermediate temperature T_{2n} leads to a suppression factor $(T_{2n+1}/T_{2n})^2$, where T_{2n+1} denotes the temperature at the start of the next matter era.

On large scales, $kt_{\text{eq}} < 1$, the energy density spectrum is flat. The best constraints on an intermediate radiation-dominated era therefore come from the largest scales, i.e. from observations of the cosmic microwave background (CMB) as we shall discuss in the next section.

We now briefly consider the case when the initial conditions differ from the vacuum case, Eq. (8). We assume an arbitrary initial state of the field given by

$$\chi_{\mathbf{k}}(t) = \frac{\alpha_0}{\sqrt{2k}} e^{-ikt} + \frac{\beta_0}{\sqrt{2k}} e^{ikt}, \quad (42)$$

together with the normalization condition which ensures that the field is canonically normalized, $|\alpha_0|^2 - |\beta_0|^2 = 1$. The same calculations as above now yield

$$\begin{pmatrix} \alpha_3 \\ \beta_3 \end{pmatrix} = M^{-1}(z_3) M(z_2) M^{-1}(z_1) \begin{pmatrix} \alpha_0 \\ \beta_0 \end{pmatrix}, \quad (43)$$

where $M(z)$ is the matrix giving the transition from matter to radiation defined in Eq. (31).

Expanding this in z_1 , z_2 and z_3 , using $z_1 < z_2 < z_3 < 1$ one finds that to lowest nonvanishing order, the final result for β_3 depends only $|\alpha_0 + \beta_0|$. However, if the phase of α_0 and β_0 are nearly opposite, i.e., $\alpha_0 \simeq -\beta_0$, and if $|\alpha_0|$ and therefore also $|\beta_0|$ are much larger than 1, a correction proportional to $|\alpha_0 - \beta_0|$ becomes important. More precisely, the last of Eqs. (36) now is replaced by

$$\beta_3 \simeq \frac{2z_2}{3z_1 z_3} \left[-\frac{1}{z_1} (\alpha_0 + \beta_0) + 2i(\alpha_0 - \beta_0) \right]. \quad (44)$$

If $\alpha_0 = 1$ and $\beta_0 = 0$, the second term can be neglected with respect to the first one and we reproduce the previous result (38). As we see from this equation, a large phase difference between α_0 and β_0 changes not only the amplitude but also the slope of the spectrum. Of course in concrete examples, like for a previous inflationary period, see Ref. [7], the coefficients α_0 and β_0 also depend on the wave number.

In Fig. 2 we show the dependence of $|\beta_3|^2$ on $|\alpha_0|$ for different values of the relative phase between α_0 and β_0 (top panel) and as a function of the relative phase for different values of $|\alpha_0|$. The difference of $|\beta_3|^2$ between the case where α_0 and β_0 are perfectly in phase and of opposite phase is of the order of $1/z_1$, if $|\alpha_0|$ is significantly larger than 1. This is already evident from Eq. (44). In the Fig. 2 we have chosen $z_1 = 0.1$, a unrealistically high value, in order to have a better visibility of the phase dependence which then changes $|\beta_3|^2$ only by 1 order of magnitude.

In conclusion, in the case of a nonvacuum initial state, $|\alpha_0|$ significantly larger than 1, graviton production is enhanced typically by a factor of order $|\alpha_0|^2 \sim |\beta_0|^2$ which is the number of initial particles. Hence in addition to the spontaneous creation we now also have induced particle creation which is proportional to the initial particle number and much larger than the spontaneous creation if




FIG. 2 (color online). In the top panel we show $|\beta_3|^2$ as a function of α_0 for α_0 and β_0 in phase (top, solid, black line), of opposite phase (lowest, dashed, red line) and with a phase difference of 0.8π (middle, dot-dashed, blue line). In the bottom panel we show $|\beta_3|^2$ as a function of the phase difference ψ for $|\alpha_0| = 80$ (top line) $|\alpha_0| = 40$ (middle line) and $|\alpha_0| = 10$ (lowest line). We have chosen $z_1 = 0.1$ in this plot and the overall vertical normalization is in units of $(2z_2/(z_1z_3))^2$.

the particle number is large. An interesting point is that the phase shift between α_0 and β_0 can significantly affect the final spectrum.

V. DISCUSSION AND CONCLUSIONS

The fact that the observed CMB anisotropies are of the order of 10^{-5} yields a strong limit on gravitational waves with wave numbers of the order of the present Hubble

scale, see e.g. [14].

$$\left. \frac{d\Omega_h}{d\log k} \right|_{k=H_0} < 10^{-15}. \quad (45)$$

With $\Omega_{\text{rad}} \approx 10^{-5}$, and Eq. (40), this implies the limit

$$\left(\frac{T_1}{m_p} \right)^2 \frac{T_{\text{eq}}}{T_2} < 10^{-5}. \quad (46)$$

We know that during nucleosynthesis the Universe was radiation-dominated, hence $T_2 \geq 0.1$ MeV. With $T_{\text{eq}} \sim 1$ eV, the above inequality reduces to $T_1 < m_p$ for the value $T_2 \approx 0.1$ MeV, and it is even less stringent for higher values of T_2 . Hence even though the production of gravitons during an intermediate matter era is of principal interest, we cannot derive stringent limits on T_1 and T_2 . On the other hand, for values of T_1 and T_2 close to the maximal, respectively, minimal value, $T_1 \sim m_p$ and $T_2 \sim 0.1$ MeV, these gravitons would leave a detectable signature in the cosmic microwave background.

We can, however, use this effect to limit any intermediate era with $q > 2$, i.e. $-1/3 < w = P/\rho < 0$. According to Eq. (26), in the general case, the particle number is of the order of $|\beta_1|^2 \approx z_1^{-2q}$, so that

$$\frac{d\Omega_h}{d\log k} \approx \Omega_{\text{rad}}(kt_1)^{4-2q} \left(\frac{T_1}{m_p} \right)^4 \left(\frac{T_{\text{eq}}}{T_2} \right)^2, \quad z_1 \ll 1. \quad (47)$$

As above, T_2 denotes the temperature at which the Universe returns to the radiation-dominated state, hence nucleosynthesis requires $T_2 > 0.1$ MeV. In this case, if $q > 2$, the spectrum becomes red and, at $k \sim H_0$, the limit can become quite interesting. Hence, graviton production can significantly limit a (noninflationary) phase with negative pressure in the early universe. For $k = H_0$, using $t_1 = \mathcal{H}_1^{-1} = 1/(a_1 H_1)$, we obtain

$$(H_0 t_1)^{-1} \approx \left(\frac{g_{\text{eff}}(T_1)}{10^5} \right)^{1/2} \left(\frac{T_1}{T_0} \right)^3,$$

where $T_0 \approx 0.2 \times 10^{-4}$ eV is the present temperature of the Universe. This can be a significant factor for large values of T_1 . Inserting this expression in Eq. (47), the limit (45) yields

$$\left(\frac{g_{\text{eff}}(T_1)}{10^5} \right)^{q-2} \left(\frac{T_1}{T_0} \right)^{6(q-2)} \left(\frac{T_1}{m_p} \right)^4 \left(\frac{T_{\text{eq}}}{T_2} \right)^2 \leq 10^{-10}. \quad (48)$$

For example, for $w = -1/21$, ($q = 7/3$), we already obtain

$$\frac{T_1}{T_2} \left(\frac{T_1}{m_p} \right)^2 \leq 10^{-9}. \quad (49)$$

E.g. for $T_2 = 1$ MeV this implies $T_1 < 10^8$ GeV. For smaller values of w the limit becomes more stringent.

In this paper we have shown that there is cosmological particle production of massless modes whenever the expansion law is not radiation-dominated so that $\ddot{a}/a \neq 0$.

This term acts like a time-dependent mass and leads to the production of modes with comoving energy $k^2 < |\ddot{a}/a| \simeq \mathcal{H}^2$, hence with physical energy $\omega = k/a \lesssim H$. One readily sees that particle production is significant only if $q < 1$, i.e. the squared mass $-\ddot{a}/a < 0$. When starting from a radiation-dominated Universe, we do have a well defined initial vacuum state also for the superhorizon modes which can be amplified by a transition to another expansion law. We have also given the expressions for the produced particle number in the case of a generic, nonvacuum initial state, Eq. (44). This can be applied to an arbitrary inflationary, pre-big bang or bouncing model, which may already contain gravitons before the first radiation era.

We have explicitly calculated the production of gravitons for a vacuum initial state and have arrived at the following main conclusions:

- (i) The gravitons produced after the standard radiation-matter transition are completely negligible.

- (ii) If we introduce a matter-dominated era in the early universe, this leads to a flat energy spectrum of gravity waves which, in the most optimistic case, can be sufficient to contribute to the CMB tensor anisotropies in an observable way.
- (iii) A phase of expansion with $-1/3 < w < 0$ leads to a red spectrum of gravitons. Such a phase in the early universe is severely constrained mainly by the amplitude of CMB anisotropies.
- (iv) A graviton spectrum present at the beginning of the radiation era can become significantly amplified and modified by intermediate not-standard evolution of the universe.

ACKNOWLEDGMENTS

We thank John Barrow for valuable comments. This work is supported by the Swiss National Science Foundation.

-
- [1] V. Mukhanov and G. Chibisov, JETP **56**, 258 (1982).
 - [2] A. A. Starobinskii, JETP Lett. **30**, 682 (1979).
 - [3] V. Mukhanov, *Physical Foundations of Cosmology* (Cambridge University Press, Cambridge, England, 2005).
 - [4] R. Durrer, *The Cosmic Microwave Background* (Cambridge University Press, Cambridge, England, 2008).
 - [5] R. Durrer, M. Gasparini, M. Sakellariadou, and G. Veneziano, Phys. Lett. B **436**, 66 (1998).
 - [6] R. Durrer, M. Gasparini, M. Sakellariadou, and G. Veneziano, Phys. Rev. D **59**, 043511 (1999).
 - [7] V. Sahni, Phys. Rev. D **42**, 453 (1990).
 - [8] V. Sahni, Phys. Rev. D **65**, 023518 (2001).
 - [9] N. D. Birrell and P. C. W. Davies, *Quantum Fields In Curved Space* (Cambridge University Press, Cambridge, England, 1982).
 - [10] L. P. Grishchuk, Sov. Phys. JETP **67**, 825 (1974) [Sov. Phys. JETP **40**, 409 (1975)].
 - [11] We use the notion “particle” in a somewhat sloppy way. These modes are particles in the standard sense of the term only once the mode has entered the horizon. Only then $N(k, t)$ is really a particle number, before it has rather to be related to the square amplitude of field correlations. Of course there is no way of measuring fluctuations with wavelengths larger than the horizon scale.
 - [12] Again, this is a physical graviton energy density only for scales well inside the horizon.
 - [13] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions* (Dover Publications, New York, 1972), 9th ed..
 - [14] L. A. Boyle and A. Buonanno, Phys. Rev. D **78**, 043531 (2008).