

ON STRASSEN'S RANK ADDITIVITY FOR SMALL THREE-WAY TENSORS *

JAROSŁAW BUCZYŃSKI[†], ELISA POSTINGHEL[‡], AND FILIP RUPNIEWSKI[§]

Abstract. We address the problem of the additivity of the tensor rank. That is, for two independent tensors we study if the rank of their direct sum is equal to the sum of their individual ranks. A positive answer to this problem was previously known as Strassen's conjecture until recent counterexamples were proposed by Shitov. The latter are not very explicit, and they are only known to exist asymptotically for very large tensor spaces. In this article we prove that for some small three-way tensors the additivity holds. For instance, if the rank of one of the tensors is at most 6, then the additivity holds. Or, if one of the tensors lives in $\mathbb{C}^k \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ for any k , then the additivity also holds. More generally, if one of the tensors is concise and its rank is at most 2 more than the dimension of one of the linear spaces, then additivity holds. In addition we also treat some cases of the additivity of the border rank of such tensors. In particular, we show that the additivity of the border rank holds if the direct sum tensor is contained in $\mathbb{C}^4 \otimes \mathbb{C}^4 \otimes \mathbb{C}^4$. Some of our results are valid over an arbitrary base field.

Key words. Tensor rank, additivity of tensor rank, Strassen's conjecture, slices of tensor, secant variety, border rank.

AMS subject classifications. Primary: 15A69, Secondary: 14M17, 68W30, 15A03.

1. Introduction. The matrix multiplication is a bilinear map $\mu_{i,j,k}: \mathcal{M}^{i \times j} \times \mathcal{M}^{j \times k} \rightarrow \mathcal{M}^{i \times k}$, where $\mathcal{M}^{l \times m}$ is the linear space of $l \times m$ matrices with coefficients in a field \mathbb{k} . In particular, $\mathcal{M}^{l \times m} \simeq \mathbb{k}^{l \cdot m}$, where \simeq denotes an isomorphism of vector spaces. We can interpret $\mu_{i,j,k}$ as a three-way tensor

$$\mu_{i,j,k} \in (\mathcal{M}^{i \times j})^* \otimes (\mathcal{M}^{j \times k})^* \otimes \mathcal{M}^{i \times k}.$$

Following the discoveries of Strassen [30], scientists started to wonder what is the minimal number of multiplications required to calculate the product of two matrices MN , for any $M \in \mathcal{M}^{i \times j}$ and $N \in \mathcal{M}^{j \times k}$. This is a question about the *tensor rank* of $\mu_{i,j,k}$.

Suppose A , B , and C are finite dimensional vector spaces over \mathbb{k} . A *simple tensor* is an element of the tensor space $A \otimes B \otimes C$ which can be written as $a \otimes b \otimes c$ for some $a \in A$, $b \in B$, $c \in C$. The rank of a tensor $p \in A \otimes B \otimes C$ is the minimal number $R(p)$ of simple tensors needed, such that p can be expressed as a linear combination of simple tensors. Thus $R(p) = 0$ if and only if $p = 0$, and $R(p) = 1$ if and only if p is a simple tensor. In general, the higher the rank is, the more complicated p "tends" to be. In particular, the minimal number of multiplications needed to calculate MN as above is equal to $R(\mu_{i,j,k})$. See for instance [17], [22], [13] and references therein for more details and further motivations to study tensor rank.

Our main interest in this article is in the *additivity* of the tensor rank. Going on with the main example, given arbitrary four matrices $M' \in \mathcal{M}^{i' \times j'}$, $N' \in \mathcal{M}^{j' \times k'}$, $M'' \in \mathcal{M}^{i'' \times j''}$, $N'' \in \mathcal{M}^{j'' \times k''}$, suppose we want to calculate both products $M'N'$ and $M''N''$ simultaneously. What is the minimal number of multiplications needed to obtain the result? Is it equal to the sum of the ranks $R(\mu_{i',j',k'}) + R(\mu_{i'',j'',k''})$? More generally, the same question can be asked for arbitrary tensors. If we are given two tensors in independent vector spaces, is the rank of their sum equal to the sum of their ranks? A positive answer to this question was widely known as Strassen's Conjecture [31, p. 194, §4, Vermutung 3], [22, §5.7], until it was disproved by Shitov [29].

PROBLEM 1 (Strassen's additivity problem). *Suppose $A = A' \oplus A''$, $B = B' \oplus B''$, and $C = C' \oplus C''$, where all A, \dots, C'' are finite dimensional vector spaces over a field \mathbb{k} . Pick*

*Submitted to the editors DATE.

[†] Institute of Mathematics of the Polish Academy of Sciences, ul. Śniadeckich 8, 00-656 Warsaw, Poland, and Faculty of Mathematics, Computer Science and Mechanics, University of Warsaw, ul. Banacha 2, 02-097 Warsaw, Poland (jabu@mimuw.edu.pl)

[‡] Department of Mathematical Sciences, Loughborough University, Leicestershire LE11 3TU, United Kingdom (e.postinghel@lboro.ac.uk)

[§] Institute of Mathematics of the Polish Academy of Sciences, ul. Śniadeckich 8, 00-656 Warsaw, Poland (f.rupniewski@impan.pl)

$p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ and let $p = p' + p''$, which we will write as $p = p' \oplus p''$. Does the following equality hold

$$(1.1) \quad R(p) = R(p') + R(p'')?$$

In this article we address several cases of Problem 1 and its generalisations. It is known that if one of the vector spaces $A', A'', B', B'', C', C''$ is at most two dimensional, then the additivity of the tensor rank (1.1) holds: see [21] for the original proof and Section 3.2 for a discussion of more recent approaches. One of our results includes the next case, that is, if say $\dim B'' = \dim C'' = 3$, then (1.1) holds. The following theorem summarises our main results.

THEOREM 1.2. *Let \mathbb{k} be any base field and let $A', A'', B', B'', C', C''$ be vector spaces over \mathbb{k} . Assume $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ and let*

$$p = p' \oplus p'' \in (A' \oplus A'') \otimes (B' \oplus B'') \otimes (C' \oplus C'').$$

If at least one of the following conditions holds, then the additivity of the rank holds for p , that is, $R(p) = R(p') + R(p'')$:

- $\mathbb{k} = \mathbb{C}$ or $\mathbb{k} = \mathbb{R}$ (complex or real numbers) and $\dim B'' \leq 3$ and $\dim C'' \leq 3$.
- $R(p'') \leq \dim A'' + 2$ and p'' is not contained in $\tilde{A}'' \otimes B'' \otimes C''$ for any linear subspace $\tilde{A}'' \subsetneq A''$ (this part of the statement is valid for any field \mathbb{k}).
- $\mathbb{k} = \mathbb{R}$ or \mathbb{k} is an algebraically closed field of characteristic $\neq 2$ and $R(p'') \leq 6$.

Analogous statements hold if we exchange the roles of A, B, C and/or of $'$ and $''$.

The theorem summarises the content of Theorems 4.17–4.19 proven in Section 4.4.

REMARK 1.3. *Although most of our arguments are characteristic free, we partially rely on some earlier results which often are proven only over the fields of the complex or the real numbers, or other special fields. Specifically, we use upper bounds on the maximal rank of small tensors, such as [6] or [33]. See Section 4.4 for a more detailed discussion. In particular, the consequence of the proof of Theorem 4.19 is that if (over any field \mathbb{k}) there are p' and p'' such that $R(p'') \leq 6$ and $R(p' \oplus p'') < R(p') + R(p'')$, then $p'' \in \mathbb{k}^3 \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$ and $R(p'') = 6$. In [6] it is shown that if $\mathbb{k} = \mathbb{Z}_2$ (the field with two elements), then such tensors p'' with $R(p'') = 6$ exist.*

Some other cases of additivity were shown in [18]. Another variant of Problem 1 asks the same question in the setting of symmetric tensors and the symmetric tensor rank, or equivalently, for homogeneous polynomials and their Waring rank. No counterexamples to this version of the problem are yet known, while some partial positive results are described in [10], [11], [12], [14], and [34]. Possible ad hoc extensions to the symmetric case of the techniques and results obtained in this article are subject of a follow-up research.

Next we turn our attention to the *border rank*. Roughly speaking, over the complex numbers, a tensor p has border rank at most r , if and only if it is a limit of tensors of rank at most r . The border rank of p is denoted by $\underline{R}(p)$. One can pose the analogue of Problem 1 for the border rank: for which tensors $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ is the border rank additive, that is, $\underline{R}(p' \oplus p'') = \underline{R}(p') + \underline{R}(p'')$?

In general, the answer is negative; in fact there exist examples for which $\underline{R}(p' \oplus p'') < \underline{R}(p') + \underline{R}(p'')$: Schönhage [28] proposed a family of counterexamples amongst which the smallest is

$$\underline{R}(\mu_{2,1,3}) = 6, \quad \underline{R}(\mu_{1,2,1}) = 2, \quad \underline{R}(\mu_{2,1,3} \oplus \mu_{1,2,1}) = 7,$$

see also [22, §11.2.2].

Nevertheless, one may be interested in special cases of the problem. We describe one instance suggested by J. Landsberg (private communication, also mentioned during his lectures at Berkeley in 2014).

PROBLEM 2 (Landsberg). *Suppose A', B', C' are vector spaces and $A'' \simeq B'' \simeq C'' \simeq \mathbb{C}$. Let $p' \in A' \otimes B' \otimes C'$ be any tensor and $p'' \in A'' \otimes B'' \otimes C''$ be a non-zero tensor. Is $\underline{R}(p' \oplus p'') > \underline{R}(p')$?*

Another interesting question is what is the smallest counterexample to the additivity of the border rank? The example of Schönhage lives in $\mathbb{C}^{2+2} \otimes \mathbb{C}^{3+2} \otimes \mathbb{C}^{6+1}$, that is, it requires using a seven dimensional vector space. Here we show that if all three spaces A, B, C have dimensions at most 4, then it is impossible to find a counterexample to the additivity of the border rank.

THEOREM 1.4. *Suppose $A', A'', B', B'', C', C''$ are vector spaces over \mathbb{C} and $A = A' \oplus A''$, $B = B' \oplus B''$, and $C = C' \oplus C''$. If $\dim A, \dim B, \dim C \leq 4$, then for any $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ the additivity of the border rank holds:*

$$\underline{R}(p' \oplus p'') = \underline{R}(p') + \underline{R}(p'').$$

We prove the theorem in Section 5 as Corollary 5.2, Propositions 5.13 and 5.14, which in fact cover a wider variety of cases.

1.1. Overview. In this article, for the sake of simplicity, we mostly restrict our presentation to the case of three-way tensors, even though some intermediate results hold more generally. In Section 2 we introduce the notation and review known methods about tensors in general. We review the translation of the rank and border rank of three-way tensors into statements about linear spaces of matrices. In Proposition 2.10 we explain that any decomposition that uses elements outside of the minimal tensor space containing a given tensor must involve more terms than the rank of that tensor. In Section 3 we present the notation related to the direct sum tensors and we prove the first results on the additivity of the tensor rank. In particular, we slightly generalise the proof of the additivity of the rank when one of the tensor spaces has dimension 2. In Section 4 we analyse rank one matrices contributing to the minimal decompositions of tensors, and we distinguish seven types of such matrices. Then we show that to prove the additivity of the tensor rank one can get rid of two of those types, that is, we can produce a smaller example, which does not have these two types, but if the additivity holds for the smaller one, then it also holds for the original one. This is the core observation to prove the main result, Theorem 1.2. Finally, in Section 5 we analyse the additivity of the border rank for small tensor spaces. For most of the possible splittings of the triple $A = \mathbb{C}^4 = A' \oplus A''$, $B = \mathbb{C}^4 = B' \oplus B''$, $C = \mathbb{C}^4 = C' \oplus C''$, there is an easy observation (Corollary 5.2) proving the additivity of the border rank. The remaining two pairs of triples are treated by more advanced methods, involving in particular the Strassen type equations for secant varieties. We conclude the article with a brief discussion of the potential analogue of Theorem 1.4 for $A = B = C = \mathbb{C}^5$.

2. Ranks and slices. This section reviews the notions of rank, border rank, slices, conciseness. Readers that are familiar to these concepts may easily skip this section. The main things to remember from here are Notation 2.2 and Proposition 2.10.

Let $A_1, A_2, \dots, A_d, A, B, C$, and V be finite dimensional vector spaces over a field k . Recall a tensor $s \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ is *simple* if and only if it can be written as $a_1 \otimes a_2 \otimes \dots \otimes a_d$ with $a_i \in A_i$. Simple tensors will also be referred to as *rank one tensors* throughout this paper. If P is a subset of V , we denote by $\langle P \rangle$ its linear span. If $P = \{p_1, \dots, p_r\}$ is a finite subset, we will write $\langle p_1, \dots, p_r \rangle$ rather than $\langle \{p_1, \dots, p_r\} \rangle$ to simplify notation.

DEFINITION 2.1. *Suppose $W \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$ is a linear subspace of the tensor product space. We define $R(W)$, the rank of W , to be the minimal number r , such that there exist simple tensors s_1, \dots, s_r with W contained in $\langle s_1, \dots, s_r \rangle$. For $p \in A_1 \otimes \dots \otimes A_d$, we write $R(p) := R(\langle p \rangle)$.*

In the setting of the definition, if $d = 1$, then $R(W) = \dim W$. If $d = 2$ and $W = \langle p \rangle$ is 1-dimensional, then $R(W)$ is the rank of p viewed as a linear map $A_1^* \rightarrow A_2$. If $d = 3$ and $W = \langle p \rangle$ is 1-dimensional, then $R(W)$ is equal to $R(p)$ in the sense of Section 1. More generally, for arbitrary d , one can relate the rank $R(p)$ of d -way tensors with the rank $R(W)$ of certain linear subspaces in the space of $(d-1)$ -way tensors. This relation is based on the *slice technique*, which we are going to review in Section 2.4.

2.1. Variety of simple tensors. As it is clear from the definition, the rank of a tensor does not depend on the non-zero rescalings of p . Thus it is natural and customary to consider the rank as a function on the projective space $\mathbb{P}(A_1 \otimes A_2 \otimes \dots \otimes A_d)$. There the set of simple tensors is naturally isomorphic to the cartesian product of projective spaces. Its embedding in the tensor space is also called the *Segre variety*:

$$\text{Seg} = \text{Seg}_{A_1, A_2, \dots, A_d} := \mathbb{P}A_1 \times \mathbb{P}A_2 \times \dots \times \mathbb{P}A_d \subset \mathbb{P}(A_1 \otimes A_2 \otimes \dots \otimes A_d).$$

We will intersect linear subspaces of the tensor space with the Segre variety. Using the language of algebraic geometry, such intersection may have a non-trivial scheme structure. In this article we just ignore the scheme structure and all our intersections are set theoretic. To avoid ambiguity of notation, we write $(\cdot)_{\text{red}}$ to underline this issue, while the reader not originating from algebraic geometry should ignore the symbol $(\cdot)_{\text{red}}$.

NOTATION 2.2. *Given a linear subspace of a tensor space, $V \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$, we denote:*

$$V_{\text{Seg}} := (\mathbb{P}V \cap \text{Seg})_{\text{red}}.$$

Thus V_{Seg} is (up to projectivisation) the set of rank one tensors in V .

In this setting, we have the following trivial rephrasing of the definition of rank:

PROPOSITION 2.3. *Suppose $W \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$ is a linear subspace. Then $R(W)$ is equal to the minimal number r , such that there exists a linear subspace $V \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$ of dimension r with $W \subset V$ and $\mathbb{P}V$ is linearly spanned by V_{Seg} . In particular,*

(i) $R(W) = \dim W$ if and only if

$$\mathbb{P}W = \langle W_{\text{Seg}} \rangle.$$

- (ii) Let U be the linear subspace such that $\mathbb{P}U := \langle W_{\text{Seg}} \rangle$. Then $\dim U$ tensors from W can be used in the minimal decomposition of W , that is, there exist $s_1, \dots, s_{\dim U} \in W_{\text{Seg}}$ such that $W \subset \langle s_1, \dots, s_{R(W)} \rangle$ and s_i are simple tensors.

2.2. Secant varieties and border rank. For this subsection (and also in Section 5) we assume $\mathbb{k} = \mathbb{C}$. See Remark 2.6 for generalisations.

In general, the set of tensors of rank at most r is neither open nor closed. One of the very few exceptions is the case of matrices, that is, tensors in $A \otimes B$. Instead, one defines the secant variety $\sigma_r(\text{Seg}_{A_1, \dots, A_d}) \subset \mathbb{P}(A_1 \otimes \dots \otimes A_d)$ as:

$$\sigma_r = \sigma_r(\text{Seg}_{A_1, \dots, A_d}) := \overline{\{p \in \mathbb{P}(A_1 \otimes \dots \otimes A_d) \mid R(p) \leq r\}}.$$

The overline $\overline{\{\cdot\}}$ denotes the closure in the Zariski topology. However in this definition, the resulting set coincides with the Euclidean closure. This is a classically studied algebraic variety [2], [26], [35], and leads to a definition of border rank of a point.

DEFINITION 2.4. For $p \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ define $\underline{R}(p)$, the border rank of p , to be the minimal number r , such that $\langle p \rangle \in \sigma_r(\text{Seg}_{A_1, \dots, A_d})$, where $\langle p \rangle$ is the underlying point of p in the projective space. We follow the standard convention that $\underline{R}(p) = 0$ if and only if $p = 0$.

Analogously we can give the same definitions for linear subspaces. Fix A_1, \dots, A_d and an integer k . Denote by $\text{Gr}(k, A_1 \otimes \dots \otimes A_d)$ the Grassmannian of k -dimensional linear subspaces of the vector space $A_1 \otimes \dots \otimes A_d$. Let $\sigma_{r,k}(\text{Seg}) \subset \text{Gr}(k, A_1 \otimes \dots \otimes A_d)$ be the Grassmann secant variety [9], [15], [16]:

$$\sigma_{r,k}(\text{Seg}) := \overline{\{W \in \text{Gr}(k, A_1 \otimes \dots \otimes A_d) \mid R(W) \leq r\}}.$$

DEFINITION 2.5. For $W \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$, a linear subspace of dimension k , define $\underline{R}(W)$, the border rank of W , to be the minimal number r , such that $W \in \sigma_{r,k}(\text{Seg}_{A_1, \dots, A_d})$.

In particular, if $k = 1$, then Definition 2.5 coincides with Definition 2.4: $\underline{R}(p) = \underline{R}(\langle p \rangle)$. An important consequence of the definitions of border rank of a point or of a linear space is that it is a semicontinuous function

$$\underline{R}: \text{Gr}(k, A_1 \otimes \dots \otimes A_d) \rightarrow \mathbb{N}$$

for every k . Moreover, $\underline{R}(p) = 1$ if and only if $\langle p \rangle \in \text{Seg}$.

REMARK 2.6. When treating the border rank and secant varieties we assume the base field is $\mathbb{k} = \mathbb{C}$. However, the results of [8, §6, Prop. 6.11] imply (roughly) that anything that we can say about a secant variety over \mathbb{C} , we can also say about the same secant variety over any field \mathbb{k} of characteristic 0. In particular, the same results for border rank over an algebraically closed field \mathbb{k} will be true. If \mathbb{k} is not algebraically closed, then the definition of border rank above might not generalise immediately, as there might be a difference between the closure in the Zariski topology or in some other topology, the latter being the Euclidean topology in the case $\mathbb{k} = \mathbb{R}$.

2.3. Independence of the rank of the ambient space. As defined above, the notions of rank and border rank of a vector subspace $W \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$, or of a tensor $p \in A_1 \otimes \dots \otimes A_d$, might seem to depend on the ambient spaces A_i . However, it is well known that the rank is actually independent of the choice of the vector spaces. We first recall this result for tensors, then we apply the slice technique to show it in general.

LEMMA 2.7 ([22, Prop. 3.1.3.1] and [9, Cor. 2.2]). Suppose $\mathbb{k} = \mathbb{C}$ and $p \in A'_1 \otimes A'_2 \otimes \dots \otimes A'_d$ for some linear subspaces $A'_i \subset A_i$. Then $R(p)$ (respectively, $\underline{R}(p)$) measured as the rank (respectively, the border rank) in $A'_1 \otimes \dots \otimes A'_d$ is equal to the rank (respectively, the border rank) measured in $A_1 \otimes \dots \otimes A_d$.

We also state a stronger fact about the rank from the same references: in the notation of Lemma 2.7, any minimal expression $W \subset \langle s_1, \dots, s_{R(W)} \rangle$, for simple tensors s_i , must be contained in $A'_1 \otimes \dots \otimes A'_d$. Here we show that the difference in the length of the decompositions must be at least the difference of the respective dimensions. For simplicity of notation, we restrict the presentation to the case $d = 3$. The reader will easily generalise the argument to any other numbers of factors. We stress that the lemma below does not depend on the base field, in particular, it does not require algebraic closedness.

LEMMA 2.8. Suppose that $p \in A' \otimes B \otimes C$, for a linear subspace $A' \subset A$, and that we have an expression $p \in \langle s_1, \dots, s_r \rangle$, where $s_i = a_i \otimes b_i \otimes c_i$ are simple tensors. Then:

$$r \geq R(p) + \dim \langle a_1, \dots, a_r \rangle - \dim A'.$$

In particular, Lemma 2.8 implies the rank part of Lemma 2.7 for any base field \mathbb{k} , which on its own can also be seen by following the proof of [22, Prop. 3.1.3.1] or [9, Cor. 2.2].

Proof. For simplicity of notation, we assume that $A' \subset \langle a_1, \dots, a_r \rangle$ (by replacing A' with a smaller subspace if needed) and that $A = \langle a_1, \dots, a_r \rangle$ (by replacing A with a smaller subspace). Set $k = \dim A - \dim A'$ and let us reorder the simple tensors s_i in such a way that the first k of the a_i 's are linearly independent and $\langle A' \sqcup \{a_1, \dots, a_k\} \rangle = A$.

Let $A'' = \langle a_1, \dots, a_k \rangle$ so that $A = A' \oplus A''$ and consider the quotient map $\pi: A \rightarrow A/A''$. Then the composition $A' \rightarrow A \xrightarrow{\pi} A/A'' \simeq A'$ is an isomorphism, denoted by ϕ . By a minor abuse of notation, let π and ϕ also denote the induced maps $\pi: A \otimes B \otimes C \rightarrow (A/A'') \otimes B \otimes C$ and $\phi: A' \otimes B \otimes C \simeq A' \otimes B \otimes C$. We have

$$\begin{aligned} \phi(p) &= \pi(p) \in \pi(\langle a_1 \otimes b_1 \otimes c_1, \dots, a_r \otimes b_r \otimes c_r \rangle) \\ &= \langle \pi(a_1) \otimes b_1 \otimes c_1, \dots, \pi(a_r) \otimes b_r \otimes c_r \rangle \\ &= \langle \pi(a_{k+1}) \otimes b_{k+1} \otimes c_{k+1}, \dots, \pi(a_r) \otimes b_r \otimes c_r \rangle. \end{aligned}$$

Using the inverse of the isomorphism ϕ , we get a presentation of p as a linear combination of $(r - k)$ simple tensors, that is, $R(p) \leq r - k$ as claimed.

2.4. Slice technique and conciseness. We define the notion of conciseness of tensors and we review a standard *slice technique* that replaces the calculation of rank of three way tensors with the calculation of rank of linear spaces of matrices.

A tensor $p \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ determines a linear map $p: A_1^* \rightarrow A_2 \otimes \dots \otimes A_d$. Consider the image $W = p(A_1^*) \subset A_2 \otimes \dots \otimes A_d$. The elements of a basis of W (or the image of a basis of A_1^*) are called *slices* of p . The point is that W essentially uniquely (up to an action of $GL(A_1)$) determines p (cfr. [9, Cor. 3.6]). Thus the subspace W captures the geometric information about p , in particular its rank and border rank.

LEMMA 2.9 ([9, Thm 2.5]). *Suppose $p \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ and $W = p(A_1^*)$ as above. Then $R(p) = R(W)$ and (if $\mathbb{k} = \mathbb{C}$) $\underline{R}(p) = \underline{R}(W)$.*

Clearly, we can also replace A_1 with any of the A_i to define slices as images $p(A_i^*)$ and obtain the analogue of the lemma.

We can now prove the analogue of Lemmas 2.7 and 2.8 for higher dimensional subspaces of the tensor space. As before, to simplify the notation, we only consider the case $d = 2$, which is our main interest.

PROPOSITION 2.10. *Suppose $W \subset B' \otimes C'$ for some linear subspaces $B' \subset B, C' \subset C$.*

- (i) *The numbers $R(W)$ and $\underline{R}(W)$ measured as the rank and border rank of W in $B' \otimes C'$ are equal to its rank and border rank calculated in $B \otimes C$ (in the statement about border rank, we assume that $\mathbb{k} = \mathbb{C}$).*
- (ii) *Moreover, if we have an expression $W \subset \langle s_1, \dots, s_r \rangle$, where $s_i = b_i \otimes c_i$ are simple tensors, then:*

$$r \geq R(W) + \dim \langle b_1, \dots, b_r \rangle - \dim B'$$

Proof. Reduce to Lemmas 2.7 and 2.8 using Lemma 2.9.

We conclude this section by recalling the following definition.

DEFINITION 2.11. *Let $p \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ be a tensor or let $W \subset A_1 \otimes A_2 \otimes \dots \otimes A_d$ be a linear subspace. We say that p or W is A_1 -concise if for all linear subspaces $V \subset A_1$, if $p \in V \otimes A_2 \otimes \dots \otimes A_d$ (respectively, $W \subset V \otimes A_2 \otimes \dots \otimes A_d$), then $V = A_1$. Analogously, we define A_i -concise tensors and spaces for $i = 2, \dots, d$. We say p or W is concise if it is A_i -concise for all $i \in \{1, \dots, n\}$.*

REMARK 2.12. *Notice that $p \in A_1 \otimes A_2 \otimes \dots \otimes A_d$ is A_1 -concise if and only if $p: A_1^* \rightarrow A_2 \otimes \dots \otimes A_d$ is injective.*

3. Direct sum tensors and spaces of matrices. Again, for simplicity of notation we restrict the presentation to the case of tensors in $A \otimes B \otimes C$ or linear subspaces of $B \otimes C$.

We introduce the following notation that will be adopted throughout this manuscript.

NOTATION 3.1. *Let $A', A'', B', B'', C', C''$ be vector spaces over \mathbb{k} of dimensions, respectively, $\mathbf{a}', \mathbf{a}'', \mathbf{b}', \mathbf{b}'', \mathbf{c}', \mathbf{c}''$. Suppose $A = A' \oplus A'', B = B' \oplus B'', C = C' \oplus C''$ and $\mathbf{a} = \dim A = \mathbf{a}' + \mathbf{a}''$, $\mathbf{b} = \dim B = \mathbf{b}' + \mathbf{b}''$ and $\mathbf{c} = \dim C = \mathbf{c}' + \mathbf{c}''$.*

For the purpose of illustration, we will interpret the two-way tensors in $B \otimes C$ as matrices in $\mathcal{M}^{\mathbf{b} \times \mathbf{c}}$. This requires choosing bases of B and C , but (whenever possible) we will refrain from naming the bases explicitly. We will refer to an element of the space of matrices $\mathcal{M}^{\mathbf{b} \times \mathbf{c}} \simeq B \otimes C$ as

a $(\mathbf{b}' + \mathbf{b}'', \mathbf{c}' + \mathbf{c}'')$ partitioned matrix. Every matrix $w \in \mathcal{M}^{\mathbf{b} \times \mathbf{c}}$ is a block matrix with four blocks of size $\mathbf{b}' \times \mathbf{c}'$, $\mathbf{b}' \times \mathbf{c}''$, $\mathbf{b}'' \times \mathbf{c}'$ and $\mathbf{b}'' \times \mathbf{c}''$ respectively.

NOTATION 3.2. As in Section 2.4, a tensor $p \in A \otimes B \otimes C$ is a linear map $p : A^* \rightarrow B \otimes C$; we denote by $W := p(A^*)$ the image of A^* in the space of matrices $B \otimes C$. Similarly, if $p = p' + p'' \in (A' \oplus A'') \otimes (B' \oplus B'') \otimes (C' \oplus C'')$ is such that $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$, we set $W' := p'(A'^*) \subset B' \otimes C'$ and $W'' := p''(A''^*) \subset B'' \otimes C''$. In such situation, we will say that $p = p' \oplus p''$ is a direct sum tensor.

We have the following direct sum decomposition:

$$W = W' \oplus W'' \subset (B' \otimes C') \oplus (B'' \otimes C'')$$

and an induced matrix partition of type $(\mathbf{b}' + \mathbf{b}'', \mathbf{c}' + \mathbf{c}'')$ on every matrix $w \in W$ such that

$$w = \begin{pmatrix} w' & \underline{0} \\ \underline{0} & w'' \end{pmatrix},$$

where $w' \in W'$ and $w'' \in W''$, and the two $\underline{0}$'s denote zero matrices of size $\mathbf{b}' \times \mathbf{c}''$ and $\mathbf{b}'' \times \mathbf{c}'$ respectively.

PROPOSITION 3.3. Suppose that p, W , etc. are as in Notation 3.2. Then the additivity of the rank holds for p , that is $R(p) = R(p') + R(p'')$, if and only if the additivity of the rank holds for W , that is, $R(W) = R(W') + R(W'')$.

Proof. It is an immediate consequence of Lemma 2.9.

3.1. Projections and decompositions. The situation we consider here again concerns the direct sums and their minimal decompositions. We fix $W' \subset B' \otimes C'$ and $W'' \subset B'' \otimes C''$ and we choose a minimal decomposition of $W' \oplus W''$, that is, a linear subspace $V \subset B \otimes C$ such that $\dim V = R(W' \oplus W'')$, $\mathbb{P}V = \langle V_{\text{Seg}} \rangle$ and $V \supset W' \oplus W''$. Such linear spaces W', W'' and V will be fixed for the rest of Sections 3 and 4.

In addition to Notations 2.2, 3.1 and 3.2 we need the following.

NOTATION 3.4. Under Notation 3.1, let $\pi_{C'}$ denote the projection

$$\pi_{C'} : C \rightarrow C'', \text{ or}$$

whose kernel is the space C' . With slight abuse of notation, we shall denote by $\pi_{C'}$ also the following projections

$$\pi_{C'} : B \otimes C \rightarrow B \otimes C'', \text{ or } \pi_{C'} : A \otimes B \otimes C \rightarrow A \otimes B \otimes C'',$$

with kernels, respectively, $B \otimes C'$ and $A \otimes B \otimes C'$. The target of the projection is regarded as a subspace of $C, B \otimes C$, or $A \otimes B \otimes C$, so that it is possible to compose such projections, for instance:

$$\pi_{C'} \pi_{B''} : B \otimes C \rightarrow B' \otimes C'', \text{ or } \pi_{C'} \pi_{B''} : A \otimes B \otimes C \rightarrow A \otimes B' \otimes C''.$$

We also let $E' \subset B'$ (resp. $E'' \subset B''$) be the minimal vector subspace such that $\pi_{C'}(V)$ (resp. $\pi_{C''}(V)$) is contained in $(E' \oplus B'') \otimes C''$ (resp. $(B' \oplus E'') \otimes C'$).

By swapping the roles of B and C , we define $F' \subset C'$ and $F'' \subset C''$ analogously. By the lowercase letters $\mathbf{e}', \mathbf{e}'', \mathbf{f}', \mathbf{f}''$ we denote the dimensions of the subspaces E', E'', F', F'' .

If the differences $R(W') - \dim W'$ and $R(W'') - \dim W''$ (which we will informally call the *gaps*) are large, then the spaces E', E'', F', F'' could be large too, in particular they can coincide with B', B'', C', C'' respectively. In fact, these spaces measure “how far” a minimal decomposition V of a direct sum $W = W' \oplus W''$ is from being a direct sum of decompositions of W' and W'' .

In particular, we will show in Proposition 4.5 and Corollary 4.16, that if $E'' = \{0\}$ or if both E'' and F'' are sufficiently small, then $R(W) = R(W') + R(W'')$. Then, as a consequence of Corollary 3.7, if one of the gaps is at most two (say, $R(W'') = \dim W'' + 2$), then the additivity of the rank holds, see Theorem 4.17.

LEMMA 3.5. In Notation 3.4 as above, with $W = W' \oplus W'' \subset B \otimes C$, the following inequalities hold.

$$\begin{aligned} R(W') + \mathbf{e}' &\leq R(W) - \dim W'', & R(W'') + \mathbf{e}' &\leq R(W) - \dim W', \\ R(W') + \mathbf{f}'' &\leq R(W) - \dim W'', & R(W'') + \mathbf{f}'' &\leq R(W) - \dim W'. \end{aligned}$$

Proof. We prove only the first inequality $R(W') + \mathbf{e}' \leq R(W) - \dim W''$, the other follow in the same way by swapping B and C or $'$ and $''$. By Proposition 2.10(i) and (ii) we may assume W' is concise: $R(W')$ or $R(W)$ are not affected by choosing the minimal subspace of B' by (i), also the minimal decomposition V cannot involve anyone from outside of the minimal subspace by (ii).

Since V is spanned by rank one matrices and the projection $\pi_{C''}$ preserves the set of matrices of rank at most one, also the vector space $\pi_{C''}(V)$ is spanned by rank one matrices, say

$$\pi_{C''}(V) = \langle b_1 \otimes c_1, \dots, b_r \otimes c_r \rangle$$

with $r = \dim \pi_{C''}(V)$. Moreover, $\pi_{C''}(V)$ contains W' . We claim that

$$B' \oplus E'' = \langle b_1, \dots, b_r \rangle.$$

Indeed, the inclusion $B' \subset \langle b_1, \dots, b_r \rangle$ follows from the conciseness of W' , as $W' \subset V \cap B' \otimes C'$. Moreover, the inclusions $E'' \subset \langle b_1, \dots, b_r \rangle$ and $B' \oplus E'' \supset \langle b_1, \dots, b_r \rangle$ follow from the definition of E'' , cf. Notation 3.4.

Thus Proposition 2.10(ii) implies that

$$(3.6) \quad r = \dim \pi_{C''}(V) \geq R(W') + \underbrace{\dim \langle b_1, \dots, b_r \rangle}_{\mathbf{b}' + \mathbf{e}''} - \mathbf{b}' = R(W') + \mathbf{e}''.$$

Since V contains W'' and $\pi_{C''}(W'') = \{0\}$, we have

$$r = \dim \pi_{C''}(V) \leq \dim V - \dim W'' = R(W) - \dim W''.$$

The claim follows from the above inequality together with (3.6). \square

Rephrasing the inequalities of Lemma 3.5, we obtain the following.

COROLLARY 3.7. *If $R(W) < R(W') + R(W'')$, then*

$$\begin{aligned} \mathbf{e}' &< R(W') - \dim W', & \mathbf{f}' &< R(W') - \dim W', \\ \mathbf{e}'' &< R(W'') - \dim W'', & \mathbf{f}'' &< R(W'') - \dim W''. \end{aligned}$$

This immediately recovers the known case of additivity, when the gap is equal to 0, that is, if $R(W') = \dim W'$, then $R(W) = R(W') + R(W'')$ (because $\mathbf{e}' \geq 0$). Moreover, it implies that if one of the gaps is equal to 1 (say $R(W') = \dim W' + 1$), then either the additivity holds or both E' and F' are trivial vector spaces. In fact, the latter case is only possible if the former case holds too.

LEMMA 3.8. *With Notation 3.4, suppose $E' = \{0\}$ and $F' = \{0\}$. Then the additivity of the rank holds $R(W) = R(W') + R(W'')$. In particular, if $R(W') \leq \dim W' + 1$, then the additivity holds.*

Proof. Since $E' = \{0\}$ and $F' = \{0\}$, by the definition of E' and F' we must have the following inclusions:

$$\pi_{B''}(V) \subset B' \otimes C' \text{ and } \pi_{C''}(V) \subset B' \otimes C'.$$

Therefore $V \subset B' \otimes C' \oplus B'' \otimes C''$ and V is obtained from the union of the decompositions of W' and W'' .

The last statement follows from Corollary 3.7 \square

Later in Proposition 4.5 we will show a stronger version of the above lemma, namely that it is sufficient to assume that only one of E' or F' is zero. In Corollary 4.16 we prove a further generalisation based on the results in the following subsection.

3.2. “Hook”-shaped spaces. It is known since [21] that the additivity of the tensor rank holds for tensors with one of the factors of dimension 2, that is, using Notation 3.1 and 3.2, if $\mathbf{a}' \leq 2$ then $R(p' + p'') = R(p') + R(p'')$. The same claim is recalled in [24, Sect. 4] after Theorem 4.1. The brief comment says that if rank of p' can be calculated by the *substitution method*, then the additivity of the rank holds. Landsberg and Michałek implicitly suggest that if $\mathbf{a}' \leq 2$, then the rank of p' can be calculated by the substitution method, [24, Items (1)–(6) after Prop. 3.1]. This is indeed the case (at least over an algebraically closed field \mathbb{k}), although rather demanding to verify, at least in the version of the algorithm presented in the cited article. In particular, to show that the substitution method can calculate the rank of $p' \in \mathbb{k}^2 \otimes B' \otimes C'$, one needs to use the normal forms of such tensors [22, §10.3] and understand all the cases, and it is hard to agree that this method is so much simpler than the original approach of [21].

Instead, probably, the intention of the authors of [24] was slightly different, with a more direct application of [24, Prop. 3.1] (or Proposition 3.11 below). This has been carefully detailed and described in [27, Prop. 3.2.12] and here we present this approach to show a stronger statement about small “hook”-shaped spaces (Proposition 3.18). We stress that our argument for Proposition 3.18, as well as [27, Prop. 3.2.12] requires the assumption of an algebraically closed base field \mathbb{k} , while the original approach of [21] works over any field. For a short while we also work over an arbitrary field.

DEFINITION 3.9. *For non-negative integers e, f , we say that a linear subspace $W \subset B \otimes C$ is (e, f) -hook shaped, if $W \subset \mathbb{k}^e \otimes C + B \otimes \mathbb{k}^f$ for some choices of linear subspaces $\mathbb{k}^e \subset B$ and $\mathbb{k}^f \subset C$.*

The name “hook shaped” space comes from the fact that under an appropriate choice of basis, the only non-zero coordinates form a shape of a hook \lrcorner situated in the upper left corner of the matrix, see Example 3.10. The integers (e, f) specify how wide the edges of the hook are. A similar name also appears in the context of Young diagrams, see for instance [5, Def. 2.3].

EXAMPLE 3.10. *A $(1, 2)$ -hook shaped subspace of $\mathbb{k}^4 \otimes \mathbb{k}^4$ has only the following possibly nonzero entries in some coordinates:*

$$\begin{bmatrix} * & * & * & * \\ * & * & 0 & 0 \\ * & * & 0 & 0 \\ * & * & 0 & 0 \end{bmatrix}.$$

The following elementary observation is presented in [24, Prop. 3.1] and in [3, Lem. B.1]. Here we have phrased it in a coordinate free way.

PROPOSITION 3.11. *Let $p \in A \otimes B \otimes C$, $R(p) = r > 0$, and pick $\alpha \in A^*$ such that $p(\alpha) \in B \otimes C$ is nonzero. Consider two hyperplanes in A : the linear hyperplane $\alpha^\perp = (\alpha = 0)$ and the affine hyperplane $(\alpha = 1)$. For any $a \in (\alpha = 1)$, denote*

$$\tilde{p}_a := p - a \otimes p(\alpha) \in \alpha^\perp \otimes B \otimes C.$$

Then:

- (i) *there exists a choice of $a \in (\alpha = 1)$ such that $R(\tilde{p}_a) \leq r - 1$,*
- (ii) *if in addition $R(p(\alpha)) = 1$, then for any choice of $a \in (\alpha = 1)$ we have $R(\tilde{p}_a) \geq r - 1$.*

See [24, Prop. 3.1] for the proof (note the statement there is over the complex numbers only, but the proof is field independent) or, alternatively, using Lemma 2.9 translate it into the following straightforward statement on linear spaces of tensors:

PROPOSITION 3.12. *Suppose $W \subset B \otimes C$ is a linear subspace, $R(W) = r$. Assume $w \in W$ is a non-zero element. Then:*

- (i) *there exists a choice of a complementary subspace $\tilde{W} \subset W$, such that $\tilde{W} \oplus \langle w \rangle = W$ and $R(\tilde{W}) \leq r - 1$, and*
- (ii) *if in addition $R(w) = 1$, then for any choice of the complementary subspace $\tilde{W} \oplus \langle w \rangle = W$ we have $R(\tilde{W}) \geq r - 1$.*

Proposition 3.11 is crucial in the proof that the additivity of the rank holds for vector spaces, one of which is $(1, 2)$ -hook shaped (provided that the base field is algebraically closed). Before taking care of that, we use the same proposition to prove a simpler statement about $(1, 1)$ -hook shaped spaces, which is valid without any assumption on the field. The proof essentially follows the idea outlined in [24, Thm 4.1].

PROPOSITION 3.13. *Suppose $W'' \subset B'' \otimes C''$ is $(1, 1)$ -hook shaped and $W' \subset B' \otimes C'$ is an arbitrary subspace. Then the additivity of the rank holds for $W' \oplus W''$.*

Before commencing the proof of the proposition we state three lemmas, which will be applied to both $(1, 1)$ and $(1, 2)$ hook shaped spaces. The first lemma is analogous to [24, Thm 4.1]. In this lemma (and also in the rest of this section) we will work with a sequence of tensors, p_0, p_1, p_2, \dots in the space $A \otimes B \otimes C$, which are not necessarily direct sums. Nevertheless, for each i , we write $p'_i = \pi_{A''} \pi_{B''} \pi_{C''}(p_i)$ (that is, this is the “corner” of p_i corresponding to A' , B' and C'). We define p''_i analogously.

LEMMA 3.14. *Suppose $W' \subset A' \otimes B' \otimes C'$ and $W'' \subset A'' \otimes B'' \otimes C''$ are two subspaces. Let $r'' = R(W'')$ and suppose that there exists a sequence of tensors $p_0, p_1, p_2, \dots, p_{r''} \in A \otimes B \otimes C$ satisfying the following properties:*

- (1) $p_0 = p$ is such that $p(A^*) = W = W' \oplus W''$,
- (2) $p'_{i+1} = p'_i$ for every $0 \leq i < r''$,
- (3) $R(p''_{i+1}) \geq R(p''_i) - 1$ for every $0 \leq i < r''$,
- (4) $R(p_{i+1}) \leq R(p_i) - 1$ for each $0 \leq i < r''$.

Then the additivity of the rank holds for $W' \oplus W''$ and for each $i < r''$ we must have $p''_i \neq 0$.

Proof. We have

$$R(W') + R(W'') \stackrel{(1),(2)}{=} R(p'_{r''}) + r'' \leq R(p_{r''}) + r'' \stackrel{(4)}{\leq} R(p_0) \stackrel{(1)}{=} R(W).$$

The nonvanishing of p''_i follows from (3). \square

The second lemma tells us how to construct a single step in the above sequence.

LEMMA 3.15. *Suppose $\Sigma \subset A \otimes B \otimes C$ is a linear subspace, $p_i \in \Sigma$ is a tensor, and $\gamma \in C''$ is such that:*

- $R(p_i''(\gamma)) = 1$,
- γ preserves Σ , that is, $\Sigma(\gamma) \otimes C \subset \Sigma$, where $\Sigma(\gamma) = \{t(\gamma) \mid t \in \Sigma\} \subset A \otimes B$.
- $\Sigma(\gamma)$ does not have entries in $A' \otimes B'$, that is $\pi_{A'} \pi_{B'}(\Sigma(\gamma)) = 0$.

Consider $\gamma^\perp \subset C$ to be the perpendicular hyperplane. Then there exists $p_{i+1} \in (\Sigma \cap A \otimes B \otimes \gamma^\perp)$ that satisfies properties (2)–(4) of Lemma 3.14 (for a fixed i).

Proof. As in Proposition 3.11 for $c \in (\gamma = 1)$ set $(\tilde{p}_i)_c = p_i - p_i(\gamma) \otimes c \in A \otimes B \otimes \gamma^\perp$. We will pick p_{i+1} among the $(\tilde{p}_i)_c$. In fact by Proposition 3.11(i) there exists a choice of c such that $p_{i+1} = (\tilde{p}_i)_c$ has rank less than $R(p_i)$, that is, (4) is satisfied. On the other hand, since γ is in $(C'')^*$, we have $p_{i+1}'' = \left(\tilde{p}_i''\right)_{c''}$ (where $c = c' + c''$ with $c' \in C'$ and $c'' \in C''$) and by Proposition 3.11(ii) also (3) is satisfied. Property (2) follows, as $\Sigma(\gamma)$ (in particular, $p_i(\gamma)$) has no entries in $A' \otimes B' \otimes C'$. Finally, $p_{i+1} \in \Sigma$ thanks to the assumption that γ preserves Σ and Σ is a linear subspace. \square

The next lemma is the common first step in the proofs of additivity for (1, 1) and (1, 2) hooks: we construct a few initial elements of the sequence needed in Lemma 3.14.

LEMMA 3.16. *Suppose $W'' \subset B'' \otimes C''$ is a $(1, f)$ -hook shaped space for some integer f and $W' \subset B' \otimes C'$ is arbitrary. Fix $\mathbb{k}^1 \subset B''$ and $\mathbb{k}^f \subset C''$ as in Definition 3.9 for W'' . Then there exists a sequence of tensors $p_0, p_1, p_2, \dots, p_k \in A \otimes B \otimes C$ for some k that satisfies properties (1)–(4) of Lemma 3.14 and in addition $p_k'' \in A'' \otimes B'' \otimes \mathbb{k}^f$ and for every i we have $p_i \in A' \otimes B' \otimes C' \oplus A'' \otimes (B'' \otimes \mathbb{k}^f + \mathbb{k}^1 \otimes C)$. In particular:*

- $p_i''((A'')^*)$ is a $(1, f)$ -hook shaped space for every $i < k$, while $p_k''((A'')^*)$ is a $(0, f)$ -hook shaped space.
- Every p_i is “almost” a direct sum tensor, that is, $p_i = (p_i' \oplus p_i'') + q_i$, where

$$q_i \in A'' \otimes \mathbb{k}^1 \otimes C' \subset A'' \otimes B'' \otimes C'.$$

Proof. To construct the sequence p_i we recursively apply Lemma 3.15. By our assumptions, $p'' \in A'' \otimes B'' \otimes \mathbb{k}^f + A'' \otimes \langle x \rangle \otimes C''$ for some choice of $x \in B''$ and fixed $\mathbb{k}^f \subset C''$. We let $\Sigma = A' \otimes B' \otimes C' \oplus A'' \otimes (B'' \otimes \mathbb{k}^f + \langle x \rangle \otimes C)$.

Tensor p_0 is defined by (1). Suppose we have already constructed p_0, \dots, p_i and that p_i'' is not yet contained in $A'' \otimes B'' \otimes \mathbb{k}^f$. Therefore there exists a hyperplane $\gamma^\perp = (\gamma = 0) \subset C'$ for some $\gamma \in (C'')^* \subset C^*$ such that $\mathbb{k}^f \subset \gamma^\perp$, but $p_i'' \notin A'' \otimes B'' \otimes \gamma^\perp$. Equivalently, $p_i''(\gamma) \neq 0$ and $p_i''(\gamma) \in A'' \otimes \langle x \rangle$. In particular, $R(p_i''(\gamma)) = 1$ and $\Sigma(\gamma) \subset A'' \otimes \langle x \rangle$. Thus γ preserves Σ as in Lemma 3.15 and $\Sigma(\gamma)$ has no entries in $A' \otimes B' \otimes C'$.

Thus we construct p_{i+1} using Lemma 3.15. Since we are gradually reducing the dimension of the third factor of the tensor space containing p_{i+1}'' , eventually we will arrive at the case $p_{i+1}'' \in A'' \otimes B'' \otimes \mathbb{k}^f$, proving the claim. \square

Proof of Proposition 3.13. We construct the sequence p_i as in Lemma 3.14. The initial elements p_0, \dots, p_k of the sequence are given by Lemma 3.16. By the lemma and our assumptions, $p_i'' \in A'' \otimes B'' \otimes \langle y \rangle + A'' \otimes \langle x \rangle \otimes C''$ for some choices of $x \in B''$ and $y \in C''$ and

$$p_k \in A' \otimes B' \otimes C' \oplus A'' \otimes (\langle x \rangle \otimes C' \oplus B'' \otimes \langle y \rangle).$$

Now suppose that we have constructed p_k, \dots, p_j for some $j \geq k$ satisfying (2)–(4), such that

$$p_j \in \Sigma = A' \otimes B' \otimes C' \oplus A'' \otimes B \otimes (C' \oplus \langle y \rangle).$$

If $p_j'' = 0$, then by Lemma 3.14 we are done, as $j = r''$. So suppose $p_j'' \neq 0$, and choose $\beta \in (B'')^*$ such that $p_j''(\beta) \neq 0$, that is, $R(p_j''(\beta)) = 1$ since $p_j''(\beta) \in A'' \otimes \langle y \rangle$. We produce p_{j+1} using Lemma 3.15 with the roles of B and C swapped (so also β takes the role of γ etc.).

We stop after constructing $p_{r''}$ and thus the desired sequence exists and proves the claim. \square

In the rest of this section we will show that an analogous statement holds for (1, 2)-hook shaped spaces under an additional assumption that the base field is algebraically closed. We need the following lemma (false for nonclosed fields), whose proof is a straightforward dimension count, see also [27, Prop. 3.2.11].

LEMMA 3.17. *Suppose \mathbb{k} is algebraically closed (of any characteristic) and $p \in A \otimes B \otimes \mathbb{k}^2$ and $p \neq 0$. Then at least one of the following holds:*

- there exists a rank one matrix in $p(A^*) \subset B \otimes \mathbb{k}^2$, or
- for any $x \in B$ there exists a rank one matrix in $p(x^\perp) \subset A \otimes \mathbb{k}^2$, where $x^\perp \subset B^*$ is the hyperplane defined by x .

Proof. If p is not \mathbb{k}^2 -concise, then both claims trivially hold (except if rank of p is one, then only the first claim holds). Thus without loss of generality, we may suppose p is concise by replacing A and B with smaller spaces if necessary. If $\dim A \geq \dim B$, then the projectivisation of the image

$\mathbb{P}(p(A^*)) \subset \mathbb{P}(B \otimes \mathbb{k}^2)$ intersects the Segre variety $\mathbb{P}(B) \times \mathbb{P}^1$ by the dimension count [20, Thm I.7.2] (note that here we use that the base field \mathbb{k} is algebraically closed). Otherwise, $\dim A < \dim B$ and the intersection

$$\mathbb{P}(p(B^*)) \cap (\mathbb{P}(A) \times \mathbb{P}^1) \subset \mathbb{P}(A \otimes \mathbb{k}^2)$$

has positive dimension by the same dimension count. In particular, any hyperplane $\mathbb{P}(p(x^\perp)) \subset \mathbb{P}(p(B^*))$ also intersects the Segre variety. \square

The next proposition reproves (under the additional assumption that \mathbb{k} is algebraically closed) and slightly strengthens the theorem of JaJa-Takche [21], which can be thought of as a theorem about $(0, 2)$ -hook shaped spaces.

PROPOSITION 3.18. *Suppose \mathbb{k} is algebraically closed, $W'' \subset B'' \otimes C''$ is $(1, 2)$ -hook shaped and $W' \subset B' \otimes C'$ is an arbitrary subspace. Then the additivity of the rank holds for $W' \oplus W''$.*

Proof. We will use Lemmas 3.14, 3.15 and 3.16 again. That is, we are looking for a sequence $p_0, \dots, p_{r''} \in A \otimes B \otimes C$ with the properties (1)–(4), and the initial elements p_0, \dots, p_k are constructed in such a way that $p_k \in A' \otimes B' \otimes C' \oplus A'' \otimes (\langle x \rangle \otimes C' \oplus B'' \otimes \mathbb{k}^2)$. Here $x \in B''$ is such that $W'' \subset \langle x \rangle \otimes C'' + B'' \otimes \mathbb{k}^2$.

We have already “cleaned” the part of the hook of size 1, and now we work with the remaining space of $\mathbf{b}'' \times 2$ matrices. Unfortunately, cleaning p_i'' produces rubbish in the other parts of the tensor, and we have to control the rubbish so that it does not affect p_i' , see (2). Note that what is left to do is not just the plain case of Strassen’s additivity in the case of $\mathbf{c}'' = 2$ proven in [21] since p_k may have already nontrivial entries in another block, the one corresponding to $A'' \otimes B'' \otimes C'$ (the small tensor q_k in the statement of Lemma 3.16).

We set $\Sigma = A' \otimes B' \otimes C' \oplus A \otimes (B \otimes \mathbb{k}^2 \oplus \langle x \rangle \otimes C')$. To construct p_{j+1} we use Lemma 3.17 (in particular, here we exploit the algebraic closedness of \mathbb{k}). Thus either there exists $\alpha \in (A'')^*$ such that $R(p_j''(\alpha)) = 1$, or there exists $\beta \in x^\perp \subset (B'')^*$ such that $R(p_j''(\beta)) = 1$. In both cases we apply Lemma 3.15 with the roles of A and C swapped or the roles of B and C swapped. The conditions in the lemma are straightforward to verify.

We stop after constructing $p_{r''}$ and thus the desired sequence exists and proves the claim. \square

4. Rank one matrices and additivity of the tensor rank. As hinted by the proof of Proposition 3.18, as long as we have a rank one matrix in the linear space W' or W'' , we have a good starting point for an attempt to prove the additivity of the rank. Throughout this section we will make a formal statement out of this observation and prove that if there is a rank one matrix in the linear spaces, then either the additivity holds or there exists a “smaller” example of failure of the additivity. In Section 4.4 we exploit several versions of this claim in order to prove Theorem 1.2.

Throughout this section we follow Notations 2.2 (denoting the rank one elements in a vector space by the subscript \cdot_{Seg}), 3.1 (introducing the vector spaces A, \dots, C'' and their dimensions $\mathbf{a}, \dots, \mathbf{c}''$), 3.2 (defining a direct sum tensor $p = p' \oplus p''$ and the corresponding vector spaces W, W', W''), and also 3.4 (which explains the conventions for projections $\pi_{A'}, \dots, \pi_{C''}$ and vector spaces E', \dots, F'' , which measure how much the fixed decomposition V of W sticks out from the direct sum $B' \otimes C' \oplus B'' \otimes C''$).

4.1. Combinatorial splitting of the decomposition. We carefully analyse the structure of the rank one matrices in V . We will distinguish seven types of such matrices.

LEMMA 4.1. *Every element of $V_{Seg} \subset \mathbb{P}(B \otimes C)$ lies in the projectivisation of one of the following subspaces of $B \otimes C$:*

- (i) $B' \otimes C', B'' \otimes C'',$ (Prime, Bis)
- (ii) $E' \otimes (C' \oplus F''), E'' \otimes (F' \oplus C''),$ (HL, HR)
- $(B' \oplus E'') \otimes F', (E' \oplus B'') \otimes F'',$ (VL, VR)
- (iii) $(E' \oplus E'') \otimes (F' \oplus F'').$ (Mix)

The spaces in (i) are purely contained in the original direct summands, hence, in some sense, they are the easiest to deal with (we will show how to “get rid” of them and construct a smaller example justifying a potential lack of additivity).¹ The spaces in (ii) stick out of the original summand, but only in one direction, either horizontal (HL, HR), or vertical (VL, VR)². The space in (iii) is mixed and it sticks out in all directions. It is the most difficult to deal with and we expect that the typical counterexamples to the additivity of the rank will have mostly (or only) such mixed matrices in their minimal decomposition. The mutual configuration and layout of those spaces in the case $\mathbf{b}', \mathbf{b}'', \mathbf{c}', \mathbf{c}'' = 3, \mathbf{e}', \mathbf{e}'', \mathbf{f}', \mathbf{f}'' = 1$ is illustrated in Figure 4.1.

¹The word Bis comes from the Polish way of pronouncing the “” symbol.

²Here the letters “H, V, L, R” stand for “horizontal, vertical, left, right” respectively.




FIGURE 4.1. We use Notation 3.4. In the case $b', b'', c', c'' = 3$, $e', e'', f', f'' = 1$ choose a basis of E' and a completion to a basis of B' and, similarly, bases for (E'', B'') , (F', C') , (F'', C'') . We can represent the elements of $V_{\text{Seg}} \subset B \otimes C$ as matrices in one of the following subspaces: Prime (corresponding to the top-left green rectangle), Bis (bottom-right blue rectangle), VL (purple with entries $v_{1,3}, \dots, v_{4,3}$), VR (purple with entries $v_{3,4}, \dots, v_{6,4}$), HL (brown with entries $v_{3,1}, \dots, v_{3,4}$), HR (brown with entries $v_{4,3}, \dots, v_{4,6}$), and Mix (middle orange square with entries $v_{3,3}, \dots, v_{4,4}$).

Proof of Lemma 4.1. Let $b \otimes c \in V_{\text{Seg}}$ be a matrix of rank one. Write $b = b' + b''$ and $c = c' + c''$, where $b' \in B', b'' \in B'', c' \in C'$ and $c'' \in C''$. We consider the image of $b \otimes c$ via the four natural projections introduced in Notation 3.4:

$$(4.2a) \quad \pi_{B'}(b \otimes c) = b'' \otimes c \in B'' \otimes (F' \oplus C''),$$

$$(4.2b) \quad \pi_{B''}(b \otimes c) = b' \otimes c \in B' \otimes (C' \oplus F''),$$

$$(4.2c) \quad \pi_{C'}(b \otimes c) = b \otimes c'' \in (E' \oplus B'') \otimes C'', \text{ and}$$

$$(4.2d) \quad \pi_{C''}(b \otimes c) = b \otimes c' \in (B' \oplus E'') \otimes C'.$$

Notice that b' and b'' cannot be simultaneously zero, since $b \neq 0$. Analogously, $(c', c'') \neq (0, 0)$.

Equations (4.2a)–(4.2d) prove that the non-vanishing of one of b', b'', c', c'' induces a restriction on another one. For instance, if $b' \neq 0$, then by (4.2b) we must have $c'' \in F''$. Or, if $b'' \neq 0$, then (4.2a) forces $c' \in F'$, and so on. Altogether we obtain the following cases:

- (1) If $b', b'', c', c'' \neq 0$, then $b \otimes c \in (E' \oplus E'') \otimes (F' \oplus F'')$ (case Mix).
- (2) if $b', b'' \neq 0$ and $c' = 0$, then $b \otimes c = b \otimes c'' \in (E' \oplus B'') \otimes F''$ (case VR).
- (3) if $b', b'' \neq 0$ and $c'' = 0$, then $b \otimes c = b \otimes c' \in (B' \oplus E'') \otimes F'$ (case VL).
- (4) If $b' = 0$, then either $c' = 0$ and therefore $b \otimes c = b'' \otimes c'' \in B'' \otimes C''$ (case Bis), or $c' \neq 0$ and $b \otimes c = b'' \otimes c \in E'' \otimes (F' \oplus C'')$ (case HR).
- (5) If $b'' = 0$, then either $c'' = 0$ and thus $b \otimes c = b' \otimes c' \in B' \otimes C'$ (case Prime), or $c'' \neq 0$ and $b \otimes c = b' \otimes c \in E' \otimes (C' \oplus F'')$ (case HL).

This concludes the proof. \square

As in Lemma 4.1 every element of $V_{\text{Seg}} \subset \mathbb{P}(B \otimes C)$ lies in one of seven subspaces of $B \otimes C$. These subspaces may have nonempty intersection. We will now explain our convention with respect to choosing a basis of V consisting of elements of V_{Seg} .

Here and throughout the article by \sqcup we denote the disjoint union.

NOTATION 4.3. We choose a basis \mathcal{B} of V in such a way that:

- \mathcal{B} consist of rank one matrices only,
- $\mathcal{B} = \text{Prime} \sqcup \text{Bis} \sqcup \text{HL} \sqcup \text{HR} \sqcup \text{VL} \sqcup \text{VR} \sqcup \text{Mix}$, where each of Prime, Bis, HL, HR, VL, VR, and Mix is a finite set of rank one matrices of the respective type as in Lemma 4.1 (for instance, $\text{Prime} \subset B' \otimes C'$, $\text{HL} \subset E' \otimes (C' \oplus F'')$, etc.).
- \mathcal{B} has as many elements of Prime and Bis as possible, subject to the first two conditions,

- \mathcal{B} has as many elements of HL, HR, VL and VR as possible, subject to all of the above conditions.

Let **prime** be the number of elements of Prime (equivalently, **prime** = dim (Prime)) and analogously define **bis**, **hl**, **hr**, **vl**, **vr**, and **mix**. The choice of \mathcal{B} need not be unique, but we fix one for the rest of the article. Instead, the numbers **prime**, **bis**, and **mix** are uniquely determined by V (there may be some non-uniqueness in dividing between **hl**, **hr**, **vl**, **vr**).

Thus to each decomposition we associated a sequence of seven non-negative integers (**prime**, ..., **mix**). We now study the inequalities between these integers and exploit them to get theorems about the additivity of the rank.

PROPOSITION 4.4. *In Notations 3.4 and 4.3 the following inequalities hold:*

- (i) **prime** + **hl** + **vl** + min(**mix**, $\mathbf{e}'\mathbf{f}'$) $\geq R(W')$,
- (ii) **bis** + **hr** + **vr** + min(**mix**, $\mathbf{e}''\mathbf{f}''$) $\geq R(W'')$,
- (iii) **prime** + **hl** + **vl** + min(**hr** + **mix**, $\mathbf{f}'(\mathbf{e}' + \mathbf{e}'')$) $\geq R(W') + \mathbf{e}''$,
- (iv) **prime** + **hl** + **vl** + min(**vr** + **mix**, $\mathbf{e}'(\mathbf{f}' + \mathbf{f}'')$) $\geq R(W') + \mathbf{f}''$,
- (v) **bis** + **hr** + **vr** + min(**hl** + **mix**, $\mathbf{f}''(\mathbf{e}' + \mathbf{e}'')$) $\geq R(W'') + \mathbf{e}'$,
- (vi) **bis** + **hr** + **vr** + min(**vl** + **mix**, $\mathbf{e}''(\mathbf{f}' + \mathbf{f}'')$) $\geq R(W'') + \mathbf{f}'$.

Proof. To prove Inequality (i) we consider the composition of projections $\pi_{B''}\pi_{C''}$. The linear space $\pi_{B''}\pi_{C''}(V)$ is spanned by rank one matrices $\pi_{B''}\pi_{C''}(\mathcal{B})$ (where $\mathcal{B} = \text{Prime} \sqcup \dots \sqcup \text{Mix}$ as in Notation 4.3), and it contains W' . Thus $\dim(\pi_{B''}\pi_{C''}(V)) \geq R(W')$. But the only elements of the basis \mathcal{B} that survive both projections (that is, they are not mapped to zero under the composition) are Prime, HL, VL, and Mix. Thus

$$\mathbf{prime} + \mathbf{hl} + \mathbf{vl} + \mathbf{mix} \geq \dim(\pi_{B''}\pi_{C''}(V)) \geq R(W').$$

On the other hand, $\pi_{B''}\pi_{C''}(\text{Mix}) \subset E' \otimes F'$, thus among $\pi_{B''}\pi_{C''}(\text{Mix})$ we can choose at most $\mathbf{e}'\mathbf{f}'$ linearly independent matrices. Thus

$$\mathbf{prime} + \mathbf{hl} + \mathbf{vl} + \mathbf{e}'\mathbf{f}' \geq \dim(\pi_{B''}\pi_{C''}(V)) \geq R(W').$$

The two inequalities prove (i).

To show Inequality (iii) we may assume that W' is concise as in the proof of Lemma 3.5. Moreover, as in that same proof (more precisely, Inequality (3.6)) we show that $\dim \pi_{C''}(V) \geq R(W') + \mathbf{e}''$. But $\pi_{C''}$ sends all matrices from Bis and VR to zero, thus

$$\mathbf{prime} + \mathbf{hl} + \mathbf{vl} + \mathbf{hr} + \mathbf{mix} \geq \dim \pi_{C''}(V) \geq R(W') + \mathbf{e}''.$$

As in the proof of Part (i), we can also replace **hr** + **mix** by $\mathbf{f}'(\mathbf{e}' + \mathbf{e}'')$, since $\pi_{C''}(\text{HR} \cup \text{Mix}) \subset (E' \oplus E'') \otimes F'$, concluding the proof of (iii).

The proofs of the remaining four inequalities are identical to one of the above, after swapping the roles of B and C or $'$ and $''$ (or swapping both pairs). \square

PROPOSITION 4.5. *With Notation 3.4, if one among E', E'', F', F'' is zero, then $R(W) = R(W') + R(W'')$.*

Proof. Let us assume without loss of generality that $E' = \{0\}$. Using the definitions of sets Prime, Bis, VR, ... as in Notation 4.3 we see that $\text{HL} = \text{VR} = \text{Mix} = \emptyset$, due to the order of choosing the elements of the basis \mathcal{B} : For instance, a potential candidate to become a member of HL, would be first elected to Prime, and similarly VR is consumed by Bis and Mix by HR. Thus:

$$R(W) = \dim(V_{Seg}) = \mathbf{prime} + \mathbf{bis} + \mathbf{hr} + \mathbf{vl}.$$

Proposition 4.4(i) and (ii) implies

$$R(W') + R(W'') \leq \mathbf{prime} + \mathbf{vl} + \mathbf{bis} + \mathbf{hr} = R(W),$$

while $R(W') + R(W'') \geq R(W)$ always holds. This shows the desired additivity. \square

COROLLARY 4.6. *Assume that the additivity fails for W' and W'' , that is, $d = R(W') + R(W'') - R(W' \oplus W'') > 0$. Then the following inequalities hold:*

- (a) **mix** $\geq d \geq 1$,
- (b) **hl** + **hr** + **mix** $\geq \mathbf{e}' + \mathbf{e}'' + d \geq 3$,
- (c) **vl** + **vr** + **mix** $\geq \mathbf{f}' + \mathbf{f}'' + d \geq 3$.

Proof. To prove (a) consider the inequalities (i) and (ii) from Proposition 4.4 and their sum:

$$(4.7) \quad \begin{aligned} \mathbf{prime} + \mathbf{hl} + \mathbf{vl} + \mathbf{mix} &\geq R(W'), \\ \mathbf{bis} + \mathbf{hr} + \mathbf{vr} + \mathbf{mix} &\geq R(W''), \\ \mathbf{prime} + \mathbf{bis} + \mathbf{hl} + \mathbf{hr} + \mathbf{vl} + \mathbf{vr} + 2\mathbf{mix} &\geq R(W') + R(W''). \end{aligned}$$

The lefthand side of (4.7) is equal to $R(W) + \mathbf{mix}$, while its righthand side is $R(W) + d$. Thus the desired claim.

Similarly, using inequalities (iii) and (v) of the same proposition we obtain (b), while (iv) and (vi) imply (c). Note that $\mathbf{e}' + \mathbf{e}'' + d \geq 3$ and $\mathbf{f}' + \mathbf{f}'' + d \geq 3$ by Proposition 4.5. \square

4.2. Replete pairs. As we hunger after inequalities involving integers **prime**, \dots , **mix** we distinguish a class of pairs W', W'' with particularly nice properties.

DEFINITION 4.8. Consider a pair of linear spaces $W' \subset B' \otimes C'$ and $W'' \subset B'' \otimes C''$ with a fixed minimal decomposition $V = \langle V_{\text{Seg}} \rangle \subset B \otimes C$ and **Prime**, \dots , **Mix** as in Notation 4.3. We say (W', W'') is replete, if **Prime** $\subset W'$ and **Bis** $\subset W''$.

REMARK 4.9. Strictly speaking, the notion of replete pair depends also on the minimal decomposition V . But as always we consider a pair W' and W'' with a fixed decomposition $V = \langle V_{\text{Seg}} \rangle \supset W' \oplus W''$, so we refrain from mentioning V in the notation.

The first important observation is that as long as we look for pairs that fail to satisfy the additivity, we are free to replenish any pair. More precisely, for any fixed W', W'' (and V) define the *repletion* of (W', W'') as the pair $({}^{\Re}W', {}^{\Re}W'')$:

$$(4.10) \quad {}^{\Re}W' := W' + \langle \text{Prime} \rangle, \quad {}^{\Re}W'' := W'' + \langle \text{Bis} \rangle, \quad {}^{\Re}W := {}^{\Re}W' \oplus {}^{\Re}W''.$$

PROPOSITION 4.11. For any (W', W'') , with Notation 4.3, we have:

$$\begin{aligned} R(W') &\leq R({}^{\Re}W') \leq R(W') + (\dim {}^{\Re}W' - \dim W'), \\ R(W'') &\leq R({}^{\Re}W'') \leq R(W'') + (\dim {}^{\Re}W'' - \dim W''), \\ R({}^{\Re}W) &= R(W). \end{aligned}$$

In particular, if the additivity of the rank fails for (W', W'') , then it also fails for $({}^{\Re}W', {}^{\Re}W'')$. Moreover,

- (i) V is a minimal decomposition of ${}^{\Re}W$; in particular, the same distinguished basis $\text{Prime} \sqcup \text{Bis} \sqcup \dots \sqcup \text{Mix}$ works for both W and ${}^{\Re}W$.
- (ii) $({}^{\Re}W', {}^{\Re}W'')$ is a replete pair.
- (iii) The gaps $R({}^{\Re}W') - \dim({}^{\Re}W')$, $R({}^{\Re}W'') - \dim({}^{\Re}W'')$, and $R({}^{\Re}W) - \dim({}^{\Re}W)$, are at most (respectively) $R(W') - \dim(W')$, $R(W'') - \dim(W'')$, and $R(W) - \dim(W)$.

Proof. Since $W' \subset {}^{\Re}W'$, the inequality $R(W') \leq R({}^{\Re}W')$ is clear. Moreover ${}^{\Re}W'$ is spanned by W' and $(\dim {}^{\Re}W' - \dim W')$ additional matrices, that can be chosen out of **Prime** — in particular, these additional matrices are all of rank 1 and $R({}^{\Re}W') \leq R(W') + (\dim {}^{\Re}W' - \dim W')$. The inequalities about $''$ and $R(W) \leq R({}^{\Re}W)$ follow similarly.

Further ${}^{\Re}W \subset V$, thus V is a decomposition of ${}^{\Re}W$. Therefore also $R({}^{\Re}W) \leq \dim V = R(W)$, showing $R({}^{\Re}W) = R(W)$ and (i). Item (ii) follows from (i), while (iii) is a rephrasing of the initial inequalities. \square

Moreover, if one of the inequalities of Lemma 3.5 is an equality, then the respective W' or W'' is not affected by the repletion.

LEMMA 4.12. If, say, $R(W') + \mathbf{e}'' = R(W) - \dim W''$, then $W'' = {}^{\Re}W''$, and analogous statements hold for the other equalities coming from replacing \leq by $=$ in Lemma 3.5.

Proof. By Lemma 3.5 applied to ${}^{\Re}W = {}^{\Re}W' \oplus {}^{\Re}W''$ and by Proposition 4.11 we have:

$$\begin{aligned} R({}^{\Re}W) - \mathbf{e}'' &\stackrel{3.5}{\geq} R({}^{\Re}W') + \dim({}^{\Re}W'') \\ &\stackrel{4.11}{\geq} R(W') + \dim W'' \\ &\stackrel{\text{assumptions of 4.12}}{=} R(W) - \mathbf{e}'' \stackrel{4.11}{=} R({}^{\Re}W) - \mathbf{e}''. \end{aligned}$$

Therefore all inequalities are in fact equalities. In particular, $\dim({}^{\Re}W'') = \dim W''$. The claim of the lemma follows from $W'' \subset {}^{\Re}W''$. \square

4.3. Digestion. For replete pairs it makes sense to consider the complement of **(Prime)** in W' , and of **(Bis)** in W'' .

DEFINITION 4.13. With Notation 4.3, suppose S' and S'' denote the following linear spaces:

$$\begin{aligned} S' &:= \langle \text{Bis} \sqcup \text{HL} \sqcup \text{HR} \sqcup \text{VL} \sqcup \text{VR} \sqcup \text{Mix} \rangle \cap W' && \text{(we omit Prime in the union) and} \\ S'' &:= \langle \text{Prime} \sqcup \text{HL} \sqcup \text{HR} \sqcup \text{VL} \sqcup \text{VR} \sqcup \text{Mix} \rangle \cap W'' && \text{(we omit Bis in the union).} \end{aligned}$$

We call the pair (S', S'') the digested version of (W', W'') .

LEMMA 4.14. *If (W', W'') is replete, then $W' = \langle \text{Prime} \rangle \oplus S'$ and $W'' = \langle \text{Bis} \rangle \oplus S''$.*

Proof. Both $\langle \text{Prime} \rangle$ and S' are contained in W' . The intersection $\langle \text{Prime} \rangle \cap S'$ is zero, since the seven sets Prime, Bis, HR, HL, VL, VR, Mix are disjoint and together they are linearly independent. Furthermore,

$$\text{codim}(S' \subset W') \leq \text{codim}(\langle \text{Bis} \sqcup \text{HR} \sqcup \text{HL} \sqcup \text{VL} \sqcup \text{VR} \sqcup \text{Mix} \rangle \subset V) = \mathbf{prime}.$$

Thus $\dim S' + \mathbf{prime} \geq \dim W'$, which concludes the proof of the first claim. The second claim is analogous. \square

These complements (S', S'') might replace the original replete pair (W', W'') : as we will show, if the additivity of the rank fails for (W', W'') , it also fails for (S', S'') . Moreover, (S', S'') is still replete, but it does not involve any Prime or Bis.

LEMMA 4.15. *Suppose (W', W'') is replete, define S' and S'' as above and set $S = S' \oplus S''$. Then*

- (i) $R(S) = R(W) - \mathbf{prime} - \mathbf{bis} = \mathbf{hl} + \mathbf{hr} + \mathbf{vl} + \mathbf{vr} + \mathbf{mix}$ and the space $\langle \text{HL}, \text{HR}, \text{VL}, \text{VR}, \text{Mix} \rangle$ determines a minimal decomposition of S . In particular, (S', S'') is replete and both spaces S' and S'' contain no rank one matrices.
- (ii) *If the additivity of the rank $R(S) = R(S') + R(S'')$ holds for S , then it also holds for W , that is, $R(W) = R(W') + R(W'')$.*

Proof. Since $W = S \oplus \langle \text{Prime}, \text{Bis} \rangle$, we must have $R(W) \leq R(S) + \mathbf{prime} + \mathbf{bis}$. On the other hand, $S \subset \langle \text{HL}, \text{HR}, \text{VL}, \text{VR}, \text{Mix} \rangle$, hence $R(S) \leq \mathbf{hl} + \mathbf{hr} + \mathbf{vl} + \mathbf{vr} + \mathbf{mix}$. These two claims show the equality for $R(S)$ in (i) and that $\langle \text{HL}, \text{HR}, \text{VL}, \text{VR}, \text{Mix} \rangle$ gives a minimal decomposition of S . Since there is no tensor of type Prime or Bis in this minimal decomposition, it follows that the pair (S', S'') is replete by definition. If, say, S' contained a rank one matrix, then by our choice of basis in Notation 4.3 it would be in the span of Prime, a contradiction.

Finally, if $R(S) = R(S') + R(S'')$, then:

$$\begin{aligned} R(W) &= R(S) + \mathbf{prime} + \mathbf{bis} \\ &= R(S') + \mathbf{prime} + R(S'') + \mathbf{bis} \geq R(W') + R(W''), \end{aligned}$$

showing the statement (ii) for W . \square

As a summary, in our search for examples of failure of the additivity of the rank, in the previous section we replaced a linear space $W = W' \oplus W''$ by its repletion ${}^{\mathfrak{R}}W = {}^{\mathfrak{R}}W' \oplus {}^{\mathfrak{R}}W''$, that is possibly larger. Here in turn, we replace ${}^{\mathfrak{R}}W$ by a smaller linear space $S = S' \oplus S''$. In fact, $\dim W' \geq \dim S'$ and $\dim W'' \geq \dim S''$, and also $R(S) \leq R(W)$ and $R(S') \leq R(W')$ etc. That is, changing W into S makes the corresponding tensors possibly “smaller”, but not larger. In addition, we gain more properties: S is replete and has no Prime’s or Bis’s in its minimal decomposition.

COROLLARY 4.16. *Suppose that $W = W' \oplus W''$ is as in Notation 3.2 and that \mathbf{e}'' and \mathbf{f}'' are as in Notation 3.4. If either:*

- (i) \mathbb{k} is an arbitrary field, $\mathbf{e}'' \leq 1$ and $\mathbf{f}'' \leq 1$, or
- (ii) \mathbb{k} is algebraically closed, $\mathbf{e}'' \leq 1$ and $\mathbf{f}'' \leq 2$,

then the additivity of the rank $R(W) = R(W') + R(W'')$ holds.

Proof. By Proposition 4.11 and Lemma 4.15, we can assume W is replete and equal to its digested version. But then (since $\text{Bis} = \emptyset$) we must have $W'' \subset E'' \otimes C'' + B'' \otimes F''$. In particular, W'' is, respectively, a (1, 1)-hook shaped space or a (1, 2)-hook shaped space. Then the claim follows from Proposition 3.13 or Proposition 3.18. \square

4.4. Additivity of the tensor rank for small tensors. We conclude our discussion of the additivity of the tensor rank with the following summarising results.

THEOREM 4.17. *Over an arbitrary base field \mathbb{k} assume $p' \in A' \otimes B' \otimes C'$ is any tensor, while $p'' \in A'' \otimes B'' \otimes C''$ is concise and $R(p'') \leq \mathbf{a}'' + 2$. Then the additivity of the rank holds:*

$$R(p' \oplus p'') = R(p') + R(p'').$$

The analogous statements with the roles of A replaced by B or C , or the roles of $'$ and $''$ swapped, hold as well.

Proof. Since p'' is concise, the corresponding vector subspace $W'' = p''((A'')^*)$ has dimension equal to \mathbf{a}'' . By Corollary 4.16(i) we may assume $\mathbf{e}'' \geq 2$ or $\mathbf{f}'' \geq 2$. Say, $\mathbf{e}'' \geq 2 \geq R(p'') - \dim W''$, then by Corollary 3.7 the additivity must hold. \square

THEOREM 4.18. *Suppose the base field is $\mathbb{k} = \mathbb{C}$ or $\mathbb{k} = \mathbb{R}$ (complex or real numbers) and assume $p' \in A' \otimes B' \otimes C'$ is any tensor, while $p'' \in A'' \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$ for an arbitrary vector space A'' . Then the additivity of the rank holds: $R(p' \oplus p'') = R(p') + R(p'')$.*

Proof. By the classical Ja'Ja'-Takche Theorem [21] (in the algebraically closed case also shown in Proposition 3.18), we can assume p'' is concise in $A'' \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$. But then by [33, Thm 5 and Thm 6] the rank of p'' is at most $\mathbf{a}'' + 2$ and the result follows from Theorem 4.17. \square

Note that in the proof above we exploit the results about maximal rank in $\mathbb{k}^{\mathbf{a}''} \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$. In [33] the authors assume that the base field is \mathbb{C} or \mathbb{R} . We are not aware of any similar results over other fields, with the unique exception of $\mathbf{a}'' = 3$, see the next proof for a discussion.

THEOREM 4.19. *Suppose the base field \mathbb{k} is such that:*

- *the maximal rank of a tensor in $\mathbb{k}^3 \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$ is at most 5.*

(For example \mathbb{k} is algebraically closed of characteristic $\neq 2$ or $\mathbb{k} = \mathbb{R}$). Furthermore assume $R(p'') \leq 6$. Then independently of p' , the additivity of the rank holds: $R(p' \oplus p'') = R(p') + R(p'')$.

Proof. Without loss of generality, we may assume p'' is concise in $A'' \otimes B'' \otimes C''$. As in the previous proof, if any of the dimensions $\dim A''$, $\dim B''$, $\dim C''$ is at most 2, then the claim follows from [21]. On the other hand, if any of the dimensions \mathbf{a}'' , \mathbf{b}'' , \mathbf{c}'' is at least 4, then the result follows from Theorem 4.17. The remaining case $\mathbf{a}'' = \mathbf{b}'' = \mathbf{c}'' = 3$ also follows from Theorem 4.17 by our assumption on the field \mathbb{k} .

The assumption is satisfied for $\mathbb{k} = \mathbb{R}, \mathbb{C}$ see [6, Thm 5.1] or [33, Thm 5]. In [6, top of p. 402] the authors say that their proof is also valid for any algebraically closed field of characteristic not equal to 2. They also provide the interesting history of this question and, furthermore, they show that the assumption about maximal rank in $\mathbb{k}^3 \otimes \mathbb{k}^3 \otimes \mathbb{k}^3$ fails for $\mathbb{k} = \mathbb{Z}_2$. \square

Assuming the base field is $\mathbb{k} = \mathbb{C}$, one of the smallest cases not covered by the above theorems would be the case of $p', p'' \in \mathbb{C}^4 \otimes \mathbb{C}^4 \otimes \mathbb{C}^3$. The generic rank (that is, the rank of a general tensor) in $\mathbb{C}^4 \otimes \mathbb{C}^4 \otimes \mathbb{C}^3$ is 6, moreover [4, p. 6] claims the maximal rank is 7 (see also [33, Prop. 2]).

EXAMPLE 4.20. *Suppose $A' = A'' = \mathbb{C}^4$ and either $B' = B'' = \mathbb{C}^4$ and $C' = C'' = \mathbb{C}^3$ or $B' = C'' = \mathbb{C}^4$ and $B'' = C' = \mathbb{C}^3$. Suppose both $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ are tensors of rank 7 and that the additivity of the rank fails for $p = p' \oplus p''$. Let W' and W'' be as in Notation 3.2, and $E', e', \text{etc.}$ be as in Notation 3.4. Then:*

- $R(p) = 13$,
- $e' = e'' = f' = f'' = 2$,
- *with Prime, hl, etc., as in Notation 4.3, we have Prime = Bis = \emptyset , and the following inequalities hold:*

if $\mathbf{b}'' = 4, \mathbf{c}'' = 3$	or	if $\mathbf{b}'' = 3, \mathbf{c}'' = 4$
$2 \leq \mathbf{hl} \leq 3$		$2 \leq \mathbf{hl} \leq 3$
$2 \leq \mathbf{hr} \leq 3$		$3 \leq \mathbf{hr} \leq 4$
$3 \leq \mathbf{vl} \leq 4$		$3 \leq \mathbf{vl} \leq 4$
$3 \leq \mathbf{vr} \leq 4$		$2 \leq \mathbf{vr} \leq 3$
$1 \leq \mathbf{mix} \leq 3$		$1 \leq \mathbf{mix} \leq 3$
$\mathbf{hl} + \mathbf{vl} \leq 6$		$\mathbf{hl} + \mathbf{vl} \leq 6$
$\mathbf{hr} + \mathbf{vr} \leq 6$		$\mathbf{hr} + \mathbf{vr} \leq 6$

Sketch of proof. For brevity we only argue in the case $\mathbf{b}'' = 4, \mathbf{c}'' = 3$, while the proof of $\mathbf{b}'' = 3, \mathbf{c}'' = 4$ is very similar. Both tensors $p', p'' \in \mathbb{C}^4 \otimes \mathbb{C}^4 \otimes \mathbb{C}^3$ must be concise, as otherwise either Theorem 4.18 or JaJa-Takche Theorem imply the additivity of the rank. By Corollary 3.7 we must have $e' \leq 2$, and similarly for f', e'', f'' . If one of them is strictly less than 2, then Corollary 4.16(ii) implies the additivity, a contradiction, thus $e' = e'' = f' = f'' = 2$.

By the failure of the additivity, we must have $R(W) \leq 13$, but Lemma 3.5 implies also $R(W) \geq 13$, showing that $R(p) = 13$.

If, say Prime $\neq \emptyset$, then the digested version (S', S'') of repletion of (W', W'') is also a counterexample to the additivity by Lemma 4.15(ii). If $S = S' \oplus S''$ has lower rank than W , then either S is not concise, contradicting Theorem 4.18 or S contradicts the above calculations of rank. Thus also $R(S) = 13$ and by Lemma 4.15(i) we must have **prime** = **bis** = 0. In fact, $S = W$.

Let $\widetilde{E}' \subset E'$ be the smallest linear subspace such that $\pi_{C''}(\text{HL}) \subset \widetilde{E}' \otimes C'$. Set $\widetilde{e}' = \dim \widetilde{E}'$. Since Prime = \emptyset , we must have

$$W' \subset \langle \pi_{C''}(\text{HL}), \pi_{B''}(\text{VL}), \pi_{B''} \pi_{C''}(\text{Mix}) \rangle \subset \widetilde{E}' \otimes C' + B' \otimes F'.$$

That is, W' is (\widetilde{e}', f') -hook shaped. Since $f' = 2$, Proposition 3.18 shows that $\mathbf{hl} \geq \widetilde{e}' \geq 2$. Similarly, **hr**, **vl**, **vr** are also at least 2. We also see that $\widetilde{E}' = E'$, that is, the elements of type HL are concise in E' .

Next, we show that **vl** $\neq 2$, which is perhaps the most interesting part of this example. For this purpose we consider the projection $\pi_{E' \oplus B''}: B \rightarrow B'/E'$. The related map $B \otimes C \rightarrow (B'/E') \otimes C$

(which by the standard abuse we also denote $\pi_{E' \oplus B''}$), kills all the rank one tensors of types HL, HR, VR and Mix, leaving only those of type VL alive. The image $\pi_{E' \oplus B''}(W) \subset (B'/E') \otimes F'$ has rank at most \mathbf{vl} and is concise (otherwise, either Proposition 3.18 shows the additivity or p' is not concise, a contradiction in both cases). Note that $(B'/E') \otimes F' \simeq \mathbb{C}^2 \otimes \mathbb{C}^2$ and there are only two (up to a change of basis) concise linear subspaces of $\mathbb{C}^2 \otimes \mathbb{C}^2$ which have rank at most 2. In both cases it is straightforward to verify that there exists $\beta' \in (B'/E')^* \subset (B')^*$ such that $\beta'(p) = \beta'(p') \in A' \otimes C'$ has rank 1. Then, by swapping the roles of A and B , the process of repletion and digestion (Lemma 4.15) leads to a smaller tensor which is also a counterexample to the additivity of the rank, again a contradiction. Thus $R(\pi_{E' \oplus B''}(W))$ must be at least 3 and consequently, $\mathbf{vl} \geq 3$. The same argument shows that $\mathbf{vr} \geq 3$.

Combining the inequalities obtained so far we also get:

$$\mathbf{mix} = 13 - (\mathbf{hl} + \mathbf{hr} + \mathbf{vl} + \mathbf{vr}) \leq 3.$$

The inequality $\mathbf{mix} \geq 1$ follows from Corollary 4.6(a), and it is left to show only the last two inequalities. To prove $\mathbf{hl} + \mathbf{vl} \leq 6$, we use Proposition 4.4(ii):

$$7 \leq \mathbf{hr} + \mathbf{vr} + \mathbf{mix} = R(W) - (\mathbf{hl} + \mathbf{vl}) = 13 - (\mathbf{hl} + \mathbf{vl}).$$

The last inequality follows from a similar argument. \square

5. Additivity of the tensor border rank. Throughout this section we will follow Notations 3.1 and 3.2. Moreover, we restrict to the base field $\mathbb{k} = \mathbb{C}$.

We turn our attention to the additivity of the border rank. That is, we ask for which tensors $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ the following equality holds:

$$\underline{R}(p' \oplus p'') = \underline{R}(p') + \underline{R}(p'').$$

Since the known counterexamples to the additivity are much smaller than in the case of the additivity of the tensor rank, our methods are more restricted to very small cases. We commence with the following elementary observation.

LEMMA 5.1. *Consider concise tensors $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ with $\underline{R}(p') \leq \mathbf{a}'$ and $\underline{R}(p'') \leq \mathbf{a}''$ (thus in fact $\underline{R}(p') = \mathbf{a}'$ and $\underline{R}(p'') = \mathbf{a}''$). Let $p = p' \oplus p''$. Then the additivity of the border rank holds $\underline{R}(p) = \underline{R}(p') + \underline{R}(p'')$.*

Proof. Since p' and p'' are concise, the linear maps $p': (A')^* \rightarrow B' \otimes C'$ and $p'': (A'')^* \rightarrow B'' \otimes C''$ are injective. Then also the map $p: A^* \rightarrow B \otimes C$ is injective and

$$\underline{R}(p) \geq \dim p(A^*) = \dim p'((A')^*) + \dim p''((A'')^*) = \underline{R}(p') + \underline{R}(p'').$$

The opposite inequality always holds. \square

COROLLARY 5.2. *Suppose both triples of integers $(\mathbf{a}', \mathbf{b}', \mathbf{c}')$ and $(\mathbf{a}'', \mathbf{b}'', \mathbf{c}'')$ fall into one of the following cases: $(a, b, 1)$, $(a, 1, c)$, $(a, b, 2)$ with $a \geq b \geq 2$, $(a, 2, c)$ with $a \geq c \geq 2$, (a, b, c) with $a \geq bc$. Then for any concise tensors $p' \in A' \otimes B' \otimes C'$ and $p'' \in A'' \otimes B'' \otimes C''$ the additivity of the border rank holds.*

Note that the list of triples in the corollary is a bit exaggerated, as some of these triples have no concise tensors. However, this phrasing is convenient for further applications and search for unsolved pairs of triples.

Proof. After removing the triples that do not admit any concise tensor the list reduces to: $(a, a, 1)$, $(a, 1, a)$, $(a, b, 2)$ (for $2 \leq b \leq a \leq 2b$), $(a, 2, c)$ (for $2 \leq c \leq a \leq 2c$), (bc, b, c) . We claim that in all these cases $\underline{R}(p') = \mathbf{a}'$ and $\underline{R}(p'') = \mathbf{a}''$. In fact:

- The claim is clear for $(a, 1, a)$, $(a, a, 1)$, and (bc, b, c) .
- For $(a, a, 2)$ and $(a, 2, a)$ the claim follows from the classification of such tensors, see the argument in the first paragraph of [7, §5.3].
- For $(a, b, 2)$ (with $2 \leq b < a \leq 2b$), and $(a, 2, c)$ (with $2 \leq c < a \leq 2c$), the claim follows from the previous case: any such concise tensor T has border rank at least a . But T is at the same time a (non-concise) tensor in a larger tensor space $\mathbb{C}^a \otimes \mathbb{C}^a \otimes \mathbb{C}^2$ or $\mathbb{C}^a \otimes \mathbb{C}^2 \otimes \mathbb{C}^a$. Thus by Lemma 2.7 the border rank of T is at most the generic (border) rank in this larger space, which is equal to a by the previous item.

Therefore we conclude using Lemma 5.1. \square

Theorem 1.4 claims that the additivity of the border rank holds for $\mathbf{a}, \mathbf{b}, \mathbf{c} \leq 4$. Most of the cases follow from Corollary 5.2, with the exception of $(3 + 1, 2 + 2, 2 + 2)$ and $(3 + 1, 3 + 1, 3 + 1)$, which are covered in Sections 5.2 and 5.3.

DEFINITION 5.3. *Assume $p, q \in A \otimes B \otimes C$ are two tensors. We say that p is more degenerate than q if $p \in \overline{GL(A)} \times \overline{GL(B)} \times \overline{GL(C)} \cdot q$.*

EXAMPLE 5.4. Any concise tensor in $\mathbb{C}^1 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ is more degenerate than any concise tensor in $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$.

EXAMPLE 5.5. Consider concise tensors in $\mathbb{C}^3 \times \mathbb{C}^2 \times \mathbb{C}^2$. According to [22, Table 10.3.1], there are two orbits of the action of $GL_3 \times GL_2 \times GL_2$ of such tensors, both orbits of border rank 3. One orbit is “generic”, the other is more degenerate. The latter is represented by:

$$p = a_1 \otimes b_1 \otimes c_1 + a_2 \otimes b_1 \otimes c_2 + a_3 \otimes b_2 \otimes c_1.$$

LEMMA 5.6. Suppose $p' \in A' \otimes B' \otimes C'$ is an arbitrary tensor and $p'', q'' \in A'' \otimes B'' \otimes C''$ are such that $\underline{R}(p'') = \underline{R}(q'')$ and p'' is more degenerate than q'' . If the additivity of the border rank holds for $p' \oplus p''$ then it also holds for $p' \oplus q''$.

Proof. Since p'' is more degenerate than q'' also $p' \oplus p''$ is more degenerate than $p' \oplus q''$. Thus

$$\underline{R}(p' \oplus q'') \geq \underline{R}(p' \oplus p'') = \underline{R}(p') + \underline{R}(p'') = \underline{R}(p') + \underline{R}(q''). \quad \square$$

5.1. Strassen's equations of secant varieties. Often as a criterion to determine whether a tensor is or is not of a given border rank, we exploit defining equations of the corresponding secant varieties. We review here one type of equations that is most important for the small cases we consider in this article.

First assume $\mathbf{b} = \mathbf{c}$ and consider the space of square matrices $B \otimes C$. Let $f_{\mathbf{b}} : (B \otimes C)^{\times 3} \rightarrow B \otimes C$ be the map of matrices defined as follows:

$$(5.7) \quad f_{\mathbf{b}}(x, y, z) = x \operatorname{adj}(y)z - z \operatorname{adj}(y)x,$$

where $\operatorname{adj}(y)$ denotes the adjoint matrix of y .

As in Section 2.4 write

$$p = \sum_{i=1}^{\mathbf{a}} a_i \otimes w_i,$$

where $w_1, \dots, w_{\mathbf{a}} \in W := p(A^*) \subset B \otimes C$ are $\mathbf{b} \times \mathbf{c}$ matrices and $\{a_1, \dots, a_{\mathbf{a}}\}$ is a basis of A .

PROPOSITION 5.8. Assume that $p \in A \otimes B \otimes C$.

- (i) [32] Suppose $\mathbf{a} = \mathbf{b} = \mathbf{c} = 3$. Then $\underline{R}(p) \leq 3$ if and only if $f_3(x, y, z) = \underline{0}$ for every $x, y, z \in W$.
- (ii) [23] Suppose $\mathbf{a} = \mathbf{b} = \mathbf{c}$ and $\underline{R}(p) \leq \mathbf{a}$. Then $f_{\mathbf{a}}(x, y, z) = \underline{0}$, for every $x, y, z \in W$.

See also [19, Thm 3.2].

We also recall Ottaviani's derivation of Strassen's equations ([25], see also [22, Sect. 3.8.1]) for secant varieties of three factor Segre embeddings.

Given a tensor $p : B^* \rightarrow A \otimes C$, consider the contraction operator

$$p_A^\wedge : A \otimes B^* \rightarrow \Lambda^2 A \otimes C,$$

obtained as composition of the map $\operatorname{Id}_A \otimes p : A \otimes B^* \rightarrow A^{\otimes 2} \otimes C$ with the natural projection $A^{\otimes 2} \otimes C \rightarrow \Lambda^2 A \otimes C$.

PROPOSITION 5.9 ([25, Theorem 4.1]). Assume $3 \leq \mathbf{a} \leq \mathbf{b}, \mathbf{c}$. If $\underline{R}(p) \leq r$, then $\operatorname{rk}(p_A^\wedge) \leq r(\mathbf{a} - 1)$.

If $\mathbf{a} = 3$, we can slice p as follows (cf. Section 2.4): $p = \sum_{i=1}^3 a_i \otimes w_i \in A \otimes B \otimes C$, with $w_i \in B \otimes C$. Then the matrix representation of p_A^\wedge in block matrices is the following $(\mathbf{b} + \mathbf{b} + \mathbf{b}, \mathbf{c} + \mathbf{c} + \mathbf{c})$ partitioned matrix

$$(5.10) \quad M_3(w_1, w_2, w_3) := \begin{pmatrix} \underline{0} & w_3 & -w_2 \\ -w_3 & \underline{0} & w_1 \\ w_2 & -w_1 & \underline{0} \end{pmatrix}.$$

PROPOSITION 5.11 ([22, Prop. 7.6.4.4]). If $\mathbf{a} = \mathbf{b} = \mathbf{c} = 3$, the degree nine equation

$$\det(p_A^\wedge) = 0$$

defines the variety $\sigma_4(\mathbb{P}A \times \mathbb{P}B \times \mathbb{P}C) \subset \mathbb{P}(A \otimes B \otimes C)$.

If $\mathbf{a} = 4$ and $p = \sum_{i=1}^4 a_i \otimes w_i \in A \otimes B \otimes C$, with $w_i \in B \otimes C$, then the matrix representation of p_A^\wedge in block matrices is the following $(4 \cdot \mathbf{b}, 6 \cdot \mathbf{c})$ partitioned matrix

$$(5.12) \quad M_4(w_1, w_2, w_3, w_4) := \begin{pmatrix} \underline{0} & w_3 & -w_2 & w_4 & \underline{0} & \underline{0} \\ -w_3 & \underline{0} & w_1 & \underline{0} & -w_4 & \underline{0} \\ w_2 & -w_1 & \underline{0} & \underline{0} & \underline{0} & w_4 \\ \underline{0} & \underline{0} & \underline{0} & -w_1 & w_2 & -w_3 \end{pmatrix}.$$

5.2. Case $(3 + 1, 2 + \mathbf{b}'', 2 + \mathbf{c}'')$. Assume $\mathbf{a}' = 3$, $\mathbf{b}' = \mathbf{c}' = 2$ and $\mathbf{a}'' = 1$.

PROPOSITION 5.13. *For any $p' \in \mathbb{C}^3 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ and $p'' \in \mathbb{C}^1 \otimes \mathbb{C}^{\mathbf{b}''} \otimes \mathbb{C}^{\mathbf{c}''}$ the additivity of the border rank holds.*

Proof. We can assume p'' is concise, so that $\underline{R}(p'') = \mathbf{b}'' = \mathbf{c}''$. Also if p' is not concise, then Corollary 5.2 shows the claim. So suppose p' is concise and thus $\underline{R}(p') = 3$.

We can write $p' = a_1 \otimes w'_1 + a_2 \otimes w'_2 + a_3 \otimes w'_3$ and $p'' = a_4 \otimes w''_4$, where w'_1, \dots, w'_3 are 2×2 matrices and w''_4 is an invertible $\mathbf{b}'' \times \mathbf{b}''$ matrix.

As for p' , by Example 5.5 and Lemma 5.6 we can choose the more degenerate tensor, which has the following normal form:

$$w'_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, w'_2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, w'_3 = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Write $p = \sum_{i=1}^4 a_i \otimes w_i$, where w_i are the following $(2 + \mathbf{b}'', 2 + \mathbf{b}'')$ partitioned matrices

$$w_i = \begin{pmatrix} w'_i & 0 \\ 0 & 0 \end{pmatrix}, i = 1, 2, 3, w_4 = \begin{pmatrix} 0 & 0 \\ 0 & w''_4 \end{pmatrix}.$$

We use the same notation as in Section 5.1. We claim that the matrix representing the contraction operator p'_4 , denoted by $M_4(w_1, w_2, w_3, w_4)$ as in (5.12), has rank $7 + 3\mathbf{b}''$. We conclude that $\underline{R}(p) \geq 3 + \mathbf{b}'' = \underline{R}(p') + \underline{R}(p'')$ by Proposition 5.9 showing the additivity.

In order to prove the claim, we observe that $M_4(w_1, w_2, w_3, w_4)$ can be transformed via permutations of rows and columns into the following $(6 + 3\mathbf{b}'' + 2 + \mathbf{b}'', 6 + 3\mathbf{b}'' + 2 + 2 + 2 + 3\mathbf{b}'')$ -partitioned matrix

$$\begin{pmatrix} M_3(w'_1, w'_2, w'_3) & 0 & 0 & 0 & 0 & 0 \\ 0 & N & 0 & 0 & 0 & 0 \\ 0 & 0 & -w'_1 & w'_2 & -w'_3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

where N is the following $3\mathbf{b}'' \times 3\mathbf{b}''$ matrix

$$N = \begin{pmatrix} w''_4 & 0 & 0 \\ 0 & -w''_4 & 0 \\ 0 & 0 & w''_4 \end{pmatrix}.$$

One can compute that the rank of $M_3(w'_1, w'_2, w'_3)$ equals 5. Moreover, since $\text{rk}(N) = 3\mathbf{b}''$ and $\text{rk}((-w'_1, w'_2, -w'_3)) = 2$, we conclude the proof of the claim. \square

5.3. Case $(3 + 1, 3 + \mathbf{b}'', 3 + \mathbf{c}'')$. Recall our usual setting: $p' \in A' \otimes B' \otimes C'$, $p'' \in A'' \otimes B'' \otimes C''$, $\mathbf{a}' := \dim A'$, etc. (Notation 3.2). In this subsection we are going to prove the following case of additivity of the border rank.

PROPOSITION 5.14. *The additivity of the border rank holds for $p' \oplus p''$ if $\mathbf{a}' = \mathbf{b}' = \mathbf{c}' = 3$, and p' is concise and $\mathbf{a}'' = 1$.*

Proof. By replacing B'' and C'' with smaller spaces we can assume p'' is also concise and in particular $\mathbf{b}'' = \mathbf{c}''$. If $\underline{R}(p') = 3$ then Lemma 5.1 implies the claim. On the other hand, by Terracini's Lemma, $\underline{R}(p') \leq 5$. Thus it is sufficient to treat the cases $\underline{R}(p') = 4$ and $\underline{R}(p') = 5$.

Let $\{a_1, a_2, a_3\}$ be a basis of A' and let $\{a_4\}$ be a basis of $A'' \simeq \mathbb{C}$. Write

$$(5.15) \quad p' = a_1 \otimes w'_1 + a_2 \otimes w'_2 + a_3 \otimes w'_3,$$

where $w'_1, w'_2, w'_3 \in W' := p'((A')^*) \subset B' \otimes C'$ are 3×3 matrices. Similarly, let

$$p = a_1 \otimes w_1 + a_2 \otimes w_2 + a_3 \otimes w_3 + a_4 \otimes w_4,$$

where $w_1, w_2, w_3, w_4 \in W := p(A^*) \subset B \otimes C$ are $(3 + \mathbf{b}'', 3 + \mathbf{b}'')$ partitioned matrices:

$$(5.16) \quad w_i = \begin{pmatrix} w'_i & 0 \\ 0 & 0 \end{pmatrix}, i = 1, 2, 3, \text{ and } w_4 = \begin{pmatrix} 0 & 0 \\ 0 & w''_4 \end{pmatrix}.$$

We now analyse the two cases $\underline{R}(p') = 4$ and $\underline{R}(p') = 5$ separately.

The additivity holds if the border rank of p' is equal to four. Assume by contradiction that $\underline{R}(p) \leq \mathbf{b}'' + 3 = \underline{R}(p') + \underline{R}(p'') - 1$. By Proposition 5.8(ii), we obtain the following equations: $f_{\mathbf{b}''+3}(x', y' + y'', z') = 0$, for every $x', y', z' \in W' = p'((A')^*)$ and $0 \neq y'' \in W'' = p''((A'')^*)$. We can see that $\text{adj}(y' + y'')$ is the following $(3 + \mathbf{b}'', 3 + \mathbf{b}'')$ partitioned matrix

$$\text{adj}(y' + y'') = \begin{pmatrix} \det(y'') \text{adj}(y') & 0 \\ 0 & \det(y') \text{adj}(y'') \end{pmatrix}.$$

Therefore we have

$$x' \text{adj}(y' + y'') z' = \begin{pmatrix} \det(y'') x' \text{adj}(y') z' & 0 \\ 0 & 0 \end{pmatrix}.$$

Since p'' is concise, $\det(y'') \neq 0$, and thus from the vanishing of $f_{\mathbf{b}''+3}(x', y' + y'', z')$ we also obtain that $f_3(x', y', z') = 0$. Therefore $\underline{R}(p') \leq 3$ by Proposition 5.8(i), a contradiction.

The additivity holds if the border rank of p' is equal to five. Consider the projection $\pi : A \otimes B \otimes C \rightarrow A' \otimes B \otimes C$ given by

$$\begin{aligned} a_i &\mapsto a_i, \quad i = 1, 2, 3 \\ a_4 &\mapsto a_1 + a_2 + a_3. \end{aligned}$$

Consider $\bar{p} := \pi(p) \in A' \otimes B \otimes C$ and write $\bar{p} = a_1 \otimes \bar{w}_1 + a_2 \otimes \bar{w}_2 + a_3 \otimes \bar{w}_3$, where, for $i = 1, 2, 3$, \bar{w}_i is the $(3 + \mathbf{b}'', 3 + \mathbf{b}'')$ partitioned matrix

$$\bar{w}_i = \begin{pmatrix} w'_i & 0 \\ 0 & w''_i \end{pmatrix}.$$

We claim that $\text{rk}(\bar{p}_{A'}) = 9 + 2\mathbf{b}''$. Indeed, by swapping both rows and columns of $M_3(\bar{w}_1, \bar{w}_2, \bar{w}_3)$ (see Equation 5.10) we obtain the following $(9 + 3\mathbf{b}'', 9 + 3\mathbf{b}'')$ partitioned matrix

$$\begin{pmatrix} p'_{A'} & 0 \\ 0 & M_3(w''_1, w''_2, w''_3) \end{pmatrix}.$$

Since $R(p') = 5$, the matrix $p'_{A'}$ has rank 9, by Proposition 5.11. Moreover $M_3(w''_1, w''_2, w''_3)$ has rank $2\mathbf{b}''$. Therefore, by Proposition 5.9, we obtain $R(\bar{p}) \geq 5 + \mathbf{b}''$. We conclude by observing that $R(p) \geq R(\bar{p})$. \square

This concludes the proof of Theorem 1.4, as all possible splittings $\mathbf{a} = \mathbf{a}' + \mathbf{a}'', \mathbf{b} = \mathbf{b}' + \mathbf{b}'', \mathbf{c} = \mathbf{c}' + \mathbf{c}''$ with $\mathbf{a}, \mathbf{b}, \mathbf{c} \leq 4$ are covered either by Corollary 5.2 or one of Propositions 5.13 or 5.14.

#	$(\mathbf{a}', \mathbf{b}', \mathbf{c}')$	$(\mathbf{a}'', \mathbf{b}'', \mathbf{c}'')$	$R(p')$	$R(p'')$
1.	3, 2, 2	2, 3, 2	3	3
2.	3, 3, 2	2, 2, 3	3	3
3.	3, 3, 3	2, 2, 2	4, 5	2
4.	4, 2, 2	1, 2, 2	4	2
5.	4, 2, 2	1, 3, 3	4	3
6.	4, 3, 2	1, 2, 2	4	2
7.	4, 3, 3	1, 1, 1	5	1
8.	4, 3, 3	1, 2, 2	5	2
9.	4, 4, 3	1, 1, 1	5, 6	1
10.	4, 4, 4	1, 1, 1	5, 6, 7	1

TABLE 5.1

The list of pairs of concise tensors and their border ranks that should be checked to determine the additivity of the border rank for $\mathbf{a}, \mathbf{b}, \mathbf{c} \leq 5$. This list contains all pairs of concise tensors not covered by Corollary 5.2, or Proposition 5.13, or Proposition 5.14, together with their possible border ranks, excluding the cases covered by Lemma 5.1. The maximal possible values of border ranks above have been obtained from [1, Sect. 4].

One could analyse the additivity for $\mathbf{a}, \mathbf{b}, \mathbf{c} \leq 5$ (so for the bound one more than in Theorem 1.4) by checking all 10 possible cases listed in Table 5.1. We conclude the article by solving also Case 3 from the table.

EXAMPLE 5.17. If $p' \in \mathbb{C}^3 \otimes \mathbb{C}^3 \otimes \mathbb{C}^3$ and $p'' \in \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ are both concise, then the additivity of the border rank holds for $p' \oplus p''$. Indeed, by Example 5.4 there exists $q'' \in \mathbb{C}^1 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ more degenerate than p'' , but of the same border rank. By Lemma 5.6 it is enough to prove the additivity for $p' \oplus q''$. This is provided by Proposition 5.14.

Acknowledgments. We are enormously grateful to Joseph Landsberg for introducing us to this topic and numerous discussions and explanations. We also thank Michael Forbes, Mateusz Michałek, Artie Prendergast-Smith, Zach Teitler, and Alan Thompson for reference suggestions and their valuable comments. We are also grateful the referees and the journal editors for their suggestions that have helped to improve the results and presentation.

The research on this project was spread across a wide period of time. It commenced once E. Postingshel was a postdoc at IMPAN in Warsaw (Poland, 2012-2013) under the project

“Secant varieties, computational complexity, and toric degenerations” realised within the Homing Plus programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund.

Also our collaboration in years 2014-2019 was possible during many meetings, in particular those that were related to special programmes, such as: the thematic semester “Algorithms and Complexity in Algebraic Geometry” at Simons Institute for the Theory of Computing (2014), the Polish Algebraic Geometry mini-Semester (miniPAGES, 2016), and the thematic semester Varieties: Arithmetic and Transformations (2018). The latter two events were supported by the grant 346300 for IMPAN from the Simons Foundation and the matching 2015-2019 Polish MNiSW fund. We are grateful to the participants of these semesters for numerous inspiring discussions and to the sponsors for supporting our participations.

We are grateful to Loughborough University for hosting our collaboration in May 2017 and to Copenhagen University for hosting us in January 2019.

In addition, J. Buczyński is supported by the Polish National Science Center project “Algebraic Geometry: Varieties and Structures”, 2013/08/A/ST1/00804, the scholarship “START” of the Foundation for Polish Science and a scholarship of Polish Ministry of Science.

E. Postinghel was supported by a grant of the Research Foundation - Flanders (FWO) between 2013-2016 and is supported by the EPSRC grant no. EP/S004130/1 from late 2018.

Finally, the paper is also a part of the activities of the AGATES research group.

REFERENCES

- [1] H. ABO, G. OTTAVIANI, AND C. PETERSON, *Induction for secant varieties of Segre varieties*, Trans. Amer. Math. Soc., 361 (2009), pp. 767–792, <https://doi.org/10.1090/S0002-9947-08-04725-9>, <http://dx.doi.org/10.1090/S0002-9947-08-04725-9>.
- [2] B. ÅDLANDSVIK, *Joins and higher secant varieties*, Math. Scand., 61 (1987), pp. 213–222, <https://doi.org/10.7146/math.scand.a-12200>, <https://doi.org/10.7146/math.scand.a-12200>.
- [3] B. ALEXEEV, M. A. FORBES, AND J. TSIMERMAN, *Tensor rank: some lower and upper bounds*, in 26th Annual IEEE Conference on Computational Complexity, IEEE Computer Soc., Los Alamitos, CA, 2011, pp. 283–291, <https://doi.org/10.1109/cc.2011.28>.
- [4] M. D. ATKINSON AND N. M. STEPHENS, *On the maximal multiplicative complexity of a family of bilinear forms*, Linear Algebra Appl., 27 (1979), pp. 1–8, [https://doi.org/10.1016/0024-3795\(79\)90026-0](https://doi.org/10.1016/0024-3795(79)90026-0), [https://doi.org/10.1016/0024-3795\(79\)90026-0](https://doi.org/10.1016/0024-3795(79)90026-0).
- [5] A. BERELE AND A. REGEV, *Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras*, Adv. in Math., 64 (1987), pp. 118–175, [https://doi.org/10.1016/0001-8708\(87\)90007-7](https://doi.org/10.1016/0001-8708(87)90007-7), [https://doi.org/10.1016/0001-8708\(87\)90007-7](https://doi.org/10.1016/0001-8708(87)90007-7).
- [6] M. R. BREMNER AND J. HU, *On Kruskal’s theorem that every $3 \times 3 \times 3$ array has rank at most 5*, Linear Algebra Appl., 439 (2013), pp. 401–421, <https://doi.org/10.1016/j.laa.2013.03.021>, <https://doi.org/10.1016/j.laa.2013.03.021>.
- [7] J. BUCZYŃSKI, K. HAN, M. MELLA, AND Z. TEITLER, *On the locus of points of high rank*, Eur. J. Math., 4 (2018), pp. 113–136, <https://doi.org/10.1007/s40879-017-0172-2>.
- [8] J. BUCZYŃSKI AND J. JELISIEJEW, *Finite schemes and secant varieties over arbitrary characteristic*, Differential Geom. Appl., 55 (2017), pp. 13–67, <https://doi.org/10.1016/j.difgeo.2017.08.004>, <https://doi.org/10.1016/j.difgeo.2017.08.004>.
- [9] J. BUCZYŃSKI AND J. M. LANDSBERG, *Ranks of tensors and a generalization of secant varieties*, Linear Algebra Appl., 438 (2013), pp. 668–689, <https://doi.org/10.1016/j.laa.2012.05.001>, <http://dx.doi.org/10.1016/j.laa.2012.05.001>.
- [10] E. CARLINI, M. V. CATALISANO, AND L. CHIANTINI, *Progress on the symmetric Strassen conjecture*, J. Pure Appl. Algebra, 219 (2015), pp. 3149–3157, <https://doi.org/10.1016/j.jpaa.2014.10.006>, <https://doi.org/10.1016/j.jpaa.2014.10.006>.
- [11] E. CARLINI, M. V. CATALISANO, L. CHIANTINI, A. V. GERAMITA, AND Y. WOO, *Symmetric tensors: rank, Strassen’s conjecture and e-computability*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 18 (2018), pp. 363–390.
- [12] E. CARLINI, M. V. CATALISANO, AND A. ONETO, *Waring loci and the Strassen conjecture*. 07 2017, <https://doi.org/10.1016/j.aim.2017.05.008>.
- [13] E. CARLINI, N. GRIEVE, AND L. OEDING, *Four lectures on secant varieties*, in Connections between algebra, combinatorics, and geometry, vol. 76 of Springer Proc. Math. Stat., Springer, New York, 2014, pp. 101–146, https://doi.org/10.1007/978-1-4939-0626-0_2, https://doi.org/10.1007/978-1-4939-0626-0_2.
- [14] A. CASAROTTI, A. MASSARENTI, AND M. MELLA, *On Comon’s and Strassen’s conjectures*. arXiv:1810.09338.

- [15] L. CHIANTINI AND M. COPPENS, *Grassmannians of secant varieties*, Forum Math., 13 (2001), pp. 615–628, <https://doi.org/10.1515/form.2001.025>, <https://doi.org/10.1515/form.2001.025>.
- [16] C. CILIBERTO AND F. COOLS, *On Grassmann secant extremal varieties*, Adv. Geom., 8 (2008), pp. 377–386, <https://doi.org/10.1515/ADVGEOM.2008.025>, <https://doi.org/10.1515/ADVGEOM.2008.025>.
- [17] P. COMON, *Tensor decompositions: state of the art and applications*, in Mathematics in signal processing, V (Coventry, 2000), vol. 71 of Inst. Math. Appl. Conf. Ser. New Ser., Oxford Univ. Press, Oxford, 2002, pp. 1–24.
- [18] E. FEIG AND S. WINOGRAD, *On the direct sum conjecture*, Linear Algebra Appl., 63 (1984), pp. 193–219, [https://doi.org/10.1016/0024-3795\(84\)90143-5](https://doi.org/10.1016/0024-3795(84)90143-5), [http://dx.doi.org/10.1016/0024-3795\(84\)90143-5](http://dx.doi.org/10.1016/0024-3795(84)90143-5).
- [19] S. FRIEDLAND, *On tensors of border rank l in $\mathbb{C}^{m \times n \times l}$* , Linear Algebra Appl., 438 (2013), pp. 713–737, <https://doi.org/10.1016/j.laa.2011.05.013>, <http://dx.doi.org/10.1016/j.laa.2011.05.013>.
- [20] R. HARTSHORNE, *Algebraic geometry*, Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [21] J. JA'JA' AND J. TAKCHE, *On the validity of the direct sum conjecture*, SIAM J. Comput., 15 (1986), pp. 1004–1020, <https://doi.org/10.1137/0215071>, <http://dx.doi.org/10.1137/0215071>.
- [22] J. M. LANDSBERG, *Tensors: geometry and applications*, vol. 128 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2012, <https://doi.org/10.1090/gsm/128>.
- [23] J. M. LANDSBERG AND L. MANIVEL, *Generalizations of Strassen's equations for secant varieties of Segre varieties*, Comm. Algebra, 36 (2008), pp. 405–422, <https://doi.org/10.1080/00927870701715746>, <http://dx.doi.org/10.1080/00927870701715746>.
- [24] J. M. LANDSBERG AND M. MICHAŁEK, *Abelian tensors*, J. Math. Pures Appl. (9), 108 (2017), pp. 333–371, <https://doi.org/10.1016/j.matpur.2016.11.004>, <http://dx.doi.org/10.1016/j.matpur.2016.11.004>.
- [25] G. OTTAVIANI, *Symplectic bundles on the plane, secant varieties and Lüroth quartics revisited*, in Vector bundles and low codimensional subvarieties: state of the art and recent developments, vol. 21 of Quad. Mat., Dept. Math., Seconda Univ. Napoli, Caserta, 2007, pp. 315–352.
- [26] F. PALATINI, *Sulle superficie algebriche i cui $s_h(h+1)$ -seganti non riempiono lo spazio ambiente*, Atti della R. Acc. delle Scienze di Torino, 41 (1906), pp. 634–640.
- [27] F. RUPNIEWSKI, *Tensors of border rank 3 and Strassen's additivity conjecture*, master's thesis, Institute of Mathematics, Warsaw University, 2017. Supervisor: Jarosław Buczyński. Available at <https://www.mimuw.edu.pl/~jabu>.
- [28] A. SCHÖNHAGE, *Partial and total matrix multiplication*, SIAM J. Comput., 10 (1981), pp. 434–455, <https://doi.org/10.1137/0210032>.
- [29] Y. SHITOV, *A counterexample to Strassen's direct sum conjecture*. arXiv:1712.08660, 2017.
- [30] V. STRASSEN, *Gaussian elimination is not optimal*, Numer. Math., 13 (1969), pp. 354–356, <https://doi.org/10.1007/bf02165411>.
- [31] V. STRASSEN, *Vermeidung von Divisionen*, J. Reine Angew. Math., 264 (1973), pp. 184–202, <https://doi.org/10.1515/crll.1973.264.184>.
- [32] V. STRASSEN, *The asymptotic spectrum of tensors*, J. Reine Angew. Math., 384 (1988), pp. 102–152, <https://doi.org/10.1515/crll.1988.384.102>, <http://dx.doi.org/10.1515/crll.1988.384.102>.
- [33] T. SUMI, M. MIYAZAKI, AND T. SAKATA, *About the maximal rank of 3-tensors over the real and the complex number field*, Ann. Inst. Statist. Math., 62 (2010), pp. 807–822, <https://doi.org/10.1007/s10463-010-0294-5>, <https://doi.org/10.1007/s10463-010-0294-5>.
- [34] Z. TEITLER, *Sufficient conditions for Strassen's additivity conjecture*, Illinois J. Math., 59 (2015), pp. 1071–1085, <https://doi.org/10.1215/ijm/1488186021>, <http://projecteuclid.org/euclid.ijm/1488186021>.
- [35] F. L. ZAK, *Tangents and secants of algebraic varieties*, vol. 127 of Translations of Mathematical Monographs, American Mathematical Society, Providence, RI, 1993. Translated from the Russian manuscript by the author.